

7. Mulak A, Bonaz B. Brain-gut-microbiota axis in Parkinson's disease. *World J Gastroenterol* 2015;21(37):10609-10620.
8. Bedarf JR, Hildebrand F, Coelho LP, et al. Functional implications of microbial and viral gut metagenome changes in early stage L-DOPA-naïve Parkinson's disease patients. *Genome Med* 2017; 9(1):39.
9. Heintz-Buschart A, Pandey U, Wicke T, et al. The nasal and gut microbiome in Parkinson's disease and idiopathic rapid eye movement sleep behavior disorder. *Mov Disord* 2018;33(1): 88-98.
10. Hill-Burns EM, Debelius JW, Morton JT, et al. Parkinson's disease and Parkinson's disease medications have distinct signatures of the gut microbiome. *Mov Disord* 2017;32(5):739-749.
11. Keshavarzian A, Green SJ, Engen PA, et al. Colonic bacterial composition in Parkinson's disease. *Mov Disord* 2015;30(10): 1351-1360.
12. Minato T, Maeda T, Fujisawa Y, et al. Progression of Parkinson's disease is associated with gut dysbiosis: two-year follow-up study. *PLoS One* 2017;12(11):e0187307.
13. Petrov VA, Saltykova IV, Zhukova IA, et al. Analysis of gut microbiota in patients with Parkinson's disease. *Bull Exp Biol Med* 2017; 162(6):734-737.
14. Sampson TR, Debelius JW, Thron T, et al. Gut microbiota regulate motor deficits and neuroinflammation in a model of Parkinson's disease. *Cell* 2016;167(6):1469-1480. e12.
15. Scheperjans F. Gut microbiota, 1013 new pieces in the Parkinson's disease puzzle. *Curr Opin Neurol* 2016;29(6):773-780.
16. Scheperjans F, Aho V, Pereira PA, et al. Gut microbiota are related to Parkinson's disease and clinical phenotype. *Mov Disord* 2015; 30(3):350-358.
17. Gray MT, Munoz DG, Gray DA, et al. Alpha-synuclein in the appendiceal mucosa of neurologically intact subjects. *Mov Disord* 2014;29(8):991-998.
18. Svensson E, Horvath-Puho E, Stokholm MG, et al. Appendectomy and risk of Parkinson's disease: A nationwide cohort study with more than 10 years of follow-up. *Mov Disord* 2016;31(12): 1918-1922.
19. Marras C, Lang AE, Austin PC, et al. Appendectomy in mid and later life and risk of Parkinson's disease: A population-based study. *Mov Disord* 2016;31(8):1243-1247.
20. Palacios N, Fitzgerald K, Roberts AL, et al. A prospective analysis of airborne metal exposures and risk of Parkinson disease in the nurses' health study cohort. *Environ Health Perspect* 2014;122(9):933-938.
21. Palacios N, Fitzgerald KC, Hart JE, et al. Air pollution and risk of Parkinson's disease in a large prospective study of men. *Environ Health Perspect* 2017;125(8):0807011.
22. Mendes A, Goncalves A, Vila-Cha N, et al. Appendectomy may delay Parkinson's disease Onset. *Mov Disord* 2015;30(10): 1404-1407.
23. Houser MC, Chang J, Factor SA, et al. Stool immune profiles evince gastrointestinal inflammation in Parkinson's disease. *Mov Disord* 2018 [Epub ahead of print].
24. de Oliveira Machado SL, Bagatini MD, da Costa P, et al. Evaluation of mediators of oxidative stress and inflammation in patients with acute appendicitis. *Biomarkers* 2016;21(6):530-537.
25. Ferris M, Quan S, Kaplan BS, et al. The global incidence of appendicitis: a systematic review of population-based studies. *Ann Surg* 2017;266(2):237-241.

Supporting Data

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.

Functional Monoamine Oxidase B Gene Intron 13 Polymorphism Predicts Putaminal Dopamine Turnover in De Novo Parkinson's Disease

Matthias Löhle, MD,^{1,2*} Graziella Mangone, MD, PhD,³ Martin Wolz, MD,⁴ Bettina Beuthien-Baumann, MD,^{5,6,7} Liane Oehme, PhD,⁵ Jörg van den Hoff, PhD,⁶ Jörg Kotzerke, MD,⁵ Heinz Reichmann, MD,⁸ Jean-Christophe Corvol, PhD³ and Alexander Storch, MD^{1,2*}

¹Department of Neurology, University of Rostock, Rostock, Germany ²German Centre for Neurodegenerative Diseases (DZNE) Standort Rostock, Rostock, Germany ³Sorbonne Université, INSERM UMR51127 and CIC-1422, CNRS UMR7225, Assistance Publique Hôpitaux de Paris, ICM, Department of Neurology, Hôpital Pitié-Salpêtrière, Paris, France ⁴Department of Neurology, Elblandklinikum Meißen, Meissen, Germany ⁵Department of Nuclear Medicine, Technische Universität Dresden, Dresden, Germany ⁶Positron Emission Tomography Division, Helmholtz-Zentrum Dresden-Rossendorf, Dresden, Germany ⁷German Cancer Research Centre (DKFZ), Radiology, Heidelberg, Germany ⁸Department of Neurology, Technische Universität Dresden, Dresden, Germany

ABSTRACT: Objective: The objective of this study was to evaluate the effects of common functional polymorphisms in genes involved in dopamine metabolism on striatal dopamine turnover in de novo Parkinson's disease (PD).

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

***Corresponding authors:** Drs. Matthias Löhle and Alexander Storch, Department of Neurology, University of Rostock, Gehlsheimer Straße 20, 18147 Rostock, Germany; matthias.loehle@med.uni-rostock.de, alexander.storch@med.uni-rostock.de

Funding agencies: The initial randomized trial was supported by an unrestricted grant from Pfizer Pharma, Berlin, Germany. The research leading to these results has received additional funding from the program "Investissements d'Avenir" ANR-10-IAIHU-06. The financial sponsors of the study had no role in the study design, data collection, data analysis, data interpretation, or writing of the report.

Relevant conflicts of interests/financial disclosures: A.S. and H.R. have received unrestricted research grants, speaker honoraria, and consultancy fees from Pfizer Pharma GmbH, Berlin, Germany. The other authors have no other conflict of interest to report related to this article.

Received: 27 March 2018; **Revised:** 23 May 2018; **Accepted:** 25 May 2018

Published online 14 September 2018 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.27466

Methods: This was an observer-blinded cohort study investigating effects of common functional polymorphisms in dopa decarboxylase (*DDC*, rs921451), monoamine oxidase B (*MAOB*; rs1799836), catechol-O-methyltransferase (*COMT*, rs4680), and dopamine transporter/solute carrier family 6 member 3 (*DAT/SLC6A3*, variable number tandem repeats) genes on ^{18}F -fluorodopa uptake and an effective distribution volume ratio (inverse of dopamine turnover) measured by ^{18}F -fluorodopa PET in 28 untreated PD patients.

Results: Patients carrying the *MAOB*^{CC/CT} genotype (low/intermediate enzyme activity) had a lower dopamine turnover in the putamen (higher mean effective distribution volume ratio) when compared with patients with *MAOB*^{TT/TT} genotype (high enzyme activity). Striatal PET measures were not different between variants in the remaining genes.

Conclusions: The *MAOB* (rs1799836) polymorphism predicts putaminal dopamine turnover in early PD with the *MAOB*^{TT} allele linked to high enzyme activity leading to higher intrinsic dopamine turnover, which has been demonstrated to constitute a risk factor for motor complications. © 2018 The Authors. Movement Disorders published by Wiley Periodicals, Inc. on behalf of International Parkinson and Movement Disorder Society

Key Words: dopamine metabolism; positron emission tomography (PET); dopamine turnover; monoamine oxidase B (*MAOB*); functional gene polymorphisms; Parkinson's disease

Parkinson's disease (PD) is characterized by a progressive loss of dopaminergic neurons of the substantia nigra pars compacta projecting to the striatum. Motor symptoms only emerge after a substantial loss of striatal dopaminergic nerve terminals has already occurred,¹ which implies that a variety of compensatory mechanisms take place in the preclinical phase of PD.² One of these compensatory mechanisms is an increase of putaminal dopamine turnover, which can be measured with ^{18}F -fluorodopa PET even earlier than changes in dopamine synthesis and storage.^{3–6} Moreover, an extensively elevated dopamine turnover in early PD has been discussed as intrinsic risk factor for the development of motor complications.^{7,8} Conversely, mild elevation of putaminal dopamine turnover in de novo PD has been shown to be associated with a lower risk for motor complications.⁹

Although previous studies have demonstrated that age and levodopa treatment as known clinical risk factors for motor complications correlate with dopamine turnover in early PD,^{8,10} it remains unclear whether functional polymorphisms in genes involved in dopamine metabolism intrinsically contribute to individual changes of dopamine turnover. Thus, we aimed to investigate whether common functional polymorphisms in the dopa decarboxylase (*DDC*, rs921451),¹¹ monoamine oxidase B (*MAOB*, rs1799836),¹² catechol-O-methyltransferase

(*COMT*, rs4680)¹³ and dopamine transporter/solute carrier family 6 member 3 (*DAT/SLC6A3*, variable number tandem repeats)^{14–16} genes were predictive for striatal ^{18}F -fluorodopa uptake and dopamine turnover measured by extended ^{18}F -fluorodopa PET imaging^{3,10,17,18} in de novo PD.

Participants and Methods

This observer-blinded cohort study analyzed participants with diagnostic criteria defined PD from a previous randomized clinical trial^{10,19} who had received extended ^{18}F -fluorodopa PET imaging prior to any antiparkinsonian treatment. Participants underwent genetic testing of selected functional polymorphisms during subsequent follow-up. We included all remaining participants from the PET study for whom blood collection was possible. From the original cohort of 40 patients, 6 had died (2 from pulmonary embolism, 4 from unknown causes), 4 were lost to follow-up, and 2 had been diagnosed with essential tremor (both had normal putaminal K_{occ} values).

Motor symptoms and activities of daily living were estimated using the Unified PD Rating Scale (UPDRS),²⁰ disease stage was assessed with the Hoehn and Yahr scale.²¹ Clinical evaluations were carried out in drug-naïve patients by movement disorder trained physicians blinded to PET data (M.L., M.W.). Clinical and genetic testing had approval from the ethics committee at Technische Universität Dresden, Dresden, Germany (EK91052003), the clinical trial was registered with ClinicalTrials.gov (NCT00153972). Written informed consent was obtained from all participants.

PET measurements were carried out in drug-naïve participants using a 3D ECAT EXACT HR + system (Siemens/CTI, Knoxville, Tennessee) as described previously^{3,10} (refer to Supplementary Methods for details). In short, we applied a tissue reference model in which the activity curve of the occipital cortex served as input function for calculation of the tissue input uptake rate constant K_{occ} and effective distribution volume ratio (EDVR).^{17,18} K_{occ} and (EDVR) from the left and right sides were averaged for each scan.¹⁰

Genotyping of the common functional polymorphisms *DDC* (rs921451),¹¹ *MAOB* (rs1799836),¹² *COMT* (rs4680),¹³ and *DAT/SLC6A3* (variable number tandem repeats, VNTR)^{14–16} was performed as described previously (refer to Supplementary Methods for details). In brief, genomic DNA was extracted by standard methods from venous blood samples and polymorphisms were analyzed by an allelic discrimination Taqman assay (Applied Biosystems, Foster City, California). Positive control DNA template and negative control (DNA/RNA-free water) were included in each genotyping panel. *DAT*-VNTR polymorphism was determined by standard polymerase chain reaction




FIG. 1. Effects of monoamine oxidase B (MAOB) rs1799836 and catechol-O-methyl transferase (COMT) rs4680 genotypes on putaminal ¹⁸F-fluorodopa-PET measures in de novo Parkinson's disease patients. The boxplots show putaminal ¹⁸F-fluorodopa-PET uptake (A,C) and putaminal effective distribution volume ratio (EDVR) as the inverse of the dopamine turnover dopamine (B,D) with respect to the MAOB rs1799836 genotype (A,B) and the COMT rs4680 genotype (C,D). In the plots, boxes represent the interquartile range, horizontal lines the medians, and the antennas the range excluding outside values (defined as values beyond lower/upper quartile ± 1.5 times interquartile range). * $P < .05$ and *** $P < .001$ when compared with MAOB^{C(C)/CT} (unpaired 2-sided t -test). [Color figure can be viewed at wileyonlinelibrary.com]

(PCR) followed by agarose gel electrophoresis using the KAPA 2G Fast ReadyMix PCR Kit (KK5101, Roche, Basel, Switzerland). To investigate effects of gene polymorphisms on PET measures, patients were dichotomized into low/intermediate enzyme/transporter activity (DDC^{CC/CT}, MAOB^{CC/(C)/CT}, COMT^{AA/AG}, DAT^{≤9/≤10}) and high enzyme/transporter activity (DDC^{TT}, MAOB^{TT/(T)}, COMT^{GG}, DAT^{10/10}; Supplementary Table S1).^{11–16}

Statistical analyses were performed with IBM SPSS statistics, version 23.0 (IBM Corporation, Armonk, New York). Comparisons of demographic, clinical, genetic, and PET data between the groups were made with unpaired t -test or Mann–Whitney U test, or χ^2 test, as appropriate. To account for potentially confounding factors on the comparison of PET measures, we additionally performed an analysis of covariance

(ANCOVA), in which age, UPDRS part III score, and symptom duration were entered as continuous covariates and gender as fixed factor. If not mentioned otherwise, data are displayed as means \pm standard deviation or numbers and percentages. The significance level (2-tailed) was set at $P < .0125$ to correct for multiple comparisons using the Bonferroni method.

Results

Our cohort included 28 patients (19 [68%] men, 9 [32%] women; mean age 60.3 ± 9.4 years), who had been diagnosed 1.8 ± 2.0 years after the onset of motor symptoms. Mean age at disease onset was 57.3 ± 9.2 years, and mean UPDRS motor score was 19.4 ± 7.6 . Genotyping revealed allele frequencies

TABLE 1. ^{18}F -fluorodopa-PET measures in de novo Parkinson's disease patients with respect to MAOB rs1799836 genotypes

	Overall cohort	MAOB ^{CC(C)/CT}	MAOB ^{TT(T)}	P ^a	F and P values from ANCOVA analysis ^b
Participants, n	28	15	13		
K _{occ} (1/min), ^{18}F -fluorodopa uptake					
Caudate nucleus	0.0162 ± 0.0024	0.0170 ± 0.0028	0.0154 ± 0.0018	.091	F = 0.781, P = .387
Putamen	0.0117 ± 0.0019	0.0124 ± 0.0020	0.0108 ± 0.0015	.023	F = 4.450, P = .047
Anterior putamen	0.0154 ± 0.0025	0.0163 ± 0.0027	0.0145 ± 0.0020	.056	F = 2.969, P = .100
Posterior putamen	0.0081 ± 0.0017	0.0084 ± 0.0017	0.0078 ± 0.0016	.341	F = 0.447, P = .511
EDVR, inverse of DA turnover					
Caudate nucleus	2.00 ± 0.38	2.09 ± 0.34	1.87 ± 0.39	.113	F = 0.781, P = .387
Putamen	1.38 ± 0.25	1.52 ± 0.23	1.20 ± 0.14	<.001	F = 13.863, P = .001
Anterior putamen	1.86 ± 0.38	2.04 ± 0.37	1.66 ± 0.30	.006	F = 5.963, P = .024
Posterior putamen	0.98 ± 0.20	1.08 ± 0.22	0.86 ± 0.10	.003	F = 8.762, P = .007

Data are means ± standard deviations. ANCOVA, analysis of covariance; EDVR, Effective distribution volume ratio from ^{18}F -fluorodopa positron emission tomography; MAOB, monoamine oxidase B. K_{occ}, ^{18}F -fluorodopa uptake from positron emission tomography.

^a P values are from unpaired 2-sided t-tests comparing MAOB^{CC(C)/CT} and MAOB^{TT(T)} alleles. P values of < .0125 (2-sided) were considered significant to correct for multiple comparisons using the Bonferroni method and have been marked in bold.

^b F and P values are from ANCOVA comparing MAOB^{CC(C)/CT} and MAOB^{TT(T)} alleles and controlling for the candidate covariates age, UPDRS part III motor score as a measure of disease severity and symptom duration as well as gender as fixed factor. P values of < .0125 (2-sided) were considered significant to correct for multiple comparisons using the Bonferroni method and have been marked in bold.

similar to those reported in databases for all polymorphisms tested (Supplementary Table S1). All genotypes were in the Hardy-Weinberg equilibrium. We did not detect significant differences of demographic and clinical variables after stratification for genotypes in the selected gene variants (Supplementary Table S2).

Patients carrying the MAOB^{CC(C)/CT} (rs1799836) genotype, which is associated with low/intermediate enzyme activity in the brain,¹² had a higher EDVR (lower dopamine turnover) in the whole putamen compared to patients carrying the MAOB^{TT(T)} genotype that has been linked to high enzyme activity (Fig. 1, Table 1). We also observed slightly higher ^{18}F -fluorodopa uptake (K_{occ}) in the MAOB^{CC(C)/CT} (rs1799836) genotype, which was not significant after correction for multiple comparisons. No differences were detected between genotypes of the remaining genes (Fig. 1, Supplementary Tables S3-S5). The difference in EDVR between MAOB (rs1799836) genotype groups was confirmed in separated analysis of anterior and posterior portions of the putamen with similar distribution between the 2 putaminal regions. In contrast, we found no differences for K_{occ} and EDVR between groups in the caudate nucleus for all genotypes (Table 1).

Because PET measures had been reported to potentially correlate with age, gender, and disease severity,^{8,9} we additionally calculated ANCOVAs to control for the candidate covariates age, UPDRS motor score, symptom duration, and gender. Adjustment for these potential confounders largely confirmed the differences of putaminal EDVR between the MAOB (rs1799836) genotype groups (Table 1). No significant differences of PET measures were observed for the remaining genotype comparisons (Supplementary Tables S3-S5). Moreover, we did not identify significant interactions between genotype groups when calculating ancillary ANCOVAs for K_{occ} and EDVR with Bonferroni correction.

Discussion

We provide first evidence that a common functional variation in the MAOB gene leads to differences in putaminal dopamine metabolism as measured by extended ^{18}F -fluorodopa PET imaging in patients with early untreated PD. We found that the MAOB^{CC(C)/CT} rs1799836 genotype (associated with low/intermediate enzyme activity in the brain) resulted in lower putaminal dopamine turnover when compared with the MAOB^{TT(T)} genotype (high enzyme activity).

^{18}F -fluorodopa uptake (K_{occ} from standard PET scanning 10-60 minutes after tracer administration) is assumed to mainly reflect tracer influx and central DDC activity.²² Previous studies have indicated that common DDC polymorphisms (including rs921451) modulate the therapeutic response to levodopa without changing levodopa metabolism in the periphery, suggesting that DDC polymorphisms and resulting variations in DDC activity influence bioavailability of dopamine in the CNS.¹¹ Indeed, subsequent PET studies in healthy volunteers provided evidence that 1 of 4 common DDC haplotypes is predictive of ^{18}F -fluorodopa uptake in healthy humans.²³ Although this haplotype contains a major allele that is in linkage disequilibrium with rs921451,²³ we did not find any difference of ^{18}F -fluorodopa uptake with respect to DDC (rs921451) genotypes in our PD cohort. However, this haplotype is not uniquely defined by this single polymorphism.²³ Further studies are needed to assess the relationship between DDC genetics, central DDC activity, and levodopa response in PD.

In contrast to DDC, COMT and MAO-B as catabolizing enzymes in the dopamine metabolism are proposed to particularly influence dopamine turnover as measured by extended ^{18}F -fluorodopa PET protocols.^{3,24-26} Accordingly, PD patients carrying the functional

COMT^{GG/GG} (rs4680) genotype associated with low enzyme activity¹³ showed lower late ¹⁸F-fluorodopa uptake (measure of dopamine turnover) when compared with *COMT*^{AA/AA} carriers in the frontal cortex but not the striatum.²⁴ In agreement, *COMT* (rs4680) genotypes in our cohort did not predict striatal dopamine turnover in early PD patients. This observation is in contrast to our findings on MAO-B as the other main dopamine catabolizing enzyme: in close agreement with biochemical analyses showing a correlation of *MAOB*^{CC/(C)/CT} rs1799836 genotype with low/intermediate enzyme activity in postmortem human brain tissue,¹² our study demonstrated that the *MAOB*^{CC/(C)/CT} genotype is associated with lower putaminal dopamine turnover. We also observed a nonsignificant increase of ¹⁸F-fluorodopa uptake with this genotype, which is in agreement with studies showing that *K_{occ}* is to some extent sensitive to dopamine turnover even if calculated in the first 90 minutes after tracer injection.¹⁸

Although previous ¹⁸F-fluorodopa PET studies reported that decreased DAT levels are directly associated with higher dopamine turnover,²⁷ we did not observe any effects of the *DAT/SLC6A3* (VNTR) genotype known to influence DAT expression^{14–16} on dopamine turnover in our cohort. The reasons for this discrepancy remain unclear but might be related to differences in DAT level estimation between studies (direct quantification by PET imaging by Sossi and colleagues,²⁷ only indirect assumption in our approach).

Although we present data on one of the largest PD cohorts with systematic EDVR measurement by PET imaging, our study has several limitations. First, the cohort size and consequently the genotype group sizes are relatively small, which generated the need for statistical adjustment to confounding variables and may limit generalizability to a less homogenous population. Second, our data were generated from white patients selected at one university-based movement disorder outpatient center and is thus primarily reflective for patients of similar background and setting. Third, although patient groups were clinically indistinguishable at the time of PET imaging, we cannot completely exclude that the compositions of the various genotype groups may relevantly differ, particularly because genotype group sizes in some cases were relatively small after dichotomization. On the contrary, statistical adjustment for major demographic and clinical covariates largely confirmed our unadjusted analyses.

Together, our study suggests that the functional *MAOB* gene intron 13 (rs1799836) polymorphism predicts putaminal dopamine metabolism in early untreated PD with the *MAOB*^{TT} allele linked to high enzyme activity leading to higher intrinsic dopamine turnover, which has been shown to be associated with a higher subsequent risk for motor complications.⁹ Future studies are warranted to investigate whether this

polymorphism modulates the risk or the time of onset of motor complications in PD and whether individual assessment of this gene polymorphism might be helpful for early risk stratification in PD patients. ■

Acknowledgments: The authors would like to thank all patients and their relatives for their willingness to participate in our study. Furthermore, we are grateful for help of the members of our movement disorders outpatient clinic at Technische Universität Dresden, the PET division of the Helmholtz-Zentrum Dresden-Rossendorf, and the Coordination Center for Clinical trials at Technische Universität Dresden (particularly Uta Schwanebeck and Xina Grähler).

References

- Bernheimer H, Birkmayer W, Hornykiewicz O, Jellinger K, Seitelberger F. Brain dopamine and the syndromes of Parkinson and Huntington. Clinical, morphological and neurochemical correlations. *J Neuro Sci* 1973;20:415-455.
- Calne DB, Zigmond MJ. Compensatory mechanisms in degenerative neurologic diseases. Insights from parkinsonism. *Arch Neurol* 1991; 48:361-363.
- Oehme L, Perick M, Beuthien-Baumann B, et al. Comparison of dopamine turnover, dopamine influx constant and activity ratio of striatum and occipital brain with (1)(8)F-dopa brain PET in normal controls and patients with Parkinson's disease. *Eur J Nucl Med Mol Imaging* 2011;38:1550-1559.
- Sossi V, de La Fuente-Fernandez R, Holden JE, et al. Increase in dopamine turnover occurs early in Parkinson's disease: evidence from a new modeling approach to PET 18 F-fluorodopa data. *J Cereb Blood Flow Metab* 2002;22:232-239.
- Sossi V, de la Fuente-Fernandez R, Holden JE, Schulzer M, Ruth TJ, Stoessl J. Changes of dopamine turnover in the progression of Parkinson's disease as measured by positron emission tomography: their relation to disease-compensatory mechanisms. *J Cereb Blood Flow Metab* 2004;24:869-876.
- Sossi V, de la Fuente-Fernandez R, Nandhagopal R, et al. Dopamine turnover increases in asymptomatic LRRK2 mutations carriers. *Mov Disord* 2010;25:2717-2723.
- de la Fuente-Fernandez R, Schulzer M, Mak E, Calne DB, Stoessl AJ. Presynaptic mechanisms of motor fluctuations in Parkinson's disease: a probabilistic model. *Brain* 2004;127:888-899.
- Sossi V, de la Fuente-Fernandez R, Schulzer M, Adams J, Stoessl J. Age-related differences in levodopa dynamics in Parkinson's: implications for motor complications. *Brain* 2006;129:1050-1058.
- Löhle M, Mende J, Wolz M, et al. Putaminal dopamine turnover in de novo Parkinson disease predicts later motor complications. *Neurology* 2016;86:231-240.
- Storch A, Wolz M, Beuthien-Baumann B, et al. Effects of dopaminergic treatment on striatal dopamine turnover in de novo Parkinson disease. *Neurology* 2013;80:1754-1761.
- Devos D, Lejeune S, Cormier-Dequaire F, et al. Dopa-decarboxylase gene polymorphisms affect the motor response to L-dopa in Parkinson's disease. *Parkinsonism Relat Disord* 2014;20:170-175.
- Balciuniene J, Emilsson L, Orelund L, Pettersson U, Jazin E. Investigation of the functional effect of monoamine oxidase polymorphisms in human brain. *Hum Genet* 2002;110:1-7.
- Syvanen AC, Tilgmann C, Rinne J, Ulanen I. Genetic polymorphism of catechol-O-methyltransferase (COMT): correlation of genotype with individual variation of S-COMT activity and comparison of the allele frequencies in the normal population and parkinsonian patients in Finland. *Pharmacogenetics* 1997;7:65-71.
- Heinz A, Goldman D, Jones DW, et al. Genotype influences in vivo dopamine transporter availability in human striatum. *Neuropsychopharmacology* 2000;22:133-139.
- Mill J, Asherson P, Browes C, D'Souza U, Craig I. Expression of the dopamine transporter gene is regulated by the 3' UTR VNTR: evidence from brain and lymphocytes using quantitative RT-PCR. *Am J Med Genet* 2002;114:975-979.

16. VanNess SH, Owens MJ, Kilts CD. The variable number of tandem repeats element in DAT1 regulates in vitro dopamine transporter density. *BMC Genet* 2005;6:55.
17. Sossi V, Doudet DJ, Holden JE. A reversible tracer analysis approach to the study of effective dopamine turnover. *J Cereb Blood Flow Metab* 2001;21:469-476.
18. Sossi V, Holden JE, de la Fuente-Fernandez R, Ruth TJ, Stoessl AJ. Effect of dopamine loss and the metabolite 3-O-methyl-[18F] fluoro-dopa on the relation between the 18F-fluorodopa tissue input uptake rate constant K_{occ} and the [18F]fluorodopa plasma input uptake rate constant K_i. *J Cereb Blood Flow Metab* 2003;23: 301-309.
19. Hughes AJ, Daniel SE, Kilford L, Lees AJ. Accuracy of clinical diagnosis of idiopathic Parkinson's disease: a clinico-pathological study of 100 cases. *J Neurol Neurosurg Psychiatry* 1992;55: 181-184.
20. Fahn S, Elton R, UPDRS Development Committee. The Unified Parkinson's Disease Rating Scale. In: Fahn S, Marsden C, Calne D, eds. *Recent Developments in Parkinson's Disease*. 2nd ed. Florham Park, NJ: Macmillan Healthcare Information 1987;153-163, 293-304.
21. Hoehn MM, Yahr MD. Parkinsonism: onset, progression and mortality. *Neurology* 1967;17:427-442.
22. Patlak CS, Blasberg RG, Fenstermacher JD. Graphical evaluation of blood-to-brain transfer constants from multiple-time uptake data. *J Cereb Blood Flow Metab* 1983;3:1-7.
23. Eisenberg DP, Kohn PD, Hegarty CE, et al. Common variation in the DOPA decarboxylase (DDC) gene and human striatal DDC activity in vivo. *Neuropsychopharmacology* 2016;41:2303-2308.
24. Wu K, O'Keefe D, Politis M, et al. The catechol-O-methyltransferase Val(158)Met polymorphism modulates fronto-cortical dopamine turnover in early Parkinson's disease: a PET study. *Brain* 2012;135:2449-2457.
25. Ceravolo R, Piccini P, Bailey DL, Jorga KM, Bryson H, Brooks DJ. 18F-dopa PET evidence that tolcapone acts as a central COMT inhibitor in Parkinson's disease. *Synapse* 2002;43:201-207.
26. Ruottinen HM, Niinivirta M, Bergman J, et al. Detection of response to COMT inhibition in FDOPA PET in advanced Parkinson's disease requires prolonged imaging. *Synapse* 2001;40:19-26.
27. Sossi V, de la Fuente-Fernandez R, Schulzer M, Troiano AR, Ruth TJ, Stoessl AJ. Dopamine transporter relation to dopamine turnover in Parkinson's disease: a positron emission tomography study. *Ann Neurol* 2007;62:468-474.

Supporting Data

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.