



## Hypothesis

---

# Strengthening the Brain—Is Resistance Training with Blood Flow Restriction an Effective Strategy for Cognitive Improvement?

---

Alexander Törpel, Fabian Herold, Dennis Hamacher, Notger G. Müller and Lutz Schega

## Special Issue

The Effects of Exercise on Cognitive Function

Edited by

Dr. Paul D. Loprinzi





*Hypothesis*

# Strengthening the Brain—Is Resistance Training with Blood Flow Restriction an Effective Strategy for Cognitive Improvement?

Alexander Törpel <sup>1,\*</sup> , Fabian Herold <sup>2</sup> , Dennis Hamacher <sup>1</sup> , Notger G. Müller <sup>2,3,4</sup> and Lutz Schega <sup>1</sup>

<sup>1</sup> Institute III, Department of Sport Science, Otto von Guericke University Magdeburg, Zschokkestr. 32, 39104 Magdeburg, Germany; dennis.hamacher@ovgu.de (D.H.); lutz.schega@ovgu.de (L.S.)

<sup>2</sup> Research Group Neuroprotection, German Center for Neurodegenerative Diseases (DZNE), Leipziger Str. 44, 39120 Magdeburg, Germany; Fabian.Herold@dzne.de (F.H.); notger.mueller@dzne.de (N.G.M.)

<sup>3</sup> Center for Behavioral Brain Sciences (CBBS), Universitätsplatz 2, 39106 Magdeburg, Germany

<sup>4</sup> Department of Neurology, Medical Faculty, Otto von Guericke University, Leipziger Str. 44, 39120 Magdeburg, Germany

\* Correspondence: alexander.toerpel@ovgu.de; Tel.: +49-391-67-56600

Received: 11 September 2018; Accepted: 5 October 2018; Published: 9 October 2018



**Abstract:** Aging is accompanied by a decrease in physical capabilities (e.g., strength loss) and cognitive decline. The observed bidirectional relationship between physical activity and brain health suggests that physical activities could be beneficial to maintain and improve brain functioning (e.g., cognitive performance). However, the exercise type (e.g., resistance training, endurance training) and their exercise variables (e.g., load, duration, frequency) for an effective physical activity that optimally enhance cognitive performance are still unknown. There is growing evidence that resistance training induces substantial brain changes which contribute to improved cognitive functions. A relative new method in the field of resistance training is blood flow restriction training (BFR). While resistance training with BFR is widely studied in the context of muscular performance, this training strategy also induces an activation of signaling pathways associated with neuroplasticity and cognitive functions. Based on this, it seems reasonable to hypothesize that resistance training with BFR is a promising new strategy to boost the effectiveness of resistance training interventions regarding cognitive performance. To support our hypothesis, we provide rationales of possible adaptation processes induced by resistance training with BFR. Furthermore, we outline recommendations for future studies planning to investigate the effects of resistance training with BFR on cognition.

**Keywords:** cognition; strength training; blood flow restriction; neuroplasticity

## 1. Introduction

From the third decade of life, degenerative changes of the human organism increase which leads on the one hand to a reduced physical performance and on the other hand to a decline of cognitive functions. In terms of physical performance, especially the loss of muscle mass [1–4] contributes to a decrease in muscular strength which, in turn, impairs activities of daily living (e.g., walking) [5,6]. However, musculature is the main effector organ for developing muscular strength which is important to ensure motion respectively locomotion (e.g., walking safely) [7–9]. Therefore, the integrity of the musculature and the muscle strength is of great importance throughout the entire life span. Moreover, the mentioned age-related decreases in muscle mass and strength (because of aging) are also associated with morphological losses in the brain and decreased cognitive functions [10–13]. Because

of those changes, especially cognitive functions such as memory and processing speed are negatively affected [14–18]. Furthermore, aging-related changes of the brain are considered risk factors for the development of neurological diseases (e.g., dementia) [19,20]. Dementia is associated with cognitive impairments negatively affecting quality of life and independent living [19,21]. Based on the limited ability of individuals with neurological diseases (e.g., dementia) to live independently, an intensive medical care is needed which, in turn, consumes a large amount of resources of the welfare systems of industrialized nations [22–25].

So far, no pharmacological interventions are sufficient to treat the mentioned age-associated declines [26–30]. But, there is growing evidence with respect to positive effects of physical activity preventing and treating morphological and functional losses in muscles [31] and the brain [32,33]. In recent years, evidence has emerged emphasizing the existence of a bidirectional relationship between physical performance and brain health [34,35]. For instance, as mentioned above, a decrease in muscular performance is associated with a decrease in cognitive functioning [36–38]. Consequently, the bidirectional relationship suggests that physical training (means a structured, planned, dosed, and systematic form of physical activity with the focused aim to increase physical performance and/or health; e.g., through resistance training) may be a valuable intervention strategy to decelerate not only physical but also cognitive decline in old age. However, the exercise type (e.g., resistance training, endurance training) and exercise variables (e.g., load, duration, frequency), which would be optimal to efficiently enhance cognitive performance are largely unknown [39–49].

A promising and cost-effective physical intervention strategy [50–52] which preserves and enhances both, physical performance (especially with regard to the musculature) [53–60] and cognitive functions [61–65], is resistance training (also known as strength training). The underlying neurobiological mechanisms and effects of resistance training on cognition are described in the following section.

## 2. Effects and Mechanisms of Resistance Training on Cognition

The underlying neurobiological processes which are triggered by resistance exercises and have been related to cognitive performance improvements, are not fully understood, yet [61,65,66]. Based on the promising framework of Stillman et al. [67] about mediators of physical activity (in this case resistance exercises) influencing cognitive performance on different levels (cellular and molecular level, structural and functional level and behavioral/socioemotional level) [67], the current knowledge of possible neurobiological mechanisms contributing to the improvement of cognitive functions in response to resistance training are summarized in the following.

On the cellular and molecular level, a possible key mechanism of resistance training that contributes to cognitive improvements is the pronounced release of the multifaceted acting insulin-like growth factor 1 (IGF-1) [61,62,66,68–70]. In response to resistance training, IGF-1 is mainly expressed by the liver (global output, ~70% of total circulating IGF-1), musculature (local output) and the brain (local output) itself [71,72]. Circulating IGF-1 can cross the blood-brain barrier (BBB) which is therefore also available to the brain [71,72]. While an increased IGF-1 level is associated with proliferation, differentiation, survival, and migration of neuronal progenitors [73,74], synaptic processes (e.g., Long-Term Potentiation) [74,75], angiogenesis in the brain, neuroprotection, axon outgrowth, dendritic maturation, and synaptogenesis [72,76], a deficiency of IGF-1 is associated with the risk of harmful cerebrovascular events (e.g., ischemic stroke or impaired neurovascular coupling) [77,78]. Consequently, it is not surprising that a relationship between cognitive functions and IGF-1 level in older individuals [79] and in individuals with mild cognitive impairments was observed [80]. Furthermore, it is assumed that there is a potential relationship between diminished IGF-1 levels and neurodegenerative diseases [73,80,81], which suggests that influencing IGF-1 levels is a promising target for efficient treatments.

In fact, serum IGF-1 concentration levels are increased after a single bout of resistance activities (short-term) [82] and long-term (also known as “chronic”; >2 exercise sessions) resistance training in

humans [83,84]. However, currently there is only low evidence postulating a solid relationship between physical exercise-induced modulation of IGF-1 release and cognitive functions [85]. Nevertheless, one study reveals that basal changes of IGF-1 concentrations after a long-term resistance exercise intervention are associated with cognitive performance improvements [83]. Hence, further studies are needed to get a deeper understanding of the relationship of exercise-induced modulation of IGF-1 release and cognition [85].

On the structural level, Fontes et al. [86] observed that in older individuals, the grey matter density increases in the posterior and anterior lobe of the cerebellum, superior frontal gyrus in the frontal lobe and anterior cingulate cortex in the limbic lobe in response to a 12 weeks resistance training [86]. After a 6 months resistance exercise training program, an increase in cortical thickness in posterior cingulate cortex was observed which correlated with improvements in an overall cognition score [87]. Furthermore, in the study of Liu-Ambrose et al. [88], a reduced whole-brain volume after the end of 12 months resistance intervention as compared to control groups (balance and tone group) was noticed [88]. The reduced brain volume might be the consequence of dissolve degenerative changes of the brain such as amyloid plaques [46,88,89]. However, the distinct neuronal adaptations in response to resistance exercise interventions with different exercise variables suggest that a certain dose-response relationship between physical exercise variables and neural adaptations exists, although this dose-response relationship is currently poorly understood and has to be investigated in further studies [42,64,90–93].

In addition, long-term resistance training is associated with decreased white matter atrophy at follow-up measurements [94] and lower white matter lesions volume was observed after 52 weeks of a resistance training exercise regime [95]. White matter changes are known to influence cognitive performance especially in processing-speed-dependent cognitive tasks [96–99].

On the functional level, changes can be quantified either by measuring the activity of brain regions (for instance with electroencephalography [EEG], functional near-infrared spectroscopy [fNIRS], or functional magnetic resonance imaging [fMRI]) and/or by testing cognitive functions. Both, brain activity and cognitive functions were investigated after short-term and long-term resistance training to identify beneficial effects of this type of exercise on brain as well as cognitive performance [64]. In response to an acute bout of moderate-load [100] and high-load resistance training, an improvement in cognitive functions (higher number of solved items and lower reaction times in neutral Stroop task condition compared to non-exercising control group) and a decrease in the tissue oxygenation index in left and right prefrontal cortex was observed [101]. In the same manner, it has been shown that resistance training lasting several months can lead to a substantial increase in cognitive functions [62–64,83,88,94,102,103]. Furthermore, after a long-term resistance training intervention, a decreased cortical activation in prefrontal areas (lower concentration of oxygenated hemoglobin and total hemoglobin index values measured by fNIRS) during a standardized cognitive test (e.g., Stroop-test) was noticed [104]. A decreased activation in prefrontal areas and a simultaneous increase in cognitive functions may point towards a higher automatization in behavioral tasks or the redistribution of resources in other task-relevant cortical areas. The notion that higher levels of strengths are beneficial for cognitive performance is further supported by numerous cross-sectional studies observing that an improvement of hand grip strength [38,105,106], quadriceps strength [37], leg power [107], or whole body muscle strength [36] are linked to higher cognitive performance. Regarding the longitudinal and cross-sectional studies, the question arises whether (baseline) strength level per se [108] or adaptation processes evoked by regularly conducted resistance training (see above mentioned adaptations on cellular, molecular and structural level) are more beneficial for cognitive performance. Based on the current available scientific literature, we cannot unequivocally answer this question. As shown, there is evidence for both approaches (baseline strength vs. adaptation processes evoked by regularly conducted resistance training). But maybe just the combination of both has positive effects on cognitive functions.


On the behavioral/socioemotional level, the improvements in cognitive functions (e.g., executive functions) and the reduced activity of the prefrontal cortex are, for instance, linked to the functioning of the motor control of activities of daily living such as walking safely [109–113]. This phenomenon underpins the need to persevere the capacity of executive functions especially in older individuals in order to ensure mobility and independent living. Furthermore, because of the relationship between cognitive functions and quality of life [114], improvements in cognitive functions might be associated with an enhanced socioemotional status (e.g., decreased symptoms of depression and anxiety, increased joyful activities of daily living). Here, positive effects of resistance training on quality of life have been noticed [115].

However, concerning the effectiveness of the type of exercise, it was reported that resistance training is less effective than aerobic exercises regarding the improvement of cognitive performance on behavioral/socioemotional level [116] or on functional level regarding the task-related oxygenation of brain regions [101,104]. Nevertheless, there are several strategies to increase the effectiveness of resistance exercise regimes. A potential strategy which is likely to be beneficial to increase the efficiency of resistance training is the application of devices (e.g., cuffs) modulating the blood flow to and away from the muscles. This type of training is known as blood flow restriction training (BFR). So far, the higher effectiveness of resistance training with BFR compared to resistance training without BFR has only been investigated in the context of muscle physiological adaptations and strength improvements [117–119]. Whether resistance training with BFR provides also positive neurocognitive effects that are potentially greater than those effects observed after “traditional” resistance training interventions (resistance training without BFR) will be discussed in detail in the following section.

### 3. Resistance Training with Blood Flow Restriction—An Added Value for Cognition?

A way to increase the efficiency of resistance training is the specific manipulation of different exercise variables such as load, volume (repetitions, sets), rest periods, repetition velocity, choice of exercise, order of exercise, frequency or muscle action. [120]. Here, a certain dose-response relationship regarding certain exercise variables (e.g., load) can be observed [61,121,122]. Another, newer “manipulation strategy” to increase the efficiency of resistance training includes the application of hypoxic stimuli [123–126]. Hypoxic stimulation during resistance exercises could be achieved by applying (i) localized hypoxia or (ii) systemic hypoxia [125]. Localized hypoxia can be achieved with applying BFR which is in the literature also referred to as *occlusion training*. The training method BFR is characterized by the restriction / manipulation of the blood flow to and away from the limbs due to the application of elastic straps or inflatable pressure cuffs (e.g., blood pressure cuffs) to the proximal portion of the limbs (see Figure 1A,B) [117,125,127–130]. The manipulation of the blood flow especially decreases the venous return, which increases the accumulation of metabolites in the muscle triggering pronounced adaptational processes [117,125,127–130]. A particular type of BFR is KAATSU where special inflatable cuffs with pressure sensors are used [131]. Even though KAATSU is considered a type of BFR, this term is in a strict sense only applicable if KAATSU-equipment in BFR training is used. As consequence of the special construction of KAATSU-cuffs and their distinct application protocol, it is likely that differences between KAATSU and other BFR methods regarding the physiological stimuli occur. So far, these possible physiological differences between KAATSU and other BFR methods have not been directly and systematically compared. In this manuscript, the term BFR will therefore also include KAATSU training studies.

In general, systemic hypoxia is provided by breathing oxygen-reduced air [125]. Here, the oxygen-reduced air can be applied, for instance, with mask-system hypoxicators or via a stay in special rooms where the fraction of inspired oxygen is decreased (also known as normobaric altitude chambers) [132].



**Figure 1.** Schematic illustration of (A) the basic principles of blood flow restriction, (B) the application places of the cuffs for blood flow restriction and (C) the possible neurobiological mechanisms of resistance training with blood flow restriction that are likely to contribute to improved cognitive functions; blood flow restriction (BFR), growth hormone (GH), hypoxia-inducible factor (HIF), insulin-like growth factor 1 (IGF-1), resistance training (RE), vascular endothelial growth factor (VEGF).

Both, localized hypoxia (induced by BFR) and systemic hypoxia are harmless (when conducted appropriately) and well feasible [133–135]. However, due to the cuffs on the limbs during the BFR, petechial haemorrhage beneath the skin and/or numbness of extremities can appear in few cases [125,134,136]. Compared to localized hypoxia (e.g., BFR), systemic hypoxia has the advantage that it is not limited to the limbs [125]. Remarkably, cross-transfer effects in muscles that were not directly affected by the application of blood flow manipulation cuffs were observed in response to resistance exercises with BFR. Both, muscles proximal to the restricted extremities and muscles distal to the



restricted extremities experience beneficial effects [137,138]. Systemic-endocrinological (e.g., expression of growth factors) as well as neuronal adaptations (e.g., higher recruitment of supportive muscles because of the increased fatigued muscles under BFR) are discussed for this phenomenon. However, regarding brain adaptations, systemic hypoxia leads to an oxygen deficit directly in the brain which is to a certain extent the decisive stimulus triggering positive neurophysiological adaptations [135,139,140]. In this regard, first studies have shown improved cognitive functions following interventions with systemic, normobaric hypoxia [141,142].

Also, for resistance training with BFR, a first investigation by Sardeli et al. [143] had observed positive effects on cognitive functions (Stroop-test) immediately after a low-load resistance training with BFR (30% of 1RM) [143]. Except for this first investigation of Sardeli et al. [143], there are to our knowledge currently no further studies available (neither short-term nor long-term study) which directly examine the effects of localized hypoxic exposure on cognitive performance. Based on the first hint that localized hypoxia is beneficial for cognition, we want to outline several reasons why localized hypoxia during a resistance training (e.g., through BFR) might be a promising intervention strategy which is likely to increase the efficiency of resistance training regarding the enhancement of cognitive functions in the following:

(i) On the cellular and molecular level: Some investigations showed a significant higher release of hormones which is associated with positive neurophysiological adaptations, such as serum IGF-1 [144,145], growth hormone (GH) [146–149] and vascular endothelial growth factor (VEGF) [145,147,150,151], in response to acute resistance activities with BFR when compared to resistance training without BFR. Regarding the IGF-1, also a long-term intervention (two weeks) of low-intensity BFR training which was provided twice a day led to a higher basal level of IGF-1 in comparison to the same resistance training without BFR [152]. As mentioned in the previous section, IGF-1 plays an important role in synaptic functioning and cognitive processes [75]. Because of the link between a deficiency in serum GH level and a cognitive impairment, increases in GH are associated with benefits for cognitive performance [153,154]. Furthermore, in older adults who regularly perform physical exercises, a higher level of GH and better cognitive performance was noticed compared to sedentary older adults [155]. VEGF is involved in angiogenesis [39,156–161] and it is speculated that a decrease in angiogenic factors (e.g., serum VEGF) might be associated with cognitive impairments (e.g., in Alzheimer disease) [162,163]. Notably, the increases in neurochemical substances (e.g., IGF-1) was predominantly observed after an acute bout of resistance activities with BFR, thus long-term studies are needed to investigate whether a pronounced release of those neurochemicals would be persistent after longer time intervals (e.g., 6 months).

Furthermore, there is a robust body of evidence suggesting that the blood lactate concentrations are higher after an acute bout of resistance activities with BFR as compared to a resistance exercise without BFR [145,148,149,164–170]. The levels of post-exercise blood lactate concentration are associated with acute improvements in cognitive functions such as short-term memory [171] and executive functions [172,173]. This phenomenon occurs because peripherally expressed lactate can cross the BBB by monocarboxylate transporters (MCTs) and will be utilized as fuel for cognitive processes due to oxygenation [174–178]. Moreover, lactate is associated with changes in peripheral brain-derived neurotrophic factor (BDNF). Here, Ferris et al. [179] showed a correlation between blood lactate concentrations and BDNF [179]. Besides, Schiffer et al. [180] observed an increase in BDNF after a lactate infusion in rest [180]. These insights suggest a potential neurobiological relationship between both neurochemical substances. BDNF is a member of neurotrophins and contributes to neuroplasticity which, in turn, facilitates cognitive performance [181,182].

In addition, systemic hypoxia [183,184] as well as local hypoxia [185] increase the hypoxia-inducible factor 1 $\alpha$  (HIF-1 $\alpha$ ) which is the master regulator for adaptations of oxygen homeostasis. An increase of HIF-1 $\alpha$  in response to systemic and/or localized hypoxia (e.g., induced by BFR) might be meaningful for cognition or the integrity of the brain considering the following two aspects: Firstly, the HIF-1 $\alpha$  has a neuroprotective effect [186] and secondly, this transcription factor

triggers the increase of neurotrophic factors such as the VEGF and IGF-1 [187,188]. Therefore, the HIF-1 $\alpha$  may be also a crucial factor for neurocognitive adaptations following a resistance training with BFR.

(ii) On the functional level: After a resistance training with BFR, increases in the cortical excitability [189] and higher levels of oxygenated hemoglobin were observed in cortical motor areas (compared to same resistance exercises without BFR) [190]. Furthermore, in prefrontal areas, a higher concentration of deoxygenated hemoglobin was observed during knee extension with BFR whereas the increase in oxygenated hemoglobin was diminished when compared to knee extensions without BFR [191]. In general, decreased levels of deoxygenated hemoglobin and increased levels of oxygenated hemoglobin are associated with increased cortical activity [192–195]. Since deoxygenated hemoglobin is assumed to be less affected by physiological artefacts than oxygenated hemoglobin [192,196–201], it is perhaps a better proxy of cortical activity (in this particular case) and it could therefore be speculated that a pronounced decrease of deoxygenated hemoglobin may point towards a higher cortical activation during knee extensions with BFR. Nevertheless, further research is necessary to verify or falsify these assumptions.

In general, higher levels of cortical activity (e.g., shown by higher concentration of oxygenated hemoglobin in the brain) after physical exercises are associated with improved cognitive performance [202,203]. It was observed that participants with an improved cognitive performance after exercise showed a higher cortical activity in prefrontal areas during the exercise sessions (termed as responders) in comparison to participants with no cognitive improvements (termed as non-responders) [204]. In consideration of these insights, the enhanced performance in the Stroop test after a low-load resistance training with BFR observed in the investigation of Sardeli et al. [143] may have been caused by higher levels of oxygenated hemoglobin in the prefrontal cortex [143].

### 3.1. Hypothesis

According to the potential neurobiological advantages of a resistance training with BFR compared to a resistance training without BFR on cellular and molecular level as well as on functional level of the brain (see Figure 1C), we hypothesize that a short-term and long-term resistance training with BFR is more efficient regarding the enhancement of cognitive functions than a “traditional” resistance exercise regime without BFR.

### 3.2. Considerations to Evaluate the Hypothesis

To test the hypothesis stated in the previous section, there are a number of general aspects that should be considered regarding (i) the participants’ characteristics, (ii) designing the resistance training program and (iii) the outcome measures.

(i) Regarding the selection of participants, it should be considered that individual characteristics moderate the outcomes and underlying neurobiological processes. Exemplarily, sex is a key moderator for the effect of physical exercise interventions on cognitive performance which is perhaps related to underlying neurobiological processes [116,205,206]. Here, it is assumed that women may benefit more from exercise than men with respect to cognitive functions like executive functions [116]. While the reason for this sex-phenomenon is not fully understood, it is assumed that those findings are related to sex-dependent neurobiological mechanisms (e.g., exercise-induced release of BDNF) and the higher level of habitual physical inactivity in older women (compared to older men) [68,116,205,206]. Another moderator which potentially influences the exercise-cognition interaction is the genotype of the participant [68,116] and through matching the individuals’ genotypes to an appropriate resistance training program, a greater outcome regarding muscular strength can be evoked [207]. However, currently there is not enough evidence available which would allow validly designing resistance training regimes/programs with or without BFR as a function of individual genotypes. Hence, further investigations in this field are needed. Here, moderator and mediator variables should be carefully assessed and their influence on outcomes measures as well as neurobiological processes evaluated.



A deeper understanding of moderator and mediator variables would assist the development of more personalized training regimes which may provide greater intervention efficiency [68].

Additionally, further studies should consider and test the “human baseline hypothesis” which proposes that the baseline values of strength (e.g., grip strength and/or knee extensor strength) assessed prior to resistance training or after a detraining period are more appropriate markers of long-term health outcomes compared to training-related strength gains [108]. Therefore, in relation to brain-health gains (brain volume, cognitive functions), the baselines of strength as well as muscle mass should be taken into account.

(ii) For designing resistance training programs, in general, the following exercise variables should be considered [120,208,209]:

Variables of a resistance training session:

- (1) *load* (amount of weight that is used for an exercise; usually given as a percentage of the one repetition maximum [1RM]);
- (2) number of repetitions;
- (3) number of sets;
- (4) inter-set rest periods;
- (5) inter-exercise rest periods;
- (6) *number of exercises* (for the whole training session or for a muscle or a muscle group with the same function);
- (7) *repetition velocity* (temporal details should be given for: concentric phase–inter-repetition rest periods–eccentric phase rest period up to the start of the next repetition, e.g., 2–0–2–1 s);
- (8) *muscle action* (concentric, eccentric, isometric);
- (9) *exercise selection* (e.g., multi-joint or single joint exercises);
- (10) *exercise order* (e.g., squat, leg extension, biceps curl and concentration curl or squat, biceps curl, leg extension and concentration curl);
- (11) *volitional muscle failure*
- (12) *range of motion*.

Variables for structuring resistance training:

- (13) *frequency* (number of training sessions per week);
- (14) *density* (distribution of training sessions across a week with regard to recovery time in-between training sessions) and
- (15) *duration* (duration over which a training program is carried out before exercise variables are changed).

It should be noted that some exercise variables are usually summarized into variables with a different designation: e.g., volume (exercise variables 2, 3 and 6) or time under tension (TUT, sum of the exercise variables 2 and 7) [120,209]. Additionally, the cuff pressure is of particular importance in resistance exercises with BFR, as it is intended to induce an appropriate level of localized hypoxia as physiological stimuli [210–217]. Here, the cuff pressure should be applied in such a way that venous pooling without an arterial occlusion would occur. To achieve this, the cuff pressure must be below the arterial occlusion pressure [124]. However, various moderator variables can influence the cuff pressure:

- (1) *Cuff width*: wide BFR-cuffs restrict arterial blood flow more than narrow BFR-cuffs using the same cuff pressure. Therefore, the cuff pressure should be applied relative to the cuff width [214,215,218–222].
- (2) *Cuff material*: it might be that the cuff material has an impact on the arterial blood flow restriction [211]. However, current investigations comparing different cuff materials (5 cm nylon vs. 3 cm elastic cuffs) do not consider the cuff width [223]. In contrast, Loenneke et al. [224]

compared nylon and elastic cuffs with the same width (5 cm) and observed no differences in the arterial occlusion pressure [224].

- (3) *Restricted extremity* (upper or lower limbs): cuff pressures should be determined individually for the upper and lower limbs [225].
- (4) *Systolic / arterial blood pressure*: the cuff pressure depends on the systolic / arterial blood pressure [213,218,226–232].
- (5) *Body composition / anthropometry*: the circumference of the limbs is the biggest predictor for the cuff pressure to reach arterial blood flow restriction and should be considered [218,225,233–235].
- (6) *Body position*: the cuff pressure to reach arterial blood flow restriction must be lower in the supine position compared to the seated position and standing position [210,212].
- (7) *Exercise protocol*: applying intermittent or continuous pressure; it might be that a BFR applied with a continuous pressure on the cuffs during the exercise leads to another physiological stimulus as compared to a BFR applied in an intermittent fashion [124,226,236–238].
- (8) *Blood flow restriction system*: different blood flow restriction systems (automatic pressure control vs. manual handheld pressure control) lead to diverging pressure on the limbs at rest and during exercise. However, one first investigation by Hughes et al. [239] compared several blood flow restriction systems with different cuff widths. Therefore, the influence of blood flow restriction systems for inducing effective BFR-stimuli needs further investigations [239].

Since those mentioned moderator variables are crucial for an effective BFR-stimulus and the physiological response, as well as the psychological response, it is likely that those also alter neurocognitive adaptations, which, in turn, influence the changes in cognitive functions. To evoke the above mentioned cognition-related neurobiological adaptations through a resistance training with BFR, it is strongly recommended to determine a personalized cuff pressure be chosen [217,240] which takes the above mentioned relationships of the moderator variables and the cuff pressure into account. From a practitioner's view, the optimal solution(s) to determine the cuff pressure ensuring an appropriate stimulus would be using a pressure that is relative to the used cuffs and individual's characteristics [117,241] or to use a BFR system that automatically adjust the cuff pressure [239,240]. Furthermore, even moderate cuff pressures induce adaptations comparable to high cuff pressures [227,242]. Hence, moderate cuff pressures are recommended because higher cuff pressures increase the risk of full arterial occlusion and in turn of adverse effects [131,243,244].

In resistance training without BFR, only the following exercise variables are considered and recommended to enhance cognitive functions, by now (to 1.) load: 60 to 80% of 1RM; (to 2.) number of repetitions: 7; (to 3.) number of sets: 2; (to 4.) inter-set rest periods: 2 min; (to 13.) frequency: at least twice per week; (to 15.) duration of a training period 2 to 12 months [61]. However, in short-term and long-term resistance training interventions with (and even without) BFR, the optimal selection of exercise variables to efficiently enhance cognitive functions are largely unknown, and should be investigated in future studies. Nevertheless, we would like to recommend the following exercise variables for a resistance training with BFR aiming to induce neurocognitive adaptations (Table 1). We chose these exercise variables because of their effectiveness to increase muscular strength as well as muscular hypertrophy [124,130,137,144,216,245–248]. As described above, functional and structural adaptations of the musculature are moderating factors for the neurocognitive status. Furthermore, based on our above mentioned deliberations, it can be assumed that these exercise variables are efficient to trigger adaptations on the above mentioned neurocognitive levels (cellular, molecular, structural and functional level).

**Table 1.** Recommendations for exercises variables for a resistance training with blood flow restriction (BFR); n.a.: not available; reps: repetitions; 1RM: one repetition maximum; s: seconds; min: minute.

Exercise Variables	Recommendations for Resistance Training with BFR
(1.) load	20 to 50% of 1RM
(2.) number of repetitions	15 to 30 per set, 50 to 80 repetitions per exercise (e.g., 30–15–15–15 reps)
(3.) number of sets	3 to 5 sets per exercise
(4.) inter-set rest periods	30 to 60 s
(5.) inter-exercise rest periods	5 min (without BFR)
(6.) number of exercises	n.a.
(7.) repetition velocity	1 (to 2)–0–1 (to 2)–1 s
(8.) muscle action	dynamic muscle action, eccentric is more effective than concentric
(9.) exercise selection	single- and multi-joint exercise
(10.) exercise order	n.a., depending on the training goal
(11.) volitional muscle failure	until volitional fatigue/repetition failure/technical failure
(12.) range of motion	full range of motion
(13.) frequency	2 to 3 sessions per week
(14.) density	n.a., depending on the performance level
(15.) duration	n.a., but according to the general physiological view, exercise variables or exercises should be changed after a mesocycle of 8 to 12 weeks

Furthermore, so far, there have been no consistent recommendations for the cuff pressure. However, the following criteria are often used to apply an optimal cuff pressure: 130% of the systolic blood pressure [226,237]; 10 mm Hg below the arterial occlusion pressure [225]; ~50% arterial occlusion pressure [243]. The most effective cuff pressure has still to be identified [124].

In general, resistance training with BFR is a harmless treatment strategy when applied appropriately [117,133,136,222,241,249–251], but in order to minimize the occurrence and/or to avoid adverse health effects, safety recommendations should be considered [134] and available risk assessment tools should be used [252]. Furthermore, we want to point out that during the practical implementation of a resistance training with BFR, the following general safety recommendations should be strictly adhered to minimize the occurrence of adverse events: We strongly recommend that an individual and adequate cuff pressure should be applied. Furthermore, based on the currently available recommendations the maximal duration for continuous BFR should in general not exceed a time period of circa 10 to 15 min for the upper limbs and circa 15 to 20 min for the lower limbs because longer time periods may increase the risk of adverse events [134].

(iii) Physical exercises influence cognitive performance on multiple levels: (1) cellular and molecular level; (2) structural and functional level and (3) behavioral/socioemotional level [67]. Based on these mentioned levels, multiple outcome measures should be considered in the study design and analysis in order to understand the complex interaction of physical exercises (e.g., resistance training with BFR) and cognition:

(1) On the cellular and molecular level, neurochemical markers such as IGF-1, GH, VEGF, blood lactate concentrations and BDNF might be used since the exhibited associations with cognitive performance (see the previous sections).

(2) On the structural and functional level, different neuroimaging modalities such as fNIRS, EEG, fMRI or a combination of those should be applied in order to understand physical exercise induced structural and functional brain changes [49,253]. Since fNIRS and EEG can in particular be used during physical exercises [254–256], both measuring systems are suitable for the evaluation of cognitive activity while performing resistance training with BFR. Here, short-term and long-term effects of this intervention strategy could be objectified. Regarding functional brain changes, it seems recommendable (a) to use standardized and established cognitive test (e.g., Stroop test [101,104], Sternberg test [257–259], Eriksen Flanker test [102]) to ensure comparability with existing studies and (b) to consider attention and perceptual tasks which were currently not in the focus of exercise-cognition research [260] but could be important for special cohorts (e.g., individuals with dementia) [48].

(3) On the behavioral/socioemotional level, established questionnaires such as “Felt Arousal Scale” [261], “Ratings of perceived exertion” [262], “Visual Analogue Scales” (e.g., to assess motivation or mental fatigue) [172,263–265], “SF-36” (to assess physical and mental components of the quality of life) [266] and “Pittsburgh Sleep Quality Index” (to assess various components of the sleep quality) [267] which are widely used in exercise–research with a neuropsychological and behavioral/socioemotional focus [141,172,263–265,268,269], should be used to elucidate the (moderating) effects of socioemotional states.

#### 4. Conclusions

The type of physical exercise (e.g., resistance training) in combination with related exercise variables (e.g., load, number of repetitions and sets) which efficiently enhance cognitive performance are largely unknown [39–49]. A promising physical exercise intervention which fends off physical and cognitive decline (e.g., due to the aging process) is resistance training. Hypothetically, the efficiency of resistance training interventions on cognition could be increased due to the application of BFR.

Resistance training with BFR is more efficient to increase muscle hypertrophy and strength as compared to the same resistance training without BFR [247,270] and for a resistance training with BFR, lower exercise loads are needed to achieve comparable muscular adaptations (e.g., increase in muscle mass) as compared to high-load resistance training [271,272]. The lower exercise load during a resistance training with BFR could be beneficial for special cohorts since those lower exercise loads pose lower mechanical stress to the joints and the risk of adverse cardiovascular effects is decreased [124,217,244,273]. The currently available evidence suggests (i) that strength gains in response to a resistance training mediate, at least partly, the cognitive improvements [274] or (ii) that strength performance per se is a more appropriate indicator regarding health outcomes (e.g., cognition) [108]. Hence, at the moment no reliable assumptions can be made whether (i) a regular participation in resistance training, (ii) a relative high individual (baseline) strength level or (iii) the combination of both (high muscular strength level and regular resistance training) are most beneficial for cognitive functions. Notably, since an optimal level of neurochemical substances (e.g., IGF-1) is beneficial for cognitive performance [275], it could be speculated that, in turn, also an optimal level of muscular strength and/or continuously performed effective resistance activities, which may contribute substantially to the maintenance of an optimal level of neurochemical substances, exists. In this manner, a low-load resistance training with BFR could be a promising strategy especially for special cohorts (e.g., older adults unable to tolerate high loads) to ensure an adequate level of strength and profit from biological mechanisms which would without BFR only be possible when (not well tolerated) high loads are applied. Furthermore, relative low muscle damage is induced by low-load resistance training with BFR [148,168,276,277], which may allow a higher training frequency than in high load-resistance training [124,125,246].

However, testing the hypothesis suggesting that short-term and long-term resistance training with BFR improve cognitive performance as well as brain health to a greater extent than resistance training without BFR may provide deeper insights into the interplay between neurobiological mechanisms and cognitive processes. A deeper understanding of underlying exercise-induced and cognition-related neurobiological mechanisms is urgently needed to develop efficient prevention strategies (e.g., decelerate cognitive decline due to aging process) and to optimize rehabilitation strategies for individuals with worsened cognitive functions (e.g., older individuals with dementia). Here, the resistance training with BFR might be a promising strategy of exercise intervention.

**Author Contributions:** A.T. and F.H. wrote the manuscript and share the first authorship. L.S., N.G.M. and D.H. reviewed the drafted versions. All authors have read and approved the final version.

**Funding:** No funding information is applicable.

**Acknowledgments:** The authors have nothing to acknowledge.

**Conflicts of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## References

1. Adams, K.J.; O'Shea, P.; O'Shea, K.L. Aging: Its effects on strength, power, flexibility, and bone density. *Strength Cond. J.* **1999**, *21*, 65–77. [[CrossRef](#)]
2. Goldspink, G. Age-related loss of muscle mass and strength. *J. Aging Res.* **2012**, *2012*, 1–11. [[CrossRef](#)] [[PubMed](#)]
3. Goodpaster, B.H.; Park, S.W.; Harris, T.B.; Kritchevsky, S.B.; Nevitt, M.; Schwartz, A.V.; Simonsick, E.M.; Tyllavsky, F.A.; Visser, M.; Newman, A.B. The loss of skeletal muscle strength, mass, and quality in older adults: The health, aging and body composition study. *J. Gerontol. A Biol. Sci. Med. Sci.* **2006**, *61*, 1059–1064. [[CrossRef](#)] [[PubMed](#)]
4. Keller, K.; Engelhardt, M. Strength and muscle mass loss with aging process. Age and strength loss. *MLTJ* **2013**, *3*, 346–350. [[PubMed](#)]
5. Davis, J.W.; Ross, P.D.; Preston, S.D.; Nevitt, M.C.; Wasnich, R.D. Strength, physical activity, and body mass index: Relationship to performance-based measures and activities of daily living among older Japanese women in Hawaii. *J. Am. Geriatr. Soc.* **1998**, *46*, 274–279. [[CrossRef](#)] [[PubMed](#)]
6. Rantanen, T.; Avlund, K.; Suominen, H.; Schroll, M.; Frändin, K.; Pertti, E. Muscle strength as a predictor of onset of ADL dependence in people aged 75 years. *Aging Clin. Exp. Res.* **2002**, *14*, 10–15. [[PubMed](#)]
7. Wolfson, L.; JUDGE, J.; Whipple, R.; King, M. Strength is a major factor in balance, gait, and the occurrence of falls. *J. Gerontol. A Biol. Sci. Med. Sci.* **1995**, *50*, 64–67. [[PubMed](#)]
8. Horlings, C.G.C.; van Engelen, B.G.M.; Allum, J.H.J.; Bloem, B.R. A weak balance: The contribution of muscle weakness to postural instability and falls. *Nat. Clin. Pract. Neurol.* **2008**, *4*, 504–515. [[CrossRef](#)] [[PubMed](#)]
9. Scott, D.; Stuart, A.L.; Kay, D.; Ebeling, P.R.; Nicholson, G.; Sanders, K.M. Investigating the predictive ability of gait speed and quadriceps strength for incident falls in community-dwelling older women at high risk of fracture. *Arch. Gerontol. Geriatr.* **2014**, *58*, 308–313. [[CrossRef](#)] [[PubMed](#)]
10. Kilgour, A.H.M.; Todd, O.M.; Starr, J.M. A systematic review of the evidence that brain structure is related to muscle structure and their relationship to brain and muscle function in humans over the lifecourse. *BMC Geriatr.* **2014**, *14*, 85. [[CrossRef](#)] [[PubMed](#)]
11. Tolea, M.I.; Galvin, J.E. Sarcopenia and impairment in cognitive and physical performance. *Clin. Interv. Aging* **2015**, *10*, 663–671. [[CrossRef](#)] [[PubMed](#)]
12. Burns, J.M.; Johnson, D.K.; Watts, A.; Swerdlow, R.H.; Brooks, W.M. Reduced lean mass in early Alzheimer disease and its association with brain atrophy. *Arch. Neurol.* **2010**, *67*, 428–433. [[CrossRef](#)] [[PubMed](#)]
13. Nourhashemi, F.; Andrieu, S.; Gillette-Guyonnet, S.; Reynish, E.; Albaredo, J.-L.; Grandjean, H.; Vellas, B. Is there a relationship between fat-free soft tissue mass and low cognitive function? Results from a study of 7,105 women. *J. Am. Geriatr. Soc.* **2002**, *50*, 1796–1801. [[CrossRef](#)] [[PubMed](#)]
14. Hedden, T.; Gabrieli, J.D.E. Insights into the ageing mind: A view from cognitive neuroscience. *Nat. Rev. Neurosci.* **2004**, *5*, 87–96. [[CrossRef](#)] [[PubMed](#)]
15. Park, D.C.; Lautenschlager, G.; Hedden, T.; Davidson, N.S.; Smith, A.D.; Smith, P.K. Models of visuospatial and verbal memory across the adult life span. *Psychol. Aging* **2002**, *17*, 299–320. [[CrossRef](#)] [[PubMed](#)]
16. Albert, M.S. The ageing brain: Normal and abnormal memory. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **1997**, *352*, 1703–1709. [[CrossRef](#)] [[PubMed](#)]
17. Buckner, R.L. Memory and executive function in aging and AD: Multiple factors that cause decline and reserve factors that compensate. *Neuron* **2004**, *44*, 195–208. [[CrossRef](#)] [[PubMed](#)]
18. Fjell, A.M.; Walhovd, K.B. Structural brain changes in aging: Courses, causes and cognitive consequences. *Rev. Neurosci.* **2010**, *21*, 187–221. [[CrossRef](#)] [[PubMed](#)]



19. Fiest, K.M.; Jette, N.; Roberts, J.I.; Maxwell, C.J.; Smith, E.E.; Black, S.E.; Blaikie, L.; Cohen, A.; Day, L.; Holroyd-Leduc, J.; et al. The prevalence and incidence of dementia: A systematic review and meta-analysis. *Can. J. Neurol. Sci.* **2016**, *43*, 35. [[CrossRef](#)] [[PubMed](#)]
20. Berr, C.; Wancata, J.; Ritchie, K. Prevalence of dementia in the elderly in Europe. *Eur. Neuropsychopharmacol.* **2005**, *15*, 463–471. [[CrossRef](#)] [[PubMed](#)]
21. Scherder, E.; Eggermont, L.; Visscher, C.; Scheltens, P.; Swaab, D. Understanding higher level gait disturbances in mild dementia in order to improve rehabilitation: ‘Last in–first out’. *Neurosci. Biobehav. Rev.* **2011**, *35*, 699–714. [[CrossRef](#)] [[PubMed](#)]
22. Wimo, A.; Jonsson, L.; Bond, J.; Prince, M.; Winblad, B. The worldwide economic impact of dementia 2010. *Alzheimers. Dement.* **2013**, *9*, 1–11. [[CrossRef](#)] [[PubMed](#)]
23. Wimo, A.; Guerchet, M.; Ali, G.-C.; Wu, Y.-T.; Prina, A.M.; Winblad, B.; Jönsson, L.; Liu, Z.; Prince, M. The worldwide costs of dementia 2015 and comparisons with 2010. *Alzheimers. Dement.* **2017**, *13*, 1–7. [[CrossRef](#)] [[PubMed](#)]
24. Jönsson, L.; Wimo, A. The cost of dementia in Europe: A review of the evidence, and methodological considerations. *Pharmacoeconomics* **2009**, *27*, 391–403. [[CrossRef](#)] [[PubMed](#)]
25. Hurd, M.D.; Martorell, P.; Delavande, A.; Mullen, K.J.; Langa, K.M. Monetary costs of dementia in the United States. *N. Engl. J. Med.* **2013**, *368*, 1326–1334. [[CrossRef](#)] [[PubMed](#)]
26. Tariot, P.N. Pharmacological treatment of neuropsychiatric symptoms of dementia. *JAMA* **2005**, *293*, 2212. [[CrossRef](#)] [[PubMed](#)]
27. Ringman, J.M.; Cummings, J.L. Current and emerging pharmacological treatment options for dementia. *Behav. Neurol.* **2006**, *17*, 5–16. [[CrossRef](#)] [[PubMed](#)]
28. Morley, J.E. Pharmacologic options for the treatment of Sarcopenia. *Calcif. Tissue Int.* **2016**, *98*, 319–333. [[CrossRef](#)] [[PubMed](#)]
29. Schwarz, S.; Froelich, L.; Burns, A. Pharmacological treatment of dementia. *Curr. Opin. Psychiatry* **2012**, *25*, 542–550. [[CrossRef](#)] [[PubMed](#)]
30. Van de Glind, E.M.M.; van Enst, W.A.; van Munster, B.C.; Olde Rikkert, M.G.M.; Scheltens, P.; Scholten, R.J.P.M.; Hooft, L. Pharmacological treatment of dementia: A scoping review of systematic reviews. *Dement. Geriatr. Cogn. Disord.* **2013**, *36*, 211–228. [[CrossRef](#)] [[PubMed](#)]
31. Burton, L.A.; Sumukadas, D. Optimal management of sarcopenia. *CIA* **2010**, *5*, 217. [[CrossRef](#)]
32. Paillard-Borg, S.; Fratiglioni, L.; Winblad, B.; Wang, H.-X. Leisure activities in late life in relation to dementia risk: Principal component analysis. *Dement. Geriatr. Cogn. Disord.* **2009**, *28*, 136–144. [[CrossRef](#)] [[PubMed](#)]
33. Rovio, S.; Kåreholt, I.; Helkala, E.-L.; Viitanen, M.; Winblad, B.; Tuomilehto, J.; Soininen, H.; Nissinen, A.; Kivipelto, M. Leisure-time physical activity at midlife and the risk of dementia and Alzheimer’s disease. *Lancet Neurol.* **2005**, *4*, 705–711. [[CrossRef](#)]
34. Loprinzi, P.D.; Herod, S.M.; Cardinal, B.J.; Noakes, T.D. Physical activity and the brain: A review of this dynamic, bi-directional relationship. *Brain Res.* **2013**, *1539*, 95–104. [[CrossRef](#)] [[PubMed](#)]
35. Ponce, P.; Loprinzi, P.D. A bi-directional model of exercise and episodic memory function. *Med. Hypotheses* **2018**, *117*, 3–6. [[CrossRef](#)] [[PubMed](#)]
36. Pentikäinen, H.; Savonen, K.; Komulainen, P.; Kiviniemi, V.; Paajanen, T.; Kivipelto, M.; Soininen, H.; Rauramaa, R. Muscle strength and cognition in ageing men and women: The DR’s EXTRA study. *Eur. Geriatr. Med.* **2017**, *8*, 275–277. [[CrossRef](#)]
37. Chen, W.-L.; Peng, T.-C.; Sun, Y.-S.; Yang, H.-F.; Liaw, F.-Y.; Wu, L.-W.; Chang, Y.-W.; Kao, T.-W. Examining the association between quadriceps strength and cognitive performance in the elderly. *Medicine* **2015**, *94*, e1335. [[CrossRef](#)] [[PubMed](#)]
38. Alfaro-Acha, A.; Al Snih, S.; Raji, M.A.; Kuo, Y.-F.; Markides, K.S.; Ottenbacher, K.J. Handgrip strength and cognitive decline in older Mexican Americans. *J. Gerontol. A Biol. Sci. Med. Sci.* **2006**, *61*, 859–865. [[CrossRef](#)] [[PubMed](#)]
39. Hillman, C.H.; Erickson, K.I.; Kramer, A.F. Be smart, exercise your heart: Exercise effects on brain and cognition. *Nat. Rev. Neurosci.* **2008**, *9*, 58–65. [[CrossRef](#)] [[PubMed](#)]
40. Lustig, C.; Shah, P.; Seidler, R.; Reuter-Lorenz, P.A. Aging, training, and the brain: A review and future directions. *Neuropsychol. Rev.* **2009**, *19*, 504–522. [[CrossRef](#)] [[PubMed](#)]
41. Rolland, Y.; van Abellan Kan, G.; Vellas, B. Healthy brain aging: Role of exercise and physical activity. *Clin. Geriatr. Med.* **2010**, *26*, 75–87. [[CrossRef](#)] [[PubMed](#)]



42. Bherer, L.; Erickson, K.I.; Liu-Ambrose, T. A review of the effects of physical activity and exercise on cognitive and brain functions in older adults. *J. Aging Res.* **2013**. [[CrossRef](#)] [[PubMed](#)]
43. Voelcker-Rehage, C.; Niemann, C. Structural and functional brain changes related to different types of physical activity across the life span. *Neurosci. Biobehav. Rev.* **2013**, *37*, 2268–2295. [[CrossRef](#)] [[PubMed](#)]
44. Cai, Y.; Abrahamson, K. Does exercise impact cognitive performance research article op in community-dwelling older adults with mild cognitive impairment? A systematic review. *Qual. Prim. Care* **2015**, *23*, 214–222.
45. Esteban-Cornejo, I.; Tejero-Gonzalez, C.M.; SALLIS, J.F.; Veiga, O.L. Physical activity and cognition in adolescents: A systematic review. *J. Sci. Med. Sport* **2015**, *18*, 534–539. [[CrossRef](#)] [[PubMed](#)]
46. Lautenschlager, N.T. Twelve months of resistance training can improve the cognitive functioning of older women living in the community. *J. Physiother.* **2010**, *56*, 200. [[CrossRef](#)]
47. Powell, K.E.; Paluch, A.E.; Blair, S.N. Physical activity for health: What kind? How much? How intense? On top of what? *Annu. Rev. Public Health* **2011**, *32*, 349–365. [[CrossRef](#)] [[PubMed](#)]
48. Müller, P.; Schmicker, M.; Müller, N.G. Präventionsstrategien gegen Demenz. *Z. Gerontol. Geriatr.* **2017**. [[CrossRef](#)] [[PubMed](#)]
49. Herold, F.; Hamacher, D.; Schega, L.; Müller, N.G. Thinking while Moving or Moving while Thinking—Concepts of motor-cognitive training for cognitive performance enhancement. *Front. Aging Neurosci.* **2018**, *10*, 1–11. [[CrossRef](#)] [[PubMed](#)]
50. Davis, J.C.; Bryan, S.; Marra, C.A.; Hsiung, G.-Y.R.; Liu-Ambrose, T. Challenges with cost-utility analyses of behavioural interventions among older adults at risk for dementia. *Br. J. Sports Med.* **2015**, *49*, 1343–1347. [[CrossRef](#)] [[PubMed](#)]
51. Davis, J.C.; Marra, C.A.; Robertson, M.C.; Najafzadeh, M.; Liu-Ambrose, T. Sustained economic benefits of resistance training in community-dwelling senior women. *J. Am. Geriatr. Soc.* **2011**, *59*, 1232–1237. [[CrossRef](#)] [[PubMed](#)]
52. Davis, J.C.; Bryan, S.; Marra, C.A.; Sharma, D.; Chan, A.; Beattie, B.L.; Graf, P.; Liu-Ambrose, T. An economic evaluation of resistance training and aerobic training versus balance and toning exercises in older adults with mild cognitive impairment. *PLoS ONE* **2013**, *8*, e63031. [[CrossRef](#)] [[PubMed](#)]
53. Ashton, R.E.; Tew, G.A.; Aning, J.J.; Gilbert, S.E.; Lewis, L.; Saxton, J.M. Effects of short-term, medium-term and long-term resistance exercise training on cardiometabolic health outcomes in adults: Systematic review with meta-analysis. *Br. J. Sports Med.* **2018**. [[CrossRef](#)]
54. Hunter, G.R.; McCarthy, J.P.; Bamman, M.M. Effects of resistance training on older adults. *Sports Med.* **2004**, *34*, 329–348. [[CrossRef](#)] [[PubMed](#)]
55. Hurley, B.F.; Hanson, E.D.; Sheaff, A.K. Strength training as a countermeasure to aging muscle and chronic disease. *Sports Med.* **2011**, *41*, 289–306. [[CrossRef](#)] [[PubMed](#)]
56. Hurley, B.F.; Roth, S.M. Strength Training in the Elderly. *Sports Med.* **2000**, *30*, 249–268. [[CrossRef](#)] [[PubMed](#)]
57. Kraemer, W.J.; Ratamess, N.A.; French, D.N. Resistance Training for Health and Performance. *Curr. Sports Med. Rep.* **2002**, *1*, 165–171. [[CrossRef](#)] [[PubMed](#)]
58. Shaw, B.S.; Shaw, I.; Brown, G.A. Resistance exercise is medicine: Strength training in health promotion and rehabilitation. *Int. J. Ther. Rehabil.* **2015**, *22*, 385–389. [[CrossRef](#)]
59. Winett, R.A.; Carpinelli, R.N. Potential health-related benefits of resistance training. *Prev. Med.* **2001**, *33*, 503–513. [[CrossRef](#)] [[PubMed](#)]
60. Westcott, W.L. Resistance training is medicine: Effects of strength training on health. *Curr. Sports Med. Rep.* **2012**, *11*, 209–216. [[CrossRef](#)] [[PubMed](#)]
61. Chang, Y.-K.; Pan, C.-Y.; Chen, F.-T.; Tsai, C.-L.; Huang, C.-C. Effect of resistance-exercise training on cognitive function in healthy older adults: A review. *J. Aging Phys. Act.* **2012**, *20*, 497–517. [[CrossRef](#)] [[PubMed](#)]
62. Liu-Ambrose, T.; Donaldson, M.G. Exercise and cognition in older adults: Is there a role for resistance training programmes? *Br. J. Sports Med.* **2009**, *43*, 25–27. [[CrossRef](#)] [[PubMed](#)]
63. O'Connor, P.J.; Herring, M.P.; Carvalho, A. Mental health benefits of strength training in adults. *Am. J. Lifestyle Med.* **2010**, *4*, 377–396. [[CrossRef](#)]
64. Soga, K.; Masaki, H.; Gerber, M.; Ludyga, S. Acute and long-term effects of resistance training on executive function. *J. Cogn. Enhanc.* **2018**, *56*, 729. [[CrossRef](#)]
65. Altug, Z. Resistance exercise to improve cognitive function. *J. Strength Cond.* **2014**, *36*, 46–50. [[CrossRef](#)]

66. Voss, M.W.; Nagamatsu, L.S.; Liu-Ambrose, T.; Kramer, A.F. Exercise, brain, and cognition across the life span. *J. Appl. Physiol.* **2011**, *111*, 1505–1513. [[CrossRef](#)] [[PubMed](#)]
67. Stillman, C.M.; Cohen, J.; Lehman, M.E.; Erickson, K.I. Mediators of physical activity on neurocognitive function: A review at multiple levels of analysis. *Front. Hum. Neurosci.* **2016**, *10*, 626. [[CrossRef](#)] [[PubMed](#)]
68. Barha, C.K.; Galea, L.A.; Nagamatsu, L.S.; Erickson, K.I.; Liu-Ambrose, T. Personalising exercise recommendations for brain health: Considerations and future directions. *Br. J. Sports Med.* **2017**, *51*, 636–639. [[CrossRef](#)] [[PubMed](#)]
69. Cassilhas, R.C.; Lee, K.S.; Venâncio, D.P.; Oliveira, M.G.M.; Tufik, S.; Mello, M.T. de. Resistance exercise improves hippocampus-dependent memory. *Braz. J. Med. Biol. Res.* **2012**, *45*, 1215–1220. [[CrossRef](#)] [[PubMed](#)]
70. Cassilhas, R.C.; Lee, K.S.; Fernandes, J.; Oliveira, M.G.M.; Tufik, S.; Meeusen, R.; Mello de, M.T. Spatial memory is improved by aerobic and resistance exercise through divergent molecular mechanisms. *Neurosci.* **2012**, *202*, 309–317. [[CrossRef](#)] [[PubMed](#)]
71. Ohlsson, C.; Mohan, S.; Sjögren, K.; Tivesten, A.; Isgaard, J.; Isaksson, O.; Jansson, J.-O.; Svensson, J. The role of liver-derived insulin-like growth factor-I. *Endocr. Rev.* **2009**, *30*, 494–535. [[CrossRef](#)] [[PubMed](#)]
72. Gubbi, S.; Quipildor, G.F.; Barzilai, N.; Huffman, D.M.; Milman, S. 40 years of IGF1: IGF1: The Jekyll and Hyde of the aging brain. *J. Mol. Endocrinol.* **2018**, *61*, T171–T185. [[CrossRef](#)] [[PubMed](#)]
73. Bassil, F.; Fernagut, P.-O.; Bezard, E.; Meissner, W.G. Insulin, IGF-1 and GLP-1 signaling in neurodegenerative disorders: Targets for disease modification? *Prog. Neurobiol.* **2014**, *118*, 1–18. [[CrossRef](#)] [[PubMed](#)]
74. Dyer, A.H.; Vahdatpour, C.; Sanfeliu, A.; Tropea, D. The role of insulin-like growth factor 1 (IGF-1) in brain development, maturation and neuroplasticity. *Neuroscience* **2016**, *325*, 89–99. [[CrossRef](#)] [[PubMed](#)]
75. Deak, F.; Sonntag, W.E. Aging, synaptic dysfunction, and insulin-like growth factor (IGF)-1. *J. Gerontol. A Biol. Sci. Med. Sci.* **2012**, *67A*, 611–625. [[CrossRef](#)] [[PubMed](#)]
76. Aberg, N.D.; Brywe, K.G.; Isgaard, J. Aspects of growth hormone and insulin-like growth factor-I related to neuroprotection, regeneration, and functional plasticity in the adult brain. *Sci. World J.* **2006**, *6*, 53–80. [[CrossRef](#)] [[PubMed](#)]
77. Toth, P.; Tarantini, S.; Ashpole, N.M.; Tucsek, Z.; Milne, G.L.; Valcarcel-Ares, N.M.; Menyhart, A.; Farkas, E.; Sonntag, W.E.; Csiszar, A.; et al. IGF-1 deficiency impairs neurovascular coupling in mice: Implications for cerebrovascular aging. *Aging Cell* **2015**, *14*, 1034–1044. [[CrossRef](#)] [[PubMed](#)]
78. Saber, H.; Himali, J.J.; Beiser, A.S.; Shoamanesh, A.; Pikula, A.; Roubenoff, R.; Romero, J.R.; Kase, C.S.; Vasan, R.S.; Seshadri, S. Serum insulin-like growth factor 1 and the risk of ischemic stroke: The framingham study. *Stroke* **2017**, *48*, 1760–1765. [[CrossRef](#)] [[PubMed](#)]
79. Arwert, L.I.; Deijen, J.B.; Drent, M.L. The relation between insulin-like growth factor I levels and cognition in healthy elderly: A meta-analysis. *Growth Horm. IGF Res.* **2005**, *15*, 416–422. [[CrossRef](#)] [[PubMed](#)]
80. Calvo, D.; Gunstad, J.; Miller, L.A.; Glickman, E.; Spitznagel, M.B. Higher serum insulin-like growth factor-1 is associated with better cognitive performance in persons with mild cognitive impairment. *Psychogeriatrics* **2013**, *13*, 170–174. [[CrossRef](#)] [[PubMed](#)]
81. Westwood, A.J.; Beiser, A.; DeCarli, C.; Harris, T.B.; Chen, T.C.; He, X.-M.; Roubenoff, R.; Pikula, A.; Au, R.; Braverman, L.E.; et al. Insulin-like growth factor-1 and risk of Alzheimer dementia and brain atrophy. *Neurology* **2014**, *82*, 1613–1619. [[CrossRef](#)] [[PubMed](#)]
82. Rojas Vega, S.; Knicker, A.; Hollmann, W.; Bloch, W.; Strüder, H.K. Effect of resistance exercise on serum levels of growth factors in humans. *Horm. Metab. Res.* **2010**, *42*, 982–986. [[CrossRef](#)] [[PubMed](#)]
83. Tsai, C.-L.; Wang, C.-H.; Pan, C.-Y.; Chen, F.-C. The effects of long-term resistance exercise on the relationship between neurocognitive performance and GH, IGF-1, and homocysteine levels in the elderly. *Front. Behav. Neurosci.* **2015**, *9*, 471. [[CrossRef](#)] [[PubMed](#)]
84. Borst, S.E.; Hoyos de, D.V.; Garzarella, L.; Vincent, K.; Pollock, B.H.; Lowenthal, D.T.; Pollock, M.L. Effects of resistance training on insulin-like growth factor-I and IGF binding proteins. *Med. Sci. Sports Exerc.* **2001**, *33*, 648–653. [[CrossRef](#)] [[PubMed](#)]
85. Stein, A.M.; Silva, T.M.V.; Coelho, F.G.D.M.; Arantes, F.J.; Costa, J.L.R.; Teodoro, E.; Santos-Galduróz, R.F. Physical exercise, IGF-1 and cognition A systematic review of experimental studies in the elderly. *Dement. Neuropsychol.* **2018**, *12*, 114–122. [[CrossRef](#)] [[PubMed](#)]

86. Fontes, E.B.; Libardi, C.A.; Castellano, G.; Okano, A.H.; Fernandes, P.T.; Chacon-Mikahil, M.P.; Ugrinowitsch, C.; Cavaglieri, C.R.; Li, L.M. Effects of resistance training in gray matter density of elderly. *Sport Sci. Health* **2017**, *13*, 233–238. [[CrossRef](#)]
87. Suo, C.; Singh, M.F.; Gates, N.; Wen, W.; Sachdev, P.; Brodaty, H.; Saigal, N.; Wilson, G.C.; Meiklejohn, J.; Singh, N.; et al. Therapeutically relevant structural and functional mechanisms triggered by physical and cognitive exercise. *Mol. Psychiatry* **2016**, *21*, 1633–1642. [[CrossRef](#)] [[PubMed](#)]
88. Liu-Ambrose, T.; Nagamatsu, L.S.; Graf, P.; Beattie, B.L.; Ashe, M.C.; Handy, T.C. Resistance training and executive functions: A 12-month randomized controlled trial. *Arch. Intern. Med.* **2010**, *170*, 170–178. [[CrossRef](#)] [[PubMed](#)]
89. Fox, N.C.; Black, R.S.; Gilman, S.; Rossor, M.N.; Griffith, S.G.; Jenkins, L.; Koller, M. Effects of Abeta immunization (AN1792) on MRI measures of cerebral volume in Alzheimer disease. *Neurology* **2005**, *64*, 1563–1572. [[CrossRef](#)] [[PubMed](#)]
90. Ströhlein, J.K.; van den Bongard, F.; Barthel, T.; Reinsberger, C. Dose-response-relationship between physical activity and cognition in elderly. *Dtsch. Z. Sportmed.* **2017**, *2017*, 234–242. [[CrossRef](#)]
91. Voss, M.W.; Vivar, C.; Kramer, A.F.; van Praag, H. Bridging animal and human models of exercise-induced brain plasticity. *Trends Cogn. Sci.* **2013**, *17*, 525–544. [[CrossRef](#)] [[PubMed](#)]
92. Stimpson, N.J.; Davison, G.; Javadi, A.-H. Joggin' the noggin: Towards a physiological understanding of exercise-induced cognitive benefits. *Neurosci. Biobehav. Rev.* **2018**, *88*, 177–186. [[CrossRef](#)] [[PubMed](#)]
93. Gomes-Osman, J.; Cabral, D.F.; Morris, T.P.; McInerney, K.; Cahalin, L.P.; Rundek, T.; Oliveira, A.; Pascual-Leone, A. Exercise for cognitive brain health in aging. *Neurol. Clin. Pract.* **2018**, *8*, 257–265. [[CrossRef](#)] [[PubMed](#)]
94. Best, J.R.; Chiu, B.K.; Liang Hsu, C.; Nagamatsu, L.S.; Liu-Ambrose, T. Long-term effects of resistance exercise training on cognition and brain volume in older women: Results from a randomized controlled trial. *J. Int. Neuropsychol. Soc.* **2015**, *21*, 745–756. [[CrossRef](#)] [[PubMed](#)]
95. Bolandzadeh, N.; Tam, R.; Handy, T.C.; Nagamatsu, L.S.; Hsu, C.L.; Davis, J.C.; Dao, E.; Beattie, B.L.; Liu-Ambrose, T. Resistance training and white matter lesion progression in older women: Exploratory analysis of a 12-month randomized controlled trial. *J. Am. Geriatr. Soc.* **2015**, *63*, 2052–2060. [[CrossRef](#)] [[PubMed](#)]
96. de Groot, J.C.; de Leeuw, F.E.; Oudkerk, M.; van Gijn, J.; Hofman, A.; Jolles, J.; Breteler, M.M. Cerebral white matter lesions and cognitive function: The rotterdam scan study. *Ann. Neurol.* **2000**, *47*, 145–151. [[CrossRef](#)]
97. Gunning-Dixon, F.M.; Raz, N. The cognitive correlates of white matter abnormalities in normal aging: A quantitative review. *Neuropsychology* **2000**, *14*, 224–232. [[CrossRef](#)] [[PubMed](#)]
98. Vernooij, M.W.; Ikram, M.A.; Vrooman, H.A.; Wielopolski, P.A.; Krestin, G.P.; Hofman, A.; Niessen, W.J.; van der Lugt, A.; Breteler, M.M.B. White matter microstructural integrity and cognitive function in a general elderly population. *Arch. Gen. Psychiatry* **2009**, *66*, 545–553. [[CrossRef](#)] [[PubMed](#)]
99. Gunning-Dixon, F.M.; Brickman, A.M.; Cheng, J.C.; Alexopoulos, G.S. Aging of cerebral white matter: A review of MRI findings. *Int. J. Geriatr. Psychiatry* **2009**, *24*, 109–117. [[CrossRef](#)] [[PubMed](#)]
100. Alves, C.R.R.; Tessaro, V.H.; Teixeira, L.A.C.; Murakava, K.; Roschel, H.; Gualano, B.; Takito, M.Y. Influence of acute high-intensity aerobic interval exercise bout on selective attention and short-term memory tasks. *Percept. Mot. Skills* **2014**, *118*, 63–72. [[CrossRef](#)] [[PubMed](#)]
101. Chang, H.; Kim, K.; Jung, Y.-J.; Kato, M. Effects of acute high-intensity resistance exercise on cognitive function and oxygenation in prefrontal cortex. *J. Exerc. Nutrition Biochem.* **2017**, *21*, 1–8. [[CrossRef](#)] [[PubMed](#)]
102. Liu-Ambrose, T.; Nagamatsu, L.S.; Voss, M.W.; Khan, K.M.; Handy, T.C. Resistance training and functional plasticity of the aging brain: A 12-month randomized controlled trial. *Neurobiol. Aging* **2012**, *33*, 1690–1698. [[CrossRef](#)] [[PubMed](#)]
103. Nagamatsu, L.S.; Handy, T.C.; Hsu, C.L.; Voss, M.; Liu-Ambrose, T. Resistance training promotes cognitive and functional brain plasticity in seniors with probable mild cognitive impairment. *Arch. Intern. Med.* **2012**, *172*, 666–668. [[CrossRef](#)] [[PubMed](#)]
104. Coetsee, C.; Terblanche, E. Cerebral oxygenation during cortical activation: The differential influence of three exercise training modalities. A randomized controlled trial. *Eur. J. Appl. Physiol.* **2017**. [[CrossRef](#)] [[PubMed](#)]
105. Firth, J.; Firth, J.A.; Stubbs, B.; Vancampfort, D.; Schuch, F.B.; Hallgren, M.; Veronese, N.; Yung, A.R.; Sarris, J. Association Between Muscular Strength and Cognition in People With Major Depression or Bipolar Disorder and Healthy Controls. *JAMA Psychiatry* **2018**. [[CrossRef](#)] [[PubMed](#)]

106. Firth, J.; Stubbs, B.; Vancampfort, D.; Firth, J.A.; Large, M.; Rosenbaum, S.; Hallgren, M.; Ward, P.B.; Sarris, J.; Yung, A.R. Grip Strength Is Associated With Cognitive Performance in Schizophrenia and the General Population: A UK Biobank Study of 476559 Participants. *Schizophrenia Bull.* **2018**, *44*, 728–736. [[CrossRef](#)] [[PubMed](#)]
107. Steves, C.J.; Mehta, M.M.; Jackson, S.H.D.; Spector, T.D. Kicking Back Cognitive Ageing: Leg Power Predicts Cognitive Ageing after Ten Years in Older Female Twins. *Gerontology* **2016**, *62*, 138–149. [[CrossRef](#)] [[PubMed](#)]
108. Buckner, S.L.; Dankel, S.J.; Mouser, J.G.; Mattocks, K.T.; Jessee, M.B.; Loenneke, J.P. Chasing the top quartile of cross-sectional data: Is it possible with resistance training? *Med. Hypotheses* **2017**, *108*, 63–68. [[CrossRef](#)] [[PubMed](#)]
109. Decker, L.M.; Cignetti, F.; Stergiou, N. Executive function orchestrates regulation of task-relevant gait fluctuations. *Gait Posture* **2013**, *38*, 537–540. [[CrossRef](#)] [[PubMed](#)]
110. IJmker, T.; Lamoth, C.J.C. Gait and cognition: The relationship between gait stability and variability with executive function in persons with and without dementia. *Gait Posture* **2012**, *35*, 126–130. [[CrossRef](#)] [[PubMed](#)]
111. Kearney, F.C.; Harwood, R.H.; Gladman, J.R.F.; Lincoln, N.; Masud, T. The relationship between executive function and falls and gait abnormalities in older adults: A systematic review. *Dement. Geriatr. Cogn. Disord.* **2013**, *36*, 20–35. [[CrossRef](#)] [[PubMed](#)]
112. Herold, F.; Wiegel, P.; Scholkmann, F.; Thiers, A.; Hamacher, D.; Schega, L. Functional near-infrared spectroscopy in movement science: A systematic review on cortical activity in postural and walking tasks. *Neurophoton* **2017**, *4*, 41403. [[CrossRef](#)] [[PubMed](#)]
113. Hamacher, D.; Herold, F.; Wiegel, P.; Hamacher, D.; Schega, L. Brain activity during walking: A systematic review. *Neurosci. Biobehav. Rev.* **2015**, *57*, 310–327. [[CrossRef](#)] [[PubMed](#)]
114. Saraçlı, Ö.; Akca, A.S.D.; Atasoy, N.; Önder, Ö.; Şenormancı, Ö.; Kaygisız, I.; Atik, L. The relationship between quality of life and cognitive functions, anxiety and depression among hospitalized elderly patients. *Clin. Psychopharmacol. Neurosci.* **2015**, *13*, 194–200. [[CrossRef](#)] [[PubMed](#)]
115. Kekäläinen, T.; Kokko, K.; Sipilä, S.; Walker, S. Effects of a 9-month resistance training intervention on quality of life, sense of coherence, and depressive symptoms in older adults: Randomized controlled trial. *Qual. Life Res.* **2018**, *27*, 455–465. [[CrossRef](#)] [[PubMed](#)]
116. Barha, C.K.; Davis, J.C.; Falck, R.S.; Nagamatsu, L.S.; Liu-Ambrose, T. Sex differences in exercise efficacy to improve cognition: A systematic review and meta-analysis of randomized controlled trials in older humans. *Front. Neuroendocrinol.* **2017**. [[CrossRef](#)] [[PubMed](#)]
117. Jessee, M.B.; Mattocks, K.T.; Buckner, S.L.; Dankel, S.J.; Mouser, J.G.; Abe, T.; Loenneke, J.P. Mechanisms of blood flow restriction: The New Testament. *Tech. Orthop.* **2018**, *33*, 72–79. [[CrossRef](#)]
118. Slys, J.; Stultz, J.; Burr, J.F. The efficacy of blood flow restricted exercise: A systematic review & meta-analysis. *J. Sci. Med. Sport* **2016**, *19*, 669–675. [[CrossRef](#)] [[PubMed](#)]
119. Pearson, S.J.; Hussain, S.R. A review on the mechanisms of blood-flow restriction resistance training-induced muscle hypertrophy. *Sports Med.* **2015**, *45*, 187–200. [[CrossRef](#)] [[PubMed](#)]
120. Bird, S.P.; Tarpenning, K.M.; Marino, F.E. Designing resistance training programmes to enhance muscular fitness: A review of the acute programme variables. *Sports Med.* **2005**, *35*, 841–851. [[CrossRef](#)] [[PubMed](#)]
121. Chang, Y.-K.; Chu, I.-H.; Chen, F.-T.; Wang, C.-C. Dose-response effect of acute resistance exercise on tower of London in middle-aged adults. *J. Sport Exerc. Psychol.* **2011**, *33*, 866–883. [[CrossRef](#)] [[PubMed](#)]
122. Chang, Y.-K.; Etner, J.L. Exploring the dose-response relationship between resistance exercise intensity and cognitive function. *J. Sport Exerc. Psychol.* **2009**, *31*, 640–656. [[CrossRef](#)] [[PubMed](#)]
123. Scott, B.R.; Loenneke, J.P.; Slattery, K.M.; Dascombe, B.J. Blood flow restricted exercise for athletes: A review of available evidence. *J. Sci. Med. Sport* **2015**. [[CrossRef](#)] [[PubMed](#)]
124. Scott, B.R.; Loenneke, J.P.; Slattery, K.M.; Dascombe, B.J. Exercise with blood flow restriction: An updated evidence-based approach for enhanced muscular development. *Sports Med.* **2015**, *45*, 313–325. [[CrossRef](#)] [[PubMed](#)]
125. Scott, B.R.; Slattery, K.M.; Sculley, D.V.; Dascombe, B.J. Hypoxia and resistance exercise: A comparison of localized and systemic methods. *Sports Med.* **2014**, *44*, 1037–1054. [[CrossRef](#)] [[PubMed](#)]
126. Ramos-Campo, D.J.; Scott, B.R.; Alcaraz, P.E.; Rubio-Arias, J.A. The efficacy of resistance training in hypoxia to enhance strength and muscle growth: A systematic review and meta-analysis. *Euro. J. Sport Sci* **2018**, *18*, 92–103. [[CrossRef](#)] [[PubMed](#)]



127. de Castro, F.M.P.; Aquino, R.; Berti, J.A.; Gonçalves, L.G.C.; Puggina, E.F. Strength training with vascular occlusion: A review of possible adaptive mechanisms. *Hum. Mov.* **2017**, *18*, 1974. [[CrossRef](#)]
128. Kawada, S. What phenomena do occur in blood flow-restricted muscle? *Int. J. KAATSU Train. Res.* **2005**, *1*, 37–44. [[CrossRef](#)]
129. Manini, T.M.; Clark, B.C. Blood flow restricted exercise and skeletal muscle health. *Exerc. Sport Sci. Rev.* **2009**, *37*, 78–85. [[CrossRef](#)] [[PubMed](#)]
130. Muhammad, K.; Hamed, I. Moderate Blood Flow Restriction Training. *MOJSM* **2017**, *1*. [[CrossRef](#)]
131. Sato, Y. The history and future of KAATSU Training. *Int. J. KAATSU Train. Res.* **2005**, *1*, 1–5. [[CrossRef](#)]
132. Lopata, V.A.; Serebrovskaya, T.V. Hypoxicators: Review of the operating principles and constructions. In *Intermittent Hypoxia and Human Diseases*; Springer: London, UK, 2012; pp. 291–302.
133. Nakajima, T.; Kurano, M.; Iida, H.; Takano, H.; Oounuma, H.; Morita, T.; Meguro, K.; Sato, Y.; Nagata, T. Use and safety of KAATSU training: Results of a national survey. *Int. J. KAATSU Train. Res.* **2006**, *2*, 5–13. [[CrossRef](#)]
134. Nakajima, T.; Morita, T.; Sato, Y. Key considerations when conducting KAATSU training. *Int. J. KAATSU Train. Res.* **2011**, *7*, 1–6. [[CrossRef](#)]
135. Navarrete-Opazo, A.; Mitchell, G.S. Therapeutic potential of intermittent hypoxia: A matter of dose. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2014**, *307*, R1181–R1197. [[CrossRef](#)] [[PubMed](#)]
136. Yasuda, T.; Meguro, M.; Sato, Y.; Nakajima, T. Use and safety of KAATSU training: Results of a national survey in 2016. *Int. J. KAATSU Train. Res.* **2017**, *13*, 1–9. [[CrossRef](#)]
137. Abe, T.; Loenneke, J.P.; Fahs, C.A.; Rossow, L.M.; Thiebaud, R.S.; Bemben, M.G. Exercise intensity and muscle hypertrophy in blood flow-restricted limbs and non-restricted muscles: A brief review. *Clin. Physiol. Funct. Imaging* **2012**, *32*, 247–252. [[CrossRef](#)] [[PubMed](#)]
138. Madarame, H.; Neya, M.; Ochi, E.; Nakazato, K.; Sato, Y.; Ishii, N. Cross-transfer effects of resistance training with blood flow restriction. *Med. Sci. Sports Exerc.* **2008**, *40*, 258–263. [[CrossRef](#)] [[PubMed](#)]
139. Manukhina, E.B.; Downey, H.F.; Shi, X.; Mallet, R.T. Intermittent hypoxia training protects cerebrovascular function in Alzheimer's disease. *Exp. Biol. Med.* **2016**, *241*, 1351–1363. [[CrossRef](#)] [[PubMed](#)]
140. Richardson, C.; Hogan, A.M.; Bucks, R.S.; Baya, A.; Virues-Ortega, J.; Holloway, J.W.; Rose-Zerilli, M.; Palmer, L.J.; Webster, R.J.; Kirkham, F.J.; et al. Neurophysiological evidence for cognitive and brain functional adaptation in adolescents living at high altitude. *Clin. Neurophysiol.* **2011**, *122*, 1726–1734. [[CrossRef](#)] [[PubMed](#)]
141. Schega, L.; Peter, B.; Törpel, A.; Mutschler, H.; Isermann, B.; Hamacher, D. Effects of intermittent hypoxia on cognitive performance and quality of life in elderly adults: A pilot study. *Gerontology* **2013**, *59*, 316–323. [[CrossRef](#)] [[PubMed](#)]
142. Schega, L.; Peter, B.; Brigadski, T.; Lessmann, V.; Isermann, B.; Hamacher, D.; Torpel, A. Effect of intermittent normobaric hypoxia on aerobic capacity and cognitive function in older people. *J. Sci. Med. Sport* **2016**. [[CrossRef](#)] [[PubMed](#)]
143. Sardeli, A.V.; Ferreira, M.L.V.; Santos, L.d.C.; Rodrigues, M.d.S.; Damasceno, A.; Cavaglieri, C.R.; Chacon-Mikahil, M.P.T. Low-load resistance exercise improves cognitive function in older adults. *Rev. Bras. Med. Esporte.* **2018**, *24*, 125–129. [[CrossRef](#)]
144. Loenneke, J.P.; Pujol, T.J. The use of occlusion training to produce muscle hypertrophy. *Strength Cond. J.* **2009**, *31*, 77–84. [[CrossRef](#)]
145. Takano, H.; Morita, T.; Iida, H.; Asada, K.-I.; Kato, M.; Uno, K.; Hirose, K.; Matsumoto, A.; Takenaka, K.; Hirata, Y.; et al. Hemodynamic and hormonal responses to a short-term low-intensity resistance exercise with the reduction of muscle blood flow. *Eur. J. Appl. Physiol.* **2005**, *95*, 65–73. [[CrossRef](#)] [[PubMed](#)]
146. Manini, T.M.; Yarrow, J.F.; Buford, T.W.; Clark, B.C.; Conover, C.F.; Borst, S.E. Growth hormone responses to acute resistance exercise with vascular restriction in young and old men. *Growth Horm. IGF Res.* **2012**, *22*, 167–172. [[CrossRef](#)] [[PubMed](#)]
147. Patterson, S.D.; Leggate, M.; Nimmo, M.A.; Ferguson, R.A. Circulating hormone and cytokine response to low-load resistance training with blood flow restriction in older men. *Eur. J. Appl. Physiol.* **2013**, *113*, 713–719. [[CrossRef](#)] [[PubMed](#)]
148. Takarada, Y.; Nakamura, Y.; Aruga, S.; Onda, T.; Miyazaki, S.; Ishii, N. Rapid increase in plasma growth hormone after low-intensity resistance exercise with vascular occlusion. *Eur. J. Appl. Physiol.* **2000**, *88*, 61–65. [[CrossRef](#)] [[PubMed](#)]

149. Takano, H.; Morita, T.; Iida, H.; Kato, M.; Uno, K.; Hirose, K.; Matsumoto, A.; Takenaka, K.; Hirata, Y.; Furuichi, T.; et al. Effects of low-intensity “KAATSU” resistance exercise on hemodynamic and growth hormone responses. *Int. J. KAATSU Train. Res.* **2005**, *1*, 13–18. [[CrossRef](#)]
150. Ferguson, R.A.; Hunt, J.E.A.; Lewis, M.P.; Martin, N.R.W.; Player, D.J.; Stangier, C.; Taylor, C.W.; Turner, M.C. The acute angiogenic signalling response to low-load resistance exercise with blood flow restriction. *Eur. J. Sport Sci.* **2018**, *18*, 397–406. [[CrossRef](#)] [[PubMed](#)]
151. Larkin, K.A.; Macneil, R.G.; Dirain, M.; Sandesara, B.; Manini, T.M.; Buford, T.W. Blood flow restriction enhances post-resistance exercise angiogenic gene expression. *Med. Sci. Sports Exerc.* **2012**, *44*, 2077–2083. [[CrossRef](#)] [[PubMed](#)]
152. Abe, T.; Yasuda, T.; Midorikawa, T.; Sato, Y.; Kearns, C.F.; Inoue, K.; Koizumi, K.; Ishii, N. Skeletal muscle size and circulating IGF-1 are increased after two weeks of twice daily “KAATSU” resistance training. *Int. J. KAATSU Train. Res.* **2005**, *1*, 6–12. [[CrossRef](#)]
153. Nyberg, F.; Hallberg, M. Growth hormone and cognitive function. *Nat. Rev. Endocrinol.* **2013**, *9*, 357–365. [[CrossRef](#)] [[PubMed](#)]
154. Van Dam, P.S.; Aleman, A.; de Vries, W.R.; Deijen, J.B.; van der Veen, E.A.; de Haan, E.H.; Koppeschaar, H.P. Growth hormone, insulin-like growth factor I and cognitive function in adults. *Growth Horm. IGF Res.* **2000**, *10*, S69–S73. [[CrossRef](#)]
155. Ari, Z.; Kutlu, N.; Uyanik, B.S.; Taneli, F.; Buyukyazi, G.; Tavli, T. Serum testosterone, growth hormone, and insulin-like growth factor-1 levels, mental reaction time, and maximal aerobic exercise in sedentary and long-term physically trained elderly males. *Int. J. Neurosci.* **2004**, *114*, 623–637. [[CrossRef](#)] [[PubMed](#)]
156. Cotman, C.W.; Berchtold, N.C.; Christie, L.-A. Exercise builds brain health: Key roles of growth factor cascades and inflammation. *Trends Neurosci.* **2007**, *30*, 464–472. [[CrossRef](#)] [[PubMed](#)]
157. Lucas, S.J.E.; Cotter, J.D.; Brassard, P.; Bailey, D.M. High-intensity interval exercise and cerebrovascular health: Curiosity, cause, and consequence. *J. Cereb. Blood Flow Metab.* **2015**, *35*, 902–911. [[CrossRef](#)] [[PubMed](#)]
158. Davenport, M.H.; Hogan, D.B.; Eskes, G.A.; Longman, R.S.; Poulin, M.J. Cerebrovascular reserve: The link between fitness and cognitive function? *Exerc. Sport Sci. Rev.* **2012**, *40*, 153–158. [[CrossRef](#)] [[PubMed](#)]
159. Basso, J.C.; Suzuki, W.A. The effects of acute exercise on mood, cognition, neurophysiology, and neurochemical pathways: A review. *BPL* **2017**, *2*, 127–152. [[CrossRef](#)] [[PubMed](#)]
160. Adams, R.H.; Alitalo, K. Molecular regulation of angiogenesis and lymphangiogenesis. *Nat. Rev. Mol. Cell Biol.* **2007**, *8*, 464–478. [[CrossRef](#)] [[PubMed](#)]
161. Eichmann, A.; Simons, M. VEGF signaling inside vascular endothelial cells and beyond. *Curr. Opin. Cell Biol.* **2012**, *24*, 188–193. [[CrossRef](#)] [[PubMed](#)]
162. Huang, L.; Jia, J.; Liu, R. Decreased serum levels of the angiogenic factors VEGF and TGF- $\beta$ 1 in Alzheimer’s disease and amnesic mild cognitive impairment. *Neurosci. Lett.* **2013**, *550*, 60–63. [[CrossRef](#)] [[PubMed](#)]
163. Hohman, T.J.; Bell, S.P.; Jefferson, A.L. The role of vascular endothelial growth factor in neurodegeneration and cognitive decline: Exploring interactions with biomarkers of Alzheimer disease. *JAMA Neurol.* **2015**, *72*, 520–529. [[CrossRef](#)] [[PubMed](#)]
164. Gentil, P.; Oliveira, E.; Bottaro, M. Time under Tension and Blood Lactate Response during Four Different Resistance Training Methods. *J. Physiol. Anthropol.* **2006**, *25*, 339–344. [[CrossRef](#)] [[PubMed](#)]
165. Fujita, S.; Abe, T.; Drummond, M.J.; Cadenas, J.G.; Dreyer, H.C.; Sato, Y.; Volpi, E.; Rasmussen, B.B. Blood flow restriction during low-intensity resistance exercise increases S6K1 phosphorylation and muscle protein synthesis. *J. Appl. Physiol.* **2007**, *103*, 903–910. [[CrossRef](#)] [[PubMed](#)]
166. Reeves, G.V.; Kraemer, R.R.; Hollander, D.B.; Clavier, J.; Thomas, C.; Francois, M.; Castracane, V.D. Comparison of hormone responses following light resistance exercise with partial vascular occlusion and moderately difficult resistance exercise without occlusion. *J. Appl. Physiol.* **2006**, *101*, 1616–1622. [[CrossRef](#)] [[PubMed](#)]
167. Yasuda, T.; Abe, T.; Brechue, W.F.; Iida, H.; Takano, H.; Meguro, K.; Kurano, M.; Fujita, S.; Nakajima, T. Venous blood gas and metabolite response to low-intensity muscle contractions with external limb compression. *Metabolism* **2010**, *59*, 1510–1519. [[CrossRef](#)] [[PubMed](#)]
168. Wilson, J.M.; Lowery, R.P.; Joy, J.M.; Loenneke, J.P.; Naimo, M.A. Practical blood flow restriction training increases acute determinants of hypertrophy without increasing indices of muscle damage. *J. Strength Cond. Res.* **2013**, *27*, 3068–3075. [[CrossRef](#)] [[PubMed](#)]



169. Loenneke, J.P.; Balapur, A.; Thrower, A.D.; Barnes, J.; Pujol, T.J. Blood flow restriction reduces time to muscular failure. *Eur. J. Sport Sci.* **2012**, *12*, 238–243. [[CrossRef](#)]
170. Loenneke, J.P.; Wilson, J.M.; Balapur, A.; Thrower, A.D.; Barnes, J.T.; Pujol, T.J. Time under tension decreased with blood flow-restricted exercise. *Clin. Physiol. Funct. Imaging* **2012**, *32*, 268–273. [[CrossRef](#)] [[PubMed](#)]
171. Lee, J.K.W.; Koh, A.C.H.; Koh, S.X.T.; Liu, G.J.X.; Nio, A.Q.X.; Fan, P.W.P. Neck cooling and cognitive performance following exercise-induced hyperthermia. *Eur. J. Appl. Physiol.* **2014**, *114*, 375–384. [[CrossRef](#)] [[PubMed](#)]
172. Tsukamoto, H.; Suga, T.; Takenaka, S.; Tanaka, D.; Takeuchi, T.; Hamaoka, T.; Isaka, T.; Ogoh, S.; Hashimoto, T. Repeated high-intensity interval exercise shortens the positive effect on executive function during post-exercise recovery in healthy young males. *Physiol. Behav.* **2016**, *160*, 26–34. [[CrossRef](#)] [[PubMed](#)]
173. Hashimoto, T.; Tsukamoto, H.; Takenaka, S.; Olesen, N.D.; Petersen, L.G.; Sørensen, H.; Nielsen, H.B.; Secher, N.H.; Ogoh, S. Maintained exercise-enhanced brain executive function related to cerebral lactate metabolism in men. *FASEB J.* **2017**. [[CrossRef](#)] [[PubMed](#)]
174. Riske, L.; Thomas, R.K.; Baker, G.B.; Dursun, S.M. Lactate in the brain: An update on its relevance to brain energy, neurons, glia and panic disorder. *Ther. Adv. Psychopharmacol.* **2017**, *7*, 85–89. [[CrossRef](#)] [[PubMed](#)]
175. Taher, M.; Leen, W.G.; Wevers, R.A.; Willemsen, M.A. Lactate and its many faces. *Eur. J. Paediatr. Neurol.* **2016**, *20*, 3–10. [[CrossRef](#)] [[PubMed](#)]
176. Van Hall, G.; Stromstad, M.; Rasmussen, P.; Jans, O.; Zaar, M.; Gam, C.; Quistorff, B.; Secher, N.H.; Nielsen, H.B. Blood lactate is an important energy source for the human brain. *J. Cereb. Blood Flow Metab.* **2009**, *29*, 1121–1129. [[CrossRef](#)] [[PubMed](#)]
177. Brooks, G.A. The Science and Translation of Lactate Shuttle Theory. *Cell Metab.* **2018**, *27*, 757–785. [[CrossRef](#)] [[PubMed](#)]
178. Quistorff, B.; Secher, N.H.; van Lieshout, J.J. Lactate fuels the human brain during exercise. *FASEB J.* **2008**, *22*, 3443–3449. [[CrossRef](#)] [[PubMed](#)]
179. Ferris, L.T.; Williams, J.S.; Shen, C.-L. The effect of acute exercise on serum brain-derived neurotrophic factor levels and cognitive function. *Med. Sci. Sports Exerc.* **2007**, *39*, 728–734. [[CrossRef](#)] [[PubMed](#)]
180. Schiffer, T.; Schulte, S.; Sperlich, B.; Achtzehn, S.; Fricke, H.; Struder, H.K. Lactate infusion at rest increases BDNF blood concentration in humans. *Neurosci. Lett.* **2011**, *488*, 234–237. [[CrossRef](#)] [[PubMed](#)]
181. Borrer, A. Brain-derived neurotrophic factor mediates cognitive improvements following acute exercise. *Med. Hypotheses* **2017**, *106*, 1–5. [[CrossRef](#)] [[PubMed](#)]
182. Hwang, J.; Brothers, R.M.; Castelli, D.M.; Glowacki, E.M.; Chen, Y.T.; Salinas, M.M.; Kim, J.; Jung, Y.; Calvert, H. Acute high-intensity exercise-induced cognitive enhancement and brain-derived neurotrophic factor in young, healthy adults. *Neurosci. Lett.* **2016**, *630*, 247–253. [[CrossRef](#)] [[PubMed](#)]
183. Semenza, G.L. Hypoxia-inducible factor 1: Master regulator of O<sub>2</sub> homeostasis. *Curr. Opin. Genet. Dev.* **1998**, *8*, 588–594. [[CrossRef](#)]
184. Schofield, C.J.; Ratcliffe, P.J. Oxygen sensing by HIF hydroxylases. *Nat. Rev. Mol. Cell Biol.* **2004**, *5*, 343–354. [[CrossRef](#)] [[PubMed](#)]
185. Laurentino, G.; Aoki, M.; Fernandes, R.; Soares, A.; Ugrinowitsch, C.; Hoschel, H.; Tricoli, V. Low-load resistance exercise with blood flow restriction changes hypoxia-induced genes expression. *FASEB J.* **2018**, *32*. [[CrossRef](#)]
186. Correia, S.C.; Moreira, P.I. Hypoxia-inducible factor 1: A new hope to counteract neurodegeneration? *J. Neurochem.* **2010**, *112*, 1–12. [[CrossRef](#)] [[PubMed](#)]
187. Ferrara, N.; Gerber, H.-P.; LeCouter, J. The biology of VEGF and its receptors. *Nat. Med.* **2003**, *9*, 669–676. [[CrossRef](#)] [[PubMed](#)]
188. Slomiany, M.G.; Rosenzweig, S.A. Hypoxia-inducible factor-1-dependent and -independent regulation of insulin-like growth factor-1-stimulated vascular endothelial growth factor secretion. *J. Pharmacol. Exp. Ther.* **2006**, *318*, 666–675. [[CrossRef](#)] [[PubMed](#)]
189. Brandner, C.R.; Warmington, S.A.; Kidgell, D.J. Corticomotor Excitability is Increased Following an Acute Bout of Blood Flow Restriction Resistance Exercise. *Front. Hum. Neurosci.* **2015**, *9*, 652. [[CrossRef](#)] [[PubMed](#)]
190. Morita, T.; Fukuda, T.; Kikuchi, H.; Ikeda, K.; Yumoto, M.; Sato, Y. Effects of blood flow restriction on cerebral blood flow during a single arm-curl resistance exercise. *Int. J. KAATSU Train. Res.* **2010**, *6*, 9–12. [[CrossRef](#)]

191. Ganesan, G.; Cotter, J.A.; Reuland, W.; Cerussi, A.E.; Tromberg, B.J.; Galassetti, P. Effect of blood flow restriction on tissue oxygenation during knee extension. *Med. Sci. Sports Exerc.* **2015**, *47*, 185–193. [[CrossRef](#)] [[PubMed](#)]
192. Obrig, H.; Villringer, A. Beyond the visible—Imaging the human brain with light. *J. Cereb. Blood Flow Metab.* **2003**, *23*, 1–18. [[CrossRef](#)] [[PubMed](#)]
193. Lindauer, U.; Dirnagl, U.; Fuchtemeier, M.; Bottiger, C.; Offenhauser, N.; Leithner, C.; Rojl, G. Pathophysiological interference with neurovascular coupling—When imaging based on hemoglobin might go blind. *Front. Neuroenergetics* **2010**, *2*. [[CrossRef](#)] [[PubMed](#)]
194. Scholkmann, F.; Wolf, M. Measuring brain activity using functional near infrared spectroscopy: A short review. *Spectrosc. Eur.* **2012**, *24*, 6–10.
195. Aslin, R.N.; Shukla, M.; Emberson, L.L. Hemodynamic correlates of cognition in human infants. *Annu. Rev. Psychol.* **2015**, *66*, 349–379. [[CrossRef](#)] [[PubMed](#)]
196. Kirilina, E.; Jelzow, A.; Heine, A.; Niessing, M.; Wabnitz, H.; Brühl, R.; Ittermann, B.; Jacobs, A.M.; Tachtsidis, I. The physiological origin of task-evoked systemic artefacts in functional near infrared spectroscopy. *NeuroImage* **2012**, *61*, 70–81. [[CrossRef](#)] [[PubMed](#)]
197. Dravida, S.; Noah, J.A.; Zhang, X.; Hirsch, J. Comparison of oxyhemoglobin and deoxyhemoglobin signal reliability with and without global mean removal for digit manipulation motor tasks. *Neurophotonics* **2018**, *5*, 11006. [[CrossRef](#)] [[PubMed](#)]
198. Obrig, H.; Neufang, M.; Wenzel, R.; Kohl, M.; Steinbrink, J.; Einhäupl, K.; Villringer, A. Spontaneous low frequency oscillations of cerebral hemodynamics and metabolism in human adults. *NeuroImage* **2000**, *12*, 623–639. [[CrossRef](#)] [[PubMed](#)]
199. Kumar, V.; Shivakumar, V.; Chhabra, H.; Bose, A.; Venkatasubramanian, G.; Gangadhar, B.N. Functional near infra-red spectroscopy (fNIRS) in schizophrenia: A review. *Asian J. Psychiatr.* **2017**, *27*, 18–31. [[CrossRef](#)] [[PubMed](#)]
200. Haeussinger, F.B.; Dresler, T.; Heinzel, S.; Schecklmann, M.; Fallgatter, A.J.; Ehlis, A.-C. Reconstructing functional near-infrared spectroscopy (fNIRS) signals impaired by extra-cranial confounds: An easy-to-use filter method. *NeuroImage* **2014**, *95*, 69–79. [[CrossRef](#)] [[PubMed](#)]
201. Heinzel, S.; Haeussinger, F.B.; Hahn, T.; Ehlis, A.-C.; Plichta, M.M.; Fallgatter, A.J. Variability of (functional) hemodynamics as measured with simultaneous fNIRS and fMRI during intertemporal choice. *NeuroImage* **2013**, *71*, 125–134. [[CrossRef](#)] [[PubMed](#)]
202. Byun, K.; Hyodo, K.; Suwabe, K.; Ochi, G.; Sakairi, Y.; Kato, M.; Dan, I.; Soya, H. Positive effect of acute mild exercise on executive function via arousal-related prefrontal activations: An fNIRS study. *NeuroImage* **2014**, *98*, 336–345. [[CrossRef](#)] [[PubMed](#)]
203. Hyodo, K.; Dan, I.; Suwabe, K.; Kyutoku, Y.; Yamada, Y.; Akahori, M.; Byun, K.; Kato, M.; Soya, H. Acute moderate exercise enhances compensatory brain activation in older adults. *Neurobiol. Aging* **2012**, *33*, 2621–2632. [[CrossRef](#)] [[PubMed](#)]
204. Yamazaki, Y.; Sato, D.; Yamashiro, K.; Tsubaki, A.; Yamaguchi, Y.; Takehara, N.; Maruyama, A. Inter-individual differences in exercise-induced spatial working memory improvement: A near-infrared spectroscopy study. *Adv. Exp. Med. Biol.* **2017**, *977*, 81–88. [[CrossRef](#)] [[PubMed](#)]
205. Barha, C.K.; Liu-Ambrose, T. Exercise and the aging brain: Considerations for sex differences. *BPL* **2018**, *3*, 1–11. [[CrossRef](#)]
206. Loprinzi, P.D.; Frith, E. The role of sex in memory function: Considerations and recommendations in the context of exercise. *J. Clin. Med.* **2018**, *7*. [[CrossRef](#)] [[PubMed](#)]
207. Jones, N.; Kiely, J.; Suraci, B.; Collins, D.; de Lorenzo, D.; Pickering, C.; Grimaldi, K. A genetic-based algorithm for personalized resistance-training. *Biol. Sport* **2016**, *33*, 117–126. [[CrossRef](#)] [[PubMed](#)]
208. Ratamess, N.; Alvar, B.A.; Evetoch, T.K.; Housh, T.J.; Kibler, W.B.; Kraemer, W.J.; Triplett, N.T. American College of Sports Medicine position stand. Progression models in resistance training for healthy adults. *Med. Sci. Sports Exerc.* **2009**, *41*, 687–708. [[CrossRef](#)]
209. Toigo, M.; Boutellier, U. New fundamental resistance exercise determinants of molecular and cellular muscle adaptations. *Eur. J. Appl. Physiol.* **2006**, *97*, 643–663. [[CrossRef](#)] [[PubMed](#)]
210. Hughes, L.; Jeffries, O.; Waldron, M.; Rosenblatt, B.; Gissane, C.; Paton, B.; Patterson, S.D. Influence and reliability of lower-limb arterial occlusion pressure at different body positions. *PeerJ* **2018**, *6*, e4697. [[CrossRef](#)] [[PubMed](#)]

211. Loenneke, J.P.; Fahs, C.A.; Rossow, L.M.; Thiebaud, R.S.; Mattocks, K.T.; Abe, T.; Bemben, M.G. Blood flow restriction pressure recommendations: A tale of two cuffs. *Front. Physiol.* **2013**, *4*, 249. [[CrossRef](#)] [[PubMed](#)]
212. Sieljacks, P.; Knudsen, L.; Wernbom, M.; Vissing, K. Body position influences arterial occlusion pressure: Implications for the standardization of pressure during blood flow restricted exercise. *Eur. J. Appl. Physiol.* **2018**, *118*, 303–312. [[CrossRef](#)] [[PubMed](#)]
213. Brown, H.; Binnie, M.J.; Dawson, B.; Bullock, N.; Scott, B.R.; Peeling, P. Factors affecting occlusion pressure and ischemic preconditioning. *Eur. J. Sport Sci.* **2018**, *18*, 387–396. [[CrossRef](#)] [[PubMed](#)]
214. Loenneke, J.P.; Fahs, C.A.; Rossow, L.M.; Sherk, V.D.; Thiebaud, R.S.; Abe, T.; Bemben, D.A.; Bemben, M.G. Effects of cuff width on arterial occlusion: Implications for blood flow restricted exercise. *Eur. J. Appl. Physiol.* **2012**, *112*, 2903–2912. [[CrossRef](#)] [[PubMed](#)]
215. Mouser, J.G.; Dankel, S.J.; Jessee, M.B.; Mattocks, K.T.; Buckner, S.L.; Counts, B.R.; Loenneke, J.P. A tale of three cuffs: The hemodynamics of blood flow restriction. *Eur. J. Appl. Physiol.* **2017**, *117*, 1493–1499. [[CrossRef](#)] [[PubMed](#)]
216. Vanwyke, W.R.; Weatherholt, A.M.; Mikesky, A.E. Blood flow restriction training: Implementation into clinical practice. *Int. J. Exerc. Sci.* **2017**, *10*, 649–654. [[PubMed](#)]
217. Thurston, B.; Burr, J.F. Blood flow restricted training: Applications, mechanisms, and future directions. *Health Fitness J. Can.* **2017**, *10*, 13–16.
218. Jessee, M.B.; Buckner, S.L.; Dankel, S.J.; Counts, B.R.; Abe, T.; Loenneke, J.P. The influence of cuff width, sex, and race on arterial occlusion: Implications for blood flow restriction research. *Sports Med.* **2016**, *46*, 913–921. [[CrossRef](#)] [[PubMed](#)]
219. Laurentino, G.C.; Loenneke, J.P.; Teixeira, E.L.; Nakajima, E.; Iared, W.; Tricoli, V. The effect of cuff width on muscle adaptations after blood flow restriction training. *Med. Sci. Sports Exerc.* **2016**, *48*, 920–925. [[CrossRef](#)] [[PubMed](#)]
220. Mittal, P.; Shenoy, S.; Sandhu, J.S. Effect of different cuff widths on the motor nerve conduction of the median nerve: An experimental study. *J. Orthop. Surg. Res.* **2008**, *3*, 1. [[CrossRef](#)] [[PubMed](#)]
221. Mouser, J.G.; Dankel, S.J.; Mattocks, K.T.; Jessee, M.B.; Buckner, S.L.; Abe, T.; Loenneke, J.P. Blood flow restriction and cuff width: Effect on blood flow in the legs. *Clin. Physiol. Funct. Imaging* **2018**. [[CrossRef](#)] [[PubMed](#)]
222. Mattocks, K.T.; Jessee, M.B.; Mouser, J.G.; Dankel, S.J.; Buckner, S.L.; Bell, Z.W.; Owens, J.G.; Abe, T.; Loenneke, J.P. The application of blood flow restriction: Lessons from the laboratory. *Curr. Sports Med. Rep.* **2018**, *17*, 129–134. [[CrossRef](#)] [[PubMed](#)]
223. Buckner, S.L.; Dankel, S.J.; Counts, B.R.; Jessee, M.B.; Mouser, J.G.; Mattocks, K.T.; Laurentino, G.C.; Abe, T.; Loenneke, J.P. Influence of cuff material on blood flow restriction stimulus in the upper body. *J. Physiol. Sci.* **2017**, *67*, 207–215. [[CrossRef](#)] [[PubMed](#)]
224. Loenneke, J.P.; Thiebaud, R.S.; Fahs, C.A.; Rossow, L.M.; Abe, T.; Bemben, M.G. Effect of cuff type on arterial occlusion. *Clin. Physiol. Funct. Imaging* **2013**, *33*, 325–327. [[CrossRef](#)] [[PubMed](#)]
225. Loenneke, J.P.; Allen, K.M.; Mouser, J.G.; Thiebaud, R.S.; Kim, D.; Abe, T.; Bemben, M.G. Blood flow restriction in the upper and lower limbs is predicted by limb circumference and systolic blood pressure. *Eur. J. Appl. Physiol.* **2015**, *115*, 397–405. [[CrossRef](#)] [[PubMed](#)]
226. Brandner, C.R.; Kidgell, D.J.; Warmington, S.A. Unilateral bicep curl hemodynamics: Low-pressure continuous vs high-pressure intermittent blood flow restriction. *Scand. J. Med. Sci. Sports* **2015**, *25*, 770–777. [[CrossRef](#)] [[PubMed](#)]
227. Counts, B.R.; Dankel, S.J.; Barnett, B.E.; Kim, D.; Mouser, J.G.; Allen, K.M.; Thiebaud, R.S.; Abe, T.; Bemben, M.G.; Loenneke, J.P. Influence of relative blood flow restriction pressure on muscle activation and muscle adaptation. *Muscle Nerve* **2016**, *53*, 438–445. [[CrossRef](#)] [[PubMed](#)]
228. Dankel, S.J.; Jessee, M.B.; Buckner, S.L.; Mouser, J.G.; Mattocks, K.T.; Loenneke, J.P. Are higher blood flow restriction pressures more beneficial when lower loads are used? *Psychophysiol. Int.* **2017**, *104*, 247–257. [[CrossRef](#)] [[PubMed](#)]
229. Fatela, P.; Reis, J.F.; Mendonca, G.V.; Avela, J.; Mil-Homens, P. Acute effects of exercise under different levels of blood-flow restriction on muscle activation and fatigue. *Eur. J. Appl. Physiol.* **2016**, *116*, 985–995. [[CrossRef](#)] [[PubMed](#)]

230. Jessee, M.B.; Buckner, S.L.; Mouser, J.G.; Mattocks, K.T.; Loenneke, J.P. Letter to the editor: Applying the blood flow restriction pressure: The elephant in the room. *Am. J. Physiol.* **2016**, *310*, H132–H133. [[CrossRef](#)] [[PubMed](#)]
231. Karabulut, M.; Leal, J.A., Jr.; Garcia, S.D.; Cavazos, C.; Bemben, M. Tissue oxygenation, strength and lactate response to different blood flow restrictive pressures. *Clin. Physiol. Funct. Imaging* **2014**, *34*, 263–269. [[CrossRef](#)] [[PubMed](#)]
232. Loenneke, J.P.; Kim, D.; Mouser, J.G.; Allen, K.M.; Thiebaud, R.S.; Abe, T.; Bemben, M.G. Are there perceptual differences to varying levels of blood flow restriction? *Physiol. Behav.* **2016**, *157*, 277–280. [[CrossRef](#)] [[PubMed](#)]
233. Barnett, B.E.; Dankel, S.J.; Counts, B.R.; Nooe, A.L.; Abe, T.; Loenneke, J.P. Blood flow occlusion pressure at rest and immediately after a bout of low load exercise. *Clin. Physiol. Funct. Imaging* **2016**, *36*, 436–440. [[CrossRef](#)] [[PubMed](#)]
234. Loenneke, J.P.; Kim, D.; Fahs, C.A.; Thiebaud, R.S.; Abe, T.; Larson, R.D.; Bemben, D.A.; Bemben, M.G. Effects of exercise with and without different degrees of blood flow restriction on torque and muscle activation. *Muscle Nerve* **2015**, *51*, 713–721. [[CrossRef](#)] [[PubMed](#)]
235. Loenneke, J.P.; Kim, D.; Fahs, C.A.; Thiebaud, R.S.; Abe, T.; Larson, R.D.; Bemben, D.A.; Bemben, M.G. The effects of resistance exercise with and without different degrees of blood-flow restriction on perceptual responses. *J. Sports Sci.* **2015**, *33*, 1472–1479. [[CrossRef](#)] [[PubMed](#)]
236. Fitschen, P.J.; Kistler, B.M.; Jeong, J.H.; Chung, H.R.; Wu, P.T.; Walsh, M.J.; Wilund, K.R. Perceptual effects and efficacy of intermittent or continuous blood flow restriction resistance training. *Clin. Physiol. Funct. Imaging* **2014**, *34*, 356–363. [[CrossRef](#)] [[PubMed](#)]
237. Neto, G.R.; Novaes, J.S.; Salerno, V.P.; Gonçalves, M.M.; Piazero, B.K.L.; Rodrigues-Rodrigues, T.; Cirilo-Sousa, M.S. Acute effects of resistance exercise with continuous and intermittent blood flow restriction on hemodynamic measurements and perceived exertion. *Percept. Mot. Skills* **2016**. [[CrossRef](#)] [[PubMed](#)]
238. Yasuda, T.; Loenneke, J.P.; Ogasawara, R.; Abe, T. Influence of continuous or intermittent blood flow restriction on muscle activation during low-intensity multiple sets of resistance exercise. *Acta Physiol. Hung.* **2013**, *100*, 419–426. [[CrossRef](#)]
239. Hughes, L.; Rosenblatt, B.; Gissane, C.; Paton, B.; Patterson, S.D. Interface pressure, perceptual, and mean arterial pressure responses to different blood flow restriction systems. *Scand. J. Med. Sci. Sports* **2018**, *28*, 1757–1765. [[CrossRef](#)] [[PubMed](#)]
240. McEwen, J.A.; Owens, J.G.; Jeyasurya, J. Why is it crucial to use personalized occlusion pressures in blood flow restriction (BFR) rehabilitation? *J. Med. Biol. Eng.* **2018**, *99*, 235. [[CrossRef](#)]
241. Day, B. Personalized Blood Flow Restriction Therapy: How, When and Where Can It Accelerate Rehabilitation After Surgery? *Arthroscopy* **2018**, *34*, 2511–2513. [[CrossRef](#)] [[PubMed](#)]
242. Lixandrão, M.E.; Ugrinowitsch, C.; Laurentino, G.; Libardi, C.A.; Aihara, A.Y.; Cardoso, F.N.; Tricoli, V.; Roschel, H. Effects of exercise intensity and occlusion pressure after 12 weeks of resistance training with blood-flow restriction. *Eur. J. Appl. Physiol.* **2015**, *115*, 2471–2480. [[CrossRef](#)] [[PubMed](#)]
243. Loenneke, J.P.; Thiebaud, R.S.; Abe, T.; Bemben, M.G. Blood flow restriction pressure recommendations: The hormesis hypothesis. *Medical Hypotheses* **2014**, *82*, 623–626. [[CrossRef](#)] [[PubMed](#)]
244. Patterson, S.D.; Hughes, L.; Head, P.; Warmington, S.; Brandner, C. Blood flow restriction training: A novel approach to augment clinical rehabilitation: How to do it. *Br. J. Sports Med.* **2017**, *51*, 1648–1649. [[CrossRef](#)] [[PubMed](#)]
245. Pope, Z.K.; Willardson, J.M.; Schoenfeld, B.J. Exercise and blood flow restriction. *J. Strength Cond. Res.* **2013**, *27*, 2914–2926. [[CrossRef](#)] [[PubMed](#)]
246. Fahs, C.A.; Loenneke, J.P.; Rossow, L.M.; Tiebaud, R.S.; Bemben, M.G. Methodological considerations for blood flow restricted resistance exercise. *J. Trainol.* **2012**, *1*, 14–22. [[CrossRef](#)]
247. Loenneke, J.P.; Wilson, J.M.; Marin, P.J.; Zourdos, M.C.; Bemben, M.G. Low intensity blood flow restriction training: A meta-analysis. *Eur. J. Appl. Physiol.* **2012**, *112*, 1849–1859. [[CrossRef](#)] [[PubMed](#)]
248. Radaelli, R.; Fleck, S.J.; Leite, T.; Leite, R.D.; Pinto, R.S.; Fernandes, L.; Simão, R. Dose-response of 1, 3, and 5 sets of resistance exercise on strength, local muscular endurance, and hypertrophy. *J. Strength Cond. Res.* **2015**, *29*, 1349–1358. [[CrossRef](#)] [[PubMed](#)]
249. Loenneke, J.P.; Wilson, J.M.; Wilson, G.J.; Pujol, T.J.; Bemben, M.G. Potential safety issues with blood flow restriction training. *Scand. J. Med. Sci. Sports* **2011**, *21*, 510–518. [[CrossRef](#)] [[PubMed](#)]



250. Patterson, S.D.; Brandner, C.R. The role of blood flow restriction training for applied practitioners: A questionnaire-based survey. *J. Sports Sci.* **2018**, *36*, 123–130. [[CrossRef](#)] [[PubMed](#)]
251. Loenneke, J.P.; Fahs, C.A.; Abe, T.; Rossow, L.M.; Ozaki, H.; Pujol, T.J.; Bemben, M.G. Hypertension risk: Exercise is medicine for most but not all. *Clin. Physiol. Funct. Imaging* **2014**, *34*, 77–81. [[CrossRef](#)] [[PubMed](#)]
252. Kacin, A.; Rosenblatt, B.; Grapar Zagri, T.; Biswas, A. Safety considerations with blood flow restricted resistance training. *Ann. Kinesiol.* **2015**, *6*, 3–26.
253. Lövdén, M.; Wenger, E.; Mårtensson, J.; Lindenberger, U.; Bäckman, L. Structural brain plasticity in adult learning and development. *Neurosci. Biobehav. Rev.* **2013**, *37*, 2296–2310. [[CrossRef](#)] [[PubMed](#)]
254. Rooks, C.R.; Thom, N.J.; McCully, K.K.; Dishman, R.K. Effects of incremental exercise on cerebral oxygenation measured by near-infrared spectroscopy: A systematic review. *Prog. Neurobiol.* **2010**, *92*, 134–150. [[CrossRef](#)] [[PubMed](#)]
255. Kenville, R.; Maudrich, T.; Carius, D.; Ragert, P. Hemodynamic Response Alterations in Sensorimotor Areas as a Function of Barbell Load Levels during Squatting: An fNIRS Study. *Front. Hum. Neurosci.* **2017**, *11*, 268. [[CrossRef](#)] [[PubMed](#)]
256. Park, J.L.; Fairweather, M.M.; Donaldson, D.I. Making the case for mobile cognition: EEG and sports performance. *Neurosci. Biobehav. Rev.* **2015**, *52*, 117–130. [[CrossRef](#)] [[PubMed](#)]
257. Tsujii, T.; Komatsu, K.; Sakatani, K. Acute effects of physical exercise on prefrontal cortex activity in older adults: A functional near-infrared spectroscopy study. *Adv. Exp. Med. Biol.* **2013**, *765*, 293–298. [[CrossRef](#)]
258. Moriya, M.; Aoki, C.; Sakatani, K. Effects of Physical Exercise on Working Memory and Prefrontal Cortex Function in Post-Stroke Patients. *Adv. Exp. Med. Biol.* **2016**, *923*, 203–208. [[CrossRef](#)] [[PubMed](#)]
259. Hsieh, S.-S.; Chang, Y.-K.; Hung, T.-M.; Fang, C.-L. The effects of acute resistance exercise on young and older males' working memory. *Psychol. Sport Exerc.* **2016**, *22*, 286–293. [[CrossRef](#)]
260. McMorris, T. *Exercise–Cognition Interaction: State of the Art and Future Research*; Elsevier: Amsterdam, The Netherlands, 2016; pp. 459–481.
261. Svebak, S.; Murgatroyd, S. Metamotivational dominance: A multimethod validation of reversal theory constructs. *J. Pers. Soc. Psychol.* **1985**, *48*, 107–116. [[CrossRef](#)]
262. Borg, G.A. Psychophysical bases of perceived exertion. *Med. Sci. Sports Exerc.* **1982**, *14*, 377–381. [[CrossRef](#)] [[PubMed](#)]
263. Tsukamoto, H.; Suga, T.; Takenaka, S.; Takeuchi, T.; Tanaka, D.; Hamaoka, T.; Hashimoto, T.; Isaka, T. An acute bout of localized resistance exercise can rapidly improve inhibitory control. *PLoS ONE* **2017**, *12*, e0184075. [[CrossRef](#)] [[PubMed](#)]
264. Tsukamoto, H.; Takenaka, S.; Suga, T.; Tanaka, D.; Takeuchi, T.; Hamaoka, T.; Isaka, T.; Hashimoto, T. Impact of exercise intensity and duration on post exercise executive function. *Med. Sci. Sports Exerc.* **2016**. [[CrossRef](#)]
265. Tsukamoto, H.; Suga, T.; Takenaka, S.; Tanaka, D.; Takeuchi, T.; Hamaoka, T.; Isaka, T.; Hashimoto, T. Greater impact of acute high-intensity interval exercise on post-exercise executive function compared to moderate-intensity continuous exercise. *Physiol. Behav.* **2016**, *155*, 224–230. [[CrossRef](#)] [[PubMed](#)]
266. Lins, L.; Carvalho, F.M. SF-36 total score as a single measure of health-related quality of life: Scoping review. *SAGE Open Med.* **2016**, *4*. [[CrossRef](#)] [[PubMed](#)]
267. Buysse, D.J.; Reynolds, C.F.; Monk, T.H.; Berman, S.R.; Kupfer, D.J. The Pittsburgh Sleep Quality Index: A new instrument for psychiatric practice and research. *Psychiatry Res.* **1989**, *28*, 193–213. [[CrossRef](#)]
268. Kovacevic, A.; Mavros, Y.; Heisz, J.J.; Fiatarone Singh, M.A. The effect of resistance exercise on sleep: A systematic review of randomized controlled trials. *Sleep Med. Rev.* **2018**, *39*, 52–68. [[CrossRef](#)] [[PubMed](#)]
269. Kato, K.; Iwamoto, K.; Kawano, N.; Noda, Y.; Ozaki, N.; Noda, A. Differential effects of physical activity and sleep duration on cognitive function in young adults. *J. Sport Health Sci.* **2017**. [[CrossRef](#)]
270. Hughes, L.; Paton, B.; Rosenblatt, B.; Gissane, C.; Patterson, S.D. Blood flow restriction training in clinical musculoskeletal rehabilitation: A systematic review and meta-analysis. *Br. J. Sports Med.* **2017**, *51*, 1003–1011. [[CrossRef](#)] [[PubMed](#)]
271. Dankel, S.J.; Jessee, M.B.; Abe, T.; Loenneke, J.P. The Effects of Blood Flow Restriction on Upper-Body Musculature Located Distal and Proximal to Applied Pressure. *Sports Med.* **2016**, *46*, 23–33. [[CrossRef](#)] [[PubMed](#)]

272. Lixandrão, M.E.; Ugrinowitsch, C.; Berton, R.; Vechin, F.C.; Conceição, M.S.; Damas, F.; Libardi, C.A.; Roschel, H. Magnitude of muscle strength and mass adaptations between high-load resistance training versus low-load resistance training associated with blood-flow restriction: A systematic review and meta-analysis. *Sports Med.* **2018**, *48*, 361–378. [[CrossRef](#)] [[PubMed](#)]
273. Karabulut, M.; Abe, T.; Sato, Y.; Bemben, M.G. The effects of low-intensity resistance training with vascular restriction on leg muscle strength in older men. *Eur. J. Appl. Physiol.* **2010**, *108*, 147–155. [[CrossRef](#)] [[PubMed](#)]
274. Mavros, Y.; Gates, N.; Wilson, G.C.; Jain, N.; Meiklejohn, J.; Brodaty, H.; Wen, W.; Singh, N.; Baune, B.T.; Suo, C.; et al. Mediation of Cognitive Function Improvements by Strength Gains After Resistance Training in Older Adults with Mild Cognitive Impairment: Outcomes of the Study of Mental and Resistance Training. *J. Am. Geriatr. Soc.* **2017**, *65*, 550–559. [[CrossRef](#)] [[PubMed](#)]
275. Tumati, S.; Burger, H.; Martens, S.; van der Schouw, Y.T.; Aleman, A. Association between Cognition and Serum Insulin-Like Growth Factor-1 in Middle-Aged & Older Men: An 8 Year Follow-Up Study. *PLoS ONE* **2016**, *11*, e0154450. [[CrossRef](#)]
276. Loenneke, J.P.; Thiebaud, R.S.; Abe, T. Does blood flow restriction result in skeletal muscle damage? A critical review of available evidence. *Scand. J. Med. Sci. Sports* **2014**, *24*, e415–e422. [[CrossRef](#)] [[PubMed](#)]
277. Abe, T.; Beekley, M.D.; Hinata, S.; Koizumi, K.; Sato, Y. Day-to-day change in muscle strength and MRI-measured skeletal muscle size during 7 days KAATSU resistance training: A case study. *Int. J. KAATSU Train. Res.* **2005**, *1*, 71–76. [[CrossRef](#)]



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).