

SHORT COMMUNICATION

Plasma neurofilament light chain as a biomarker for fatal familial insomnia

Peter Hermann¹  | Sezgi Canaslan¹ | Anna Villar-Piqué^{2,3} | Timothy Bunck¹ | Stefan Goebel¹ | Franc Llorens^{1,2,3} | Matthias Schmitz¹ | Inga Zerr^{1,4}

¹Department of Neurology, National Reference Center for CJD Surveillance, Göttingen University Medical Center, Göttingen, Germany

²Bellvitge Biomedical Research Institute, Hospitalet de Llobregat, Spain

³Network Center for Biomedical Research in Neurodegenerative Diseases, Carlos III Institute of Health, Madrid, Spain

⁴German Center for Neurodegenerative Diseases, Göttingen campus, Göttingen, Germany

Correspondence

Peter Hermann, Department of Neurology, Göttingen University Medical Center, Georg August University, Robert Koch Straße 40, 37075 Göttingen, Germany.
Email: peter.hermann@med.uni-goettingen.de

Funding information

CJD Foundation; Alzheimer Forschung Initiative, Grant/Award Number: 20026; Robert-Koch-Institute, Grant/Award Number: 1369-341

Abstract

Background and purpose: Fatal familial insomnia is a rare hereditary prion disease associated with the D178N-129M *PRNP* mutation. Early diagnosis is difficult, because the clinical syndrome may overlap with affective disorders. In addition, most known cerebrospinal fluid biomarkers for prion diseases and magnetic resonance imaging do not show a good diagnostic accuracy for fatal familial insomnia. In this context, data on plasma biomarkers are scarce.

Methods: We analyzed levels of neurofilament light chain, glial fibrillary acidic protein, chitinase-3-like protein 1, calcium-binding protein B, and total tau protein in six serial plasma samples from a patient with fatal familial insomnia. Subsequently, plasma neurofilament light chain was analyzed in $n = 25$ patients and $n = 19$ controls. The diagnostic accuracy and associations with disease stage and duration were explored.

Results: Among all biomarker candidates in the case study, only neurofilament light chain levels showed a constant evolution and increased over time. They discriminated fatal familial insomnia from controls with an area under the curve of 0.992 (95% confidence interval [CI] = 0.974–1) in the case-control study. Higher concentrations were associated with methionine homozygosity at codon 129 *PRNP* ($p = 0.006$), shorter total disease duration ($\rho = -0.467$, $p = 0.019$, 95% CI = -0.790 to -0.015), and shorter time from sampling to death ($\rho = -0.467$, $p = 0.019$, 95% CI = -0.773 to -0.019).

Conclusions: Plasma neurofilament light chain may be a valuable minimally invasive diagnostic biomarker for fatal familial insomnia after clinical onset. Most important, stage-related increase and association with disease duration indicate potential as a prognostic marker and as a surrogate marker in clinical trials.

KEYWORDS

biomarker, fatal familial insomnia, neurofilament light chain, plasma, prion disease

INTRODUCTION

Fatal familial insomnia (FFI) is a rare hereditary prion disease associated with the D178N-129M *PRNP* mutation, primarily thalamic

pathologic changes, and fatal outcome. The survival time is influenced by the polymorphism of methionine/valine (M/V) at codon 129 *PRNP* [1]. Early FFI is characterized by sleep disorder, nonspecific psychiatric, and vegetative symptoms [2]. Thus, early diagnosis may

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2022 The Authors. *European Journal of Neurology* published by John Wiley & Sons Ltd on behalf of European Academy of Neurology.

be difficult, even in known mutation carriers. Unfortunately, cerebrospinal fluid (CSF) and imaging biomarkers for sporadic Creutzfeldt-Jakob disease (sCJD) [3] do not perform well in the diagnosis of FFI [4]. The only sensitive diagnostic marker is thalamic hypometabolism in positron emission tomography [5], a laborious method. Minimally invasive, blood-based biomarkers are needed for early diagnosis and trial monitoring. Candidates include markers for neuroaxonal injury such as total tau and neurofilament light chain (NfL), as well as glial markers. Some of these proteins were shown to be of diagnostic value when measured in plasma of patients with sCJD [6,7] and some genetic prion diseases [8]. Data regarding FFI are scarce.

We analyzed plasma samples from a patient with FFI at various time points before and after disease onset to demonstrate the evolution of candidate biomarker proteins. Subsequently, we analyzed plasma samples from a historical cohort of FFI patients to evaluate the diagnostic accuracy and potential prognostic utility of plasma NfL, a neuron-specific protein that is able to indicate neuroaxonal injury in neurodegenerative diseases [9].

METHODS AND MATERIALS

FFI and control patients

A patient with D178N *PRNP* mutation was followed from 3 years before to 8 months after FFI onset in the outpatient clinic of the Göttingen University Medical Center. Six consecutive plasma samples were analyzed for NfL, glial fibrillary acidic protein (GFAP), chitinase-3-like protein 1 (YKL-40), calcium-binding protein B (S100B), and total tau.

Subsequently, patients with prion disease and D178N *PRNP* mutation were selected from the biobank of the German National Reference Center for Transmissible Spongiform Encephalopathies when plasma samples and clinical data were available ($n = 28$), excluding the patient from the case study. Healthy carriers of D178N ($n = 5$), P102L ($n = 2$), and 120 base pair insert ($n = 2$) *PRNP* mutation (HMC), as well as patients with neuropsychiatric disorders (depression, $n = 7$; psychosis, $n = 1$, polyneuropathy, $n = 2$), in whom FFI or other neurodegenerative disorders had been considered during the diagnostic process (neuropsychiatric controls [NPC]), were assigned as controls.

Biomarker analyses

Plasma NfL, total tau, and GFAP levels were measured with SIMOA assays (Quanterix). MicroVueYKL-40 EIA ELISA kit from Quidel was used for quantification of YKL-40 and LIAISON SANGTEC 100 for S100B. Manufacturers' instructions were followed. Test performers were blind to clinical information and clinical investigators to test performance.

Statistical methods

Comparisons were performed with multivariate linear regression models, including age and sex, followed by Tukey contrasts for multiple comparisons. Data were normalized by log transformation. Receiver operating characteristic (ROC) curves and area under the curve (AUC) with 95% confidence intervals (CIs) were calculated. Associations of NfL levels were explored with Spearman correlation. SPSS and R multComp were used for analyses and Excel for graphic items. Statistical significance was considered at $p < 0.05$.

Ethical standards


The study was conducted according to the revised Declaration of Helsinki and Good Clinical Practice guidelines, approved by the Göttingen University Medical Center ethics committee (No. 11/11/93). All patients or their legal representatives gave informed consent.

RESULTS

Case report and evaluation of biomarker candidates

Longitudinal evaluation of plasma NfL, GFAP, YKL-40, S100B, and total tau in a D178N (codon 129: MM) mutation carrier revealed that only NfL concentrations showed a stage-related increase. We obtained plasma close to the exact point of clinical onset. The patient

FIGURE 1 Evolution of plasma biomarker candidates in a case of fatal familial insomnia (FFI). (a) Development of plasma biomarker levels (percentage) in relation to Time Point 1 (100%). Time Point 1: 3 years before onset; Time Point 2: 2 years before onset; Time Point 3: 1 year before onset; Time Point 4: clinical onset (very mild sleep disturbance); Time Point 5: 4 months after onset (insomnia, hyperhidrosis, depressive mood); Time Point 6: 8 months after onset (mild dementia, ataxia, upper limb rigor). GFAP, glial fibrillary acidic protein; NfL, neurofilament light chain; S100B, calcium-binding protein B; tau, total Tau protein; YKL-40, chitinase-3-like protein 1. (b) Box plots displaying NfL concentrations in FFI MM, FFI MV, and control patients. To improve readability, logarithmic scale was chosen for the y-axis. HMC, healthy *PRNP* mutation carriers; MM, methionine homozygosity at codon 129 *PRNP*; MV, methionine/valine heterozygosity at codon 129 *PRNP*; NPC, neuropsychiatric controls. Bars and p -values above boxes indicate p -values from linear regression and Tukey contrasts. (c) Receiver operation characteristic (ROC) curve and area under the curve (AUC) with 95% confidence intervals (CIs) in the discrimination of FFI patients and all controls (HMC plus NPC). (d) Scatter plot displaying association between plasma NfL (pg/ml) and time from disease onset to death (days). Spearman coefficients (ρ), 95% CIs, and corresponding p -values are indicated. (e) Scatter plot displaying association between plasma NfL (pg/ml) and time from blood uptake to death (days). Spearman coefficients (ρ), 95% CI, and corresponding p -values are indicated



visited the outpatient clinic for a routine appointment and reported very mild sleep disturbance during the prior week. No further symptoms were present. The NfL level was almost twice as high (9.8 pg/ml) as 1 year before (5.49 pg/ml). At follow-up (4 months later; the

patient showed insomnia, vegetative, and affective symptoms but no focal neurological signs or dementia), NfL levels had increased further (20.75 pg/ml) and peaked in the last sample (57.15 pg/ml) 8 months after onset (Figure 1a, Table S1).

Diagnostic accuracy of NfL and association with disease duration of FFI

From $n = 28$ patients with D178N PRNP mutation, $n = 25$ showed an FFI phenotype and three presented with a CJD syndrome. Among FFI cases, $n = 18$ were homozygous for methionine at codon 129 PRNP (MM, mean = 124.35 pg/ml, SD = ± 73.21) and $n = 7$ were heterozygous (MV, mean = 52.19 pg/ml, SD = ± 27.85). Patients with CJD syndrome showed lower NfL concentrations (63.49 pg/ml, SD = ± 30.63) than FFI patients (102.15 pg/ml, SD = ± 71.95) but were excluded from further analyses due to the low case number. Mean NfL levels were similar in HMC with D178N mutation (6.53 pg/ml, SD = ± 2.07), other PRNP mutations (7.28 pg/ml, SD = ± 6.81), and NPC (10.88 pg/ml, SD = ± 7.26 ; Table 1).

A pooled control group (HMC and NPC) was used for comparison models and ROC analyses. FFI-MV patients showed longer total disease duration ($p = 0.023$) and lower NfL concentrations ($p = 0.006$) than FFI-MM patients (Figure 1b). Both FFI groups showed higher NfL than controls ($p < 0.001$). ROC analyses displayed an AUC of 0.992 (95% CI = 0.974–1) in the discrimination of all FFI patients from controls (Figure 1c). Regarding subgroups, NfL discriminated FFI-MM with an AUC of 1 (95% CI = 1–1) and FFI-MV with an AUC of 0.970 (95% CI = 0.912–1) from controls.

Samples from patients in the first disease tertial showed low NfL mean concentrations (25.90 pg/ml, SD = ± 13.12) compared to the second (105.65 pg/ml, SD = ± 73.89) and third (118.14 pg/ml, SD = ± 74.66) tertials. Due to the low case number in the first group ($n = 2$), statistical comparisons were not performed (Table 1). Spearman calculations showed significant negative correlations between NfL and total disease duration ($\rho = -0.4670$, $p = 0.0186$, 95% CI = -0.790 to -0.015 ; Figure 1d), as well as time from blood sampling to death

($\rho = -0.4673$, $p = 0.0185$, 95% CI = -0.773 to -0.019) in FFI cases (Figure 1e), but no significant correlation between NfL and time from onset to blood sampling ($\rho = -0.214$, $p = 0.305$, 95% CI = -0.677 to 0.264) was observed.

DISCUSSION

Most prion disease CSF biomarkers and magnetic resonance imaging show low diagnostic accuracy in FFI [4]. In contrast, CSF NfL was reported to have good diagnostic accuracy for prion diseases with the D178N PRNP mutation [10] and is also elevated in sporadic fatal insomnia [11]. Previous studies on plasma biomarkers for prion diseases included only few individuals with FFI. Nonetheless, probable elevation of plasma NfL was reported in FFI [6,8] but not in healthy D178N PRNP mutation carriers [12]. In our FFI patient, NfL increased over time, whereas other biomarker candidates showed no clear tendency. Although, for example, total tau showed a decrease during the observation time, we did not consider this alteration as relevant, because the course offers no clear tendency and all concentrations were at a generally low level.

Our retrospective study showed a significant elevation of plasma NfL in symptomatic FFI patients. Although NfL was significantly higher in FFI-MM than in FFI-MV, an excellent discriminatory value versus controls could be observed for both groups, indicating a high potential value of plasma NfL as a diagnostic biomarker. Additionally, we validated a longer disease duration in FFI-MV patients compared to FFI-MM patients [1]. Higher NfL may be associated with faster disease progression in the FFI-MM group. Control groups with other encephalopathic syndromes were not available. However, compared to previous investigations, mean concentration in FFI (102.15 pg/ml)

TABLE 1 Demographic and NfL data in the study cohort

Group	<i>n</i>	Sex, female/male	Age, median (min–max)	Disease duration, median days (min–max)	Plasma NfL, mean pg/ml $\pm$ SD
Controls	19	13/6	59 (32–73)	–	8.97 $\pm$ 6.27
D178N PRNP mutation	5	4/1	49 (32–70)	–	6.53 $\pm$ 2.07
Other PRNP mutations	4	3/1	48 (36–60)	–	7.28 $\pm$ 6.81
Other disorders	10	6/4	63 (48–73)	–	10.88 $\pm$ 7.26
Cases [D178N-FFI]	25	9/16	53 (24–70)	381 (178–765)	102.15 $\pm$ 71.95
FFI-MM	18	7/11	51 (24–65)	343 (178–550) ^a	124.35 $\pm$ 73.21
FFI-MV	7	2/5	67 (43–70)	495 (285–765) ^a	52.19 $\pm$ 27.85
First disease tertial	2	1/1	– (52–70)	– (285–679)	25.90 $\pm$ 13.12
Second disease tertial	11	5/6	46 (24–69)	391 (226–765)	105.65 $\pm$ 73.89
Third disease tertial	12	3/9	57 (27–70)	370 (178–550)	118.14 $\pm$ 74.66
Cases [D178N-gCJD] ^b	3	1/2	57 (52–60)	654 (453–661)	63.49 $\pm$ 30.63

Note: Total disease duration was divided into three time spans (tertiars), and patients were assigned according to time of blood sampling.

Abbreviations: FFI, fatal familial insomnia; gCJD, genetic Creutzfeldt-Jakob disease; M, methionine at codon 129 PRNP; min–max, minimum–maximum; NfL, neurofilament light chain; V, valine at codon 129 PRNP.

^aComparison of median disease duration with Mann–Whitney *U* test showed longer duration in FFI-MV (*U*:25, *z*-score = -2.270 , $p = 0.023$).

^bIn gCJD, two patients had codon 129 MM genotype (NfL = 97.97 and 36.88 pg/ml) and one patient had codon 129 MV genotype (NfL = 56.63 pg/ml).

was lower than what had been shown for sCJD (349.7 pg/ml) and nonneurodegenerative encephalopathies (182.2 pg/ml). In contrast, it was two- to threefold higher than in Alzheimer disease (34.9 pg/ml) and other dementia syndromes [7].

Most strikingly, the current observation shows significant association of plasma NfL with FFI stage (time from sampling to death) and duration, in line with observations in other neurodegenerative disorders [13]. To our knowledge, we investigated the largest symptomatic cohort with D178N PRNP mutation regarding plasma biomarkers. On the other hand, the cohort was still too small to perform certain subgroup analyses. Only two samples in the retrospective study were from the first disease tertial. Those showed higher NfL concentrations than HMC but still lower levels than FFI patients in later stages. Unfortunately, the case study could not show apparent elevation of NfL prior to disease onset. Thus, fluorodeoxyglucose positron emission tomography may have superior sensitivity in pre-clinical and very early clinical stages [5].

As a marker of neuroaxonal damage, plasma NfL is elevated in various neurologic diseases [9] and its specificity depends on the diagnostic context. Regarding FFI, affective disorders that may cause behavioral changes or sleeping disturbance are important differential diagnoses in early disease stages. Our control group was chosen accordingly. Individuals from families with known genetic prion diseases have a high risk for such disorders, because they suffer from higher stress and anxiety levels than controls [14]. In this context, plasma NfL may be useful for diagnosis of disease manifestation or as an individual screening test before positron emission tomography is applied. Most importantly, the stage-related increase and the strong association with survival time corroborate the potential role of plasma NfL as an urgently needed surrogate marker [15] in clinical trials for genetic prion diseases. The prognostic potential of plasma NfL in FFI may be improved by consideration of the codon 129 genotype. Here, we performed an exploratory evaluation in a small cohort. Larger investigations using international cohorts are needed to characterize the biomarker further.

ACKNOWLEDGMENTS

This study was supported by the Robert Koch Institute through funds of the Federal Ministry of Health (grant 1369-341), the CJD Foundation (grant title: "Implementation of a Blood-Based Biomarker Test for Sporadic Creutzfeldt-Jakob in Clinical Practice"), and the Alzheimer Forschung Initiative (project 20026). The funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, and approval of the manuscript; and decision to submit the manuscript for publication. Open access funding enabled and organized by ProjektDEAL.

CONFLICT OF INTEREST

A.V.-P. is the recipient of a postdoctoral grant of the Beatriz de Pinós program (2018-BP-00129) from the Ministry of Business and Knowledge of the Government of Catalonia, cofunded by the EU Horizon 2020 program under an MSCA grant agreement (801370).

F.L. has received funding from the CJD Foundation (grant title: "Implementation of a Blood-Based Biomarker Test for Sporadic Creutzfeldt-Jakob in Clinical Practice") related to this work, and the Alzheimer's Drug Discovery Foundation (grant 201810-2017419), and is the recipient of research grants through the Carlos III Institute of Health from the Government of Spain (grants CP16/00041 and PI19/00144). M.S. has received funds from the Alzheimer Forschung Initiative (project 20026) for work related to this article. I.Z. has received funding from the CJD Foundation (grant title: "Implementation of a Blood-Based Biomarker Test for Sporadic Creutzfeldt-Jakob in Clinical Practice"), the Robert Koch Institute through funds from the German Federal Ministry of Health (grant 1369-341) related to this work, and the Alzheimer's Drug Discovery Foundation (grant 201810-2017419). None of the other authors has any conflict of interest to disclose.

AUTHOR CONTRIBUTIONS

Peter Hermann: Conceptualization (equal), formal analysis (lead), investigation (equal), writing—original draft (lead). **Sezgi Canaslan:** Investigation (equal), writing—review & editing (equal). **Anna Villar-Piqué:** Investigation (equal), writing—review & editing (equal). **Timothy Bunck:** Formal analysis (supporting), investigation (equal), writing—review & editing (equal). **Stefan Goebel:** Investigation (equal), writing—review & editing (equal). **Franc Llorens:** Investigation (equal), writing—review & editing (equal). **Matthias Schmitz:** Investigation (equal), writing—review & editing (equal). **Inga Zerr:** Conceptualization (equal), formal analysis (supporting), supervision (lead), writing—review & editing (equal).

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Peter Hermann  <https://orcid.org/0000-0003-1889-8325>

REFERENCES

1. Parchi P, Petersen RB, Chen SG, et al. Molecular pathology of fatal familial insomnia. *Brain Pathol.* 1998;8:539-548.
2. Krasnianski A, Sanchez Juan P, Ponto C, et al. A proposal of new diagnostic pathway for fatal familial insomnia. *J Neurol Neurosurg Psychiatry.* 2014;85:654-659.
3. Hermann P, Appleby B, Brandel JP, et al. Biomarkers and diagnostic guidelines for sporadic Creutzfeldt-Jakob disease. *Lancet Neurol.* 2021;20:235-246.
4. Schmitz M, Villar-Piqué A, Hermann P, et al. Diagnostic accuracy and prognostic utility of CSF biomarkers in genetic prion diseases. *Brain.* 2021. doi:10.1093/brain/awab350
5. Cortelli P, Perani D, Montagna P, et al. Pre-symptomatic diagnosis in fatal familial insomnia: serial neurophysiological and 18FDG-PET studies. *Brain.* 2006;29:668-675.
6. Abu-Rumeileh S, Baiardi S, Ladogana A, et al. Comparison between plasma and cerebrospinal fluid biomarkers for the early diagnosis and association with survival in prion disease. *J Neurol Neurosurg Psychiatry.* 2020;91:1181-1188.

7. Zerr I, Villar-Piqué A, Hermann P, et al. Diagnostic and prognostic value of plasma neurofilament light and total-tau in sporadic Creutzfeldt-Jakob disease. *Alzheimers Res Ther.* 2021;13:86.
8. Thompson AGB, Anastasiadis P, Druyeh R, et al. Evaluation of plasma tau and neurofilament light chain biomarkers in a 12-year clinical cohort of human prion diseases. *Mol Psychiatry.* 2021;26(10):5955-5966. doi:[10.1038/s41380-021-01045-w](https://doi.org/10.1038/s41380-021-01045-w)
9. Khalil M, Teunissen CE, Otto M, et al. Neurofilaments as biomarkers in neurological disorders. *Nat Rev Neurol.* 2018;14:577-589.
10. Zerr I, Schmitz M, Karch A, et al. Cerebrospinal fluid neurofilament light levels in neurodegenerative dementia: evaluation of diagnostic accuracy in the differential diagnosis of prion diseases. *Alzheimers Dement.* 2018;14:751-763.
11. Abu-Rumeileh S, Redaelli V, Baiardi S, et al. Sporadic fatal insomnia in Europe: phenotypic features and diagnostic challenges. *Ann Neurol.* 2018;84:347-360.
12. Vallabh SM, Minikel EV, Williams VJ, et al. Cerebrospinal fluid and plasma biomarkers in individuals at risk for genetic prion disease. *BMC Med.* 2020;18:140.
13. Mattsson N, Cullen NC, Andreasson U, Zetterberg H, Blennow K. Association between longitudinal plasma neurofilament light and neurodegeneration in patients with Alzheimer disease. *JAMA Neurol.* 2019;76:791-799.
14. Schwartz M, Brandel JP, Babonneau ML, et al. Genetic testing in prion disease: psychological consequences of the decisions to know or not to know. *Front Genet.* 2019;10:895.
15. Vallabh SM, Minikel EV, Schreiber SL, Lander ES. Towards a treatment for genetic prion disease: trials and biomarkers. *Lancet Neurol.* 2020;19:361-368.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Hermann P, Canaslan S, Villar-Piqué A, et al. Plasma neurofilament light chain as a biomarker for fatal familial insomnia. *Eur J Neurol.* 2022;29:1841-1846. doi:[10.1111/ene.15302](https://doi.org/10.1111/ene.15302)

MANAGE-PD

Tool for Making Informed Decisions to
Aid Timely Management of Parkinson's Disease



MANAGE-PD allows you to:

- Identify PD patients inadequately controlled on oral medications
- Determine which patients with PD may be adequately controlled on their current treatment regimen or may require changes to their treatment regimen



Scan the QR code to
access to the web

Click here to
access to the web



MANAGE-PD is an AbbVie Inc. registered Medical Device. It is a collaborative research and development effort between AbbVie Medical Affairs and Health Economics and Outcomes, the Parkinson's Foundation and an international panel of Movement Disorder Specialists.

©2022 AbbVie Inc. All rights reserved. The Parkinson's Foundation logo is the sole property of the Parkinson's Foundation used with written permission. Any use of the Parkinson's Foundation name or logo without Foundation permission is prohibited. All content in <https://www.managepd.eu/> is intended only for informational use by healthcare professionals and is not offered as or intended to be medical advice for any particular patient. This information is not intended for patients. Only a healthcare professional exercising independent clinical judgement can make decisions regarding appropriate patient care and treatment options considering the unique characteristics of each patient.

PD: Parkinson's Disease