

RESEARCH ARTICLE

Dopamine Pathway and Parkinson's Risk Variants Are Associated with Levodopa-Induced Dyskinesia

Yuri L. Soser, MD, PhD,^{1,2} Sara Bandres-Ciga, PhD,³  Bart Ferwerda, PhD,⁴ Maria T.P. Tocino, MSc,^{5,6} Diaz R. Belloso, MD,^{5,6} Pilar Gómez-Garre, PhD,^{5,6}  Johann Faouzi, PhD,^{7,8} Pille Taba, MD,⁹ Lukas Pavelka, MD,^{10,11} Tainà M. Marques, PhD,¹⁰ Clarissa P.C. Gomes, PhD,¹² Alexey Kolodkin, PhD,¹⁰ Patrick May, PhD,^{10,11}  Lukas M. Milanowski, MD, PhD,^{13,14} Zbigniew K. Wszolek, MD,¹⁴  Ryan J. Uitti, MD,¹⁴ Peter Heutink, PhD,¹⁵ Jacobus J. van Hilten, MD, PhD,¹⁶ David K. Simon, MD, PhD,¹⁷  Shirley Eberly, MSc,¹⁸ Ignacio Alvarez, MSc,¹⁹ Lynne Krohn, PhD,^{1,2} Eric Yu, MSc,^{1,2} Kathryn Freeman, BSc,^{1,2} Uladzislau Rudakou, MSc,^{1,2} Jennifer A. Ruskey, MSc,^{2,20} Farnaz Asayesh, MSc,^{2,20} Manuel Menéndez-González, MD,^{21,22,23} Pau Pastor, MD, PhD,^{19,24} Owen A. Ross, PhD,¹⁴  Reijo Krüger, MD,^{10,11,12} on behalf of the NCER-PD Consortium, Jean-Christophe Corvol, MD, PhD,⁸  Sulev Koks, MD, PhD,^{25,26}  Pablo Mir, MD, PhD,^{5,6,27}  Rob M.A. De Bie, MD, PhD,²⁸  Hirotaka Iwaki, MD, PhD,^{3,29,30}  Ziv Gan-Or, MD, PhD,^{1,2,20*}  on behalf of the International Parkinson's Disease Genomic Consortium,

¹Department of Human Genetics, McGill University, Montréal, Canada

²Department of Neurology, The Neuro (Montreal Neurological Institute-Hospital), McGill University, Montréal, Canada

³Department of Health and Human Services, Center for Alzheimer's and Related Dementias (CARD), National Institute on Aging and National Institute of Neurological Disorders and Stroke, National Institutes on Health, Bethesda, Maryland, USA

⁴Department of Clinical Epidemiology and Biostatistics, Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands

⁵Servicio de Neurología y Neurofisiología Clínica, Unidad de Trastornos del Movimiento, Instituto de Biomedicina de Sevilla, Hospital Universitario Virgen del Rocío/CSIC/Universidad de Sevilla, Sevilla, Spain

⁶Department of Neurobiology, Centro de Investigación Biomédica en Red sobre Enfermedades Neurodegenerativas (CIBERNED), Madrid, Spain

⁷Sorbonne Université, Paris Brain Institute—ICM, Inserm, CNRS, Assistance Publique Hôpitaux de Paris, Department of Neurology, Pitié-Salpêtrière Hospital, Paris, France

⁸Department of Economics and Statistics, CREST, ENSAI, Campus de Ker-Lann, Bruz Cedex, France

⁹Department of Neurology and Neurosurgery, Institute of Clinical Medicine, University of Tartu, Tartu, Estonia

¹⁰Transversal Translational Medicine, Luxembourg Institute of Health (LIH), Strassen, Luxembourg

¹¹Parkinson Research Clinic, Centre Hospitalier de Luxembourg (CHL), Strassen, Luxembourg

¹²Translational Neuroscience, Luxembourg Centre for Systems Biomedicine (LCSB), University of Luxembourg, Esch-sur-Alzette, Luxembourg

¹³Department of Neurology Faculty of Health Science, Medical University of Warsaw, Warsaw, Poland

¹⁴Department of Neurology, Mayo Clinic Florida, Jacksonville, Florida, USA

¹⁵German Center for Neurodegenerative Diseases (DZNE), Göttingen, Germany

¹⁶Department of Neurology, Leiden University Medical Center, Leiden, The Netherlands

¹⁷Department of Neurology, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, Massachusetts, USA

¹⁸Department of Biostatistics and Computational Biology, University of Rochester School of Medicine and Dentistry, Rochester, New York, USA

¹⁹Department of Neurology, Hospital Universitari Mutua de Terrassa, Barcelona, Spain

²⁰Department of Neurology and Neurosurgery, McGill University, Montreal, Canada

²¹Facultad de Medicina y Ciencias de la Salud, Universidad de Oviedo, Oviedo, Spain

²²Department of Neurology, Hospital Universitario Central de Asturias, Oviedo, Spain

²³Instituto de Investigación Sanitaria del Principado de Asturias, Oviedo, Spain

²⁴Unit of Neurodegenerative Diseases, Department of Neurology, University Hospital Germans Trias i Pujol and The Germans Trias i Pujol Research Institute (IGTP) Badalona, Barcelona, Spain

²⁵Centre for Molecular Medicine and Innovative Therapeutics, Murdoch University, Murdoch, Australia

²⁶Neurological and Translational Science, Perron Institute, Nedlands, Australia

²⁷Departamento de Medicina, Facultad de Medicina, Universidad de Sevilla, Sevilla, Spain

²⁸Department of Neurology and Clinical Neurophysiology, Amsterdam University Medical Centers, Amsterdam Neuroscience, University of Amsterdam, Amsterdam, The Netherlands

²⁹Laboratory of Neurogenetics, National Institute on Aging, National Institutes of Health, Bethesda, Maryland, USA

³⁰Data Tecnica International, Washington, District of Columbia, USA

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](#) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

*Correspondence to: Dr. Z. Gan-Or, Montreal Neurological Institute, McGill University, 1033 Pine Avenue, West, Ludmer Pavilion, Room 312, Montréal, QC, H3A 1A1, Canada; E-mail: ziv.gan-or@mcgill.ca

Relevant conflicts of interest/financial disclosures: The authors declare that there are no additional disclosures to report.

Funding agencies: This work was financially supported by The Michael J. Fox Foundation, Parkinson's Society Canada, the Canadian Consortium on Neurodegeneration in Aging (CCNA), and the Canada First Research Excellence Fund (CFREF), awarded to McGill University for the Healthy Brains for Healthy Lives (HBHL) program.

Received: 20 September 2023; **Revised:** 10 July 2024; **Accepted:** 15 July 2024

Published online 12 August 2024 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.29960

ABSTRACT: Background: Levodopa-induced dyskinesia (LID) is a common adverse effect of levodopa, one of the main therapeutics used to treat the motor symptoms of Parkinson's disease (PD). Previous evidence suggests a connection between LID and a disruption of the dopaminergic system as well as genes implicated in PD, including *GBA1* and *LRRK2*.

Objectives: Our goal was to investigate the effects of genetic variants on risk and time to LID.

Methods: We performed a genome-wide association study (GWAS) and analyses focused on *GBA1* and *LRRK2* variants. We also calculated polygenic risk scores (PRS) including risk variants for PD and variants in genes involved in the dopaminergic transmission pathway. To test the influence of genetics on LID risk we used logistic regression, and to examine its impact on time to LID we performed Cox regression including 1612 PD patients with and 3175 without LID.

Results: We found that *GBA1* variants were associated with LID risk (odds ratio [OR] = 1.65; 95% confidence

interval [CI], 1.21–2.26; $P = 0.0017$) and *LRRK2* variants with reduced time to LID onset (hazard ratio [HR] = 1.42; 95% CI, 1.09–1.84; $P = 0.0098$). The fourth quartile of the PD PRS was associated with increased LID risk ($OR_{\text{fourth_quartile}} = 1.27$; 95% CI, 1.03–1.56; $P = 0.0210$). The third and fourth dopamine pathway PRS quartiles were associated with a reduced time to development of LID ($HR_{\text{third_quartile}} = 1.38$; 95% CI, 1.07–1.79; $P = 0.0128$; $HR_{\text{fourth_quartile}} = 1.38$; 95% CI = 1.06–1.78; $P = 0.0147$).

Conclusions: This study suggests that variants implicated in PD and in the dopaminergic transmission pathway play a role in the risk/time to develop LID. Further studies will be necessary to examine how these findings can inform clinical care. © 2024 The Author(s). *Movement Disorders* published by Wiley Periodicals LLC on behalf of International Parkinson and Movement Disorder Society.

Key Words: levodopa-induced dyskinesia; Parkinson's disease; dopamine; *GBA1*; *LRRK2*

Levodopa is one of the most commonly administered therapies for Parkinson's disease (PD), particularly to treat motor symptoms.¹ However, as the disease progresses and patients are exposed to long-term levodopa therapy, a significant proportion develops levodopa-induced dyskinesia (LID), a debilitating side effect characterized by involuntary, uncontrolled, and often choreiform movements.² LID is estimated to affect ~40% to 50% of PD patients within 4 to 6 years of initiating levodopa therapy,^{3,4} however, a subset of them manifests LID also within the first year of the therapy,⁵ demonstrating the broad variability of LID risk and onset. The main pathophysiologic hypothesis behind LID development suggests that the presynaptic nigrostriatal degeneration, in combination with the brief half-life of levodopa, may produce an aberrant pulsatile stimulation of dopamine receptors.⁶ The loss of dopaminergic neurons results in reduced dopamine storage capacity, leading to a radical increase in dopamine release for each dose of levodopa.⁷ Other mechanisms have also been implicated in LID development, including the involvement of the glutamatergic, serotonergic, and noradrenergic neural circuits. In particular, as a result of the nigrostriatal degeneration, levodopa might be preferentially metabolized in serotonergic and noradrenergic terminals, which are unable to regulate effectively dopamine release because of the absence of D2 autoreceptors and appropriate dopamine transporters.^{8,9} Another parallel pathophysiologic hypothesis suggests that the development of LID

is not only a consequence of an altered dopaminergic transmission but also a reflection of aberrant plastic changes in the corticobasal ganglia system.¹⁰

Multiple environmental risk factors affecting LID have been identified, including levodopa dosage and duration of the therapy, use of dopamine agonists, PD age at onset (AAO), disease duration and severity, female sex and lower body mass index (BMI).^{11–15} Most of the suggested genetic risk factors for LID are related to the dopamine pathway, including genes encoding the dopamine receptors, especially *DRD2* and *DRD3*,^{16–18} the dopamine transporter *SLC6A3*,^{19,20} or enzymes that metabolize dopamine and are targeted by PD therapeutics,^{21,22} catechol-O-methyltransferase (*COMT*)^{23–25} and monoamine oxidases A and B (*MAOA*, *MAOB*).^{24–26} Interestingly, variants in *GBA1* and *LRRK2*, among the most frequent genetic risk factors for PD,^{27,28} have also been identified as potential risk factors for LID,^{29–36} and PD patients carrying *LRRK2* variants display preserved to enhanced serotonergic activity compared to idiopathic PD.³⁷ Carriers of *GBA1* and *LRRK2* variants show distinctive clinical presentations in PD, with *GBA1* variants being associated with a more rapidly progressive PD with earlier onset,³⁸ and *LRRK2* variants with an overall more benign disease course, but with also more frequent postural instability and gait difficulty as well as slightly earlier AAO compared to sporadic PD.³⁹ Other variants reported in LID include those in *BDNF*, involved in neural plasticity,^{40,41} *GRIN2A*, encoding a glutamatergic receptor,⁴² and *ADORA2A*, encoding the adenosine A2a receptor gene.⁴³ However, the association between LID

and most of the above-mentioned putative genetic risk factors is still controversial, with most findings reported deriving from candidate gene studies that failed to be confirmed in replication studies.⁴⁴⁻⁴⁹

Here, we aimed to systematically evaluate how genetics affect the risk and rate of progression to LID including a total of 4787 PD patients from multiple centers. For this purpose, we performed genome-wide association studies (GWAS) and downstream analyses focused on specific genes previously implicated in LID. In addition, we tested the effect produced by cumulative genetic risk on the occurrence and rate of progression to LID, including risk variants previously associated with PD and variants in genes involved in the dopaminergic transmission pathway.

Methods

Population

The study population included a total of 4787 PD patients, of which 1612 with and 3175 without LID (Supplementary Table S1). PD was diagnosed by movement disorder specialists according to the United Kingdom (UK) Brain Bank or International Parkinson Disease and Movement Disorders Society criteria.⁵⁰ In each participating center, LID diagnosis was made by a movement disorder specialist. We collected data about LID based on the specialist clinical evaluation in multiple appointments or the Unified Parkinson's Disease Rating Scale (UPDRS) part IV produced after clinical assessment of the patient. In the McGill cohort, LID was self-reported in a specific questionnaire where the patient was asked if they experienced LID symptoms. The participants were of European ancestry and their clinical and genetic data were collected from 15 different cohorts (Supplementary Table S1), 12 of which were from the International Parkinson's Disease Genomics Consortium (IPDGC) and three from the Accelerating Medicines Partnership Parkinson's Disease (AMP-PD, <https://amp-pd.org/>). The latter includes the Parkinson's Disease Biomarkers Program (PDBP), Parkinson's Progression Markers Initiative (PPMI), and Harvard Biomarker Study (HBS) cohorts. The cohorts were included in the different analyses depending on data availability. The cohorts included in each analysis are specified in Supplementary Table S2.

Genetic Analyses

The study design included two stages. First, we performed an analysis focused on the *GBA1* and *LRRK2* genes, which are known to affect the risk for PD and have been suggested to affect LID in previous studies. In the second stage of the analysis, we performed a GWAS to examine genome-wide potential associations with risk

and time to develop LID, followed by polygenic risk score (PRS) analyses as detailed below. Excluding the AMP-PD cohorts, with whole genome sequencing (WGS) data, the other centers (Supplementary Table S1) were genotyped using the OmniExpress, NeuroX,⁵¹ NeuroChip,⁵² or MegaChip GWAS array according to the manufacturer's instructions (Illumina). Quality control was performed following standard pipelines (detailed in <https://github.com/neurogenetics/GWAS-pipeline>) using Plink 1.9.⁵³ In brief, we filtered out heterozygosity outliers using an *F*-statistic cutoff of <-0.15 or >0.15 . Individuals with a variant call rate $<95\%$ and sex mismatch were excluded. Variants missing in $>5\%$ of the participants, with disparate missingness between cases and controls ($P < 1e-04$), or significantly deviating from Hardy-Weinberg equilibrium in controls ($P < 1e-04$) were also removed. We used genome-wide complex trait analysis to check for relatedness closer than first cousins between participants (genome-wide pairwise relatedness [PIHAT] >0.125). We performed imputation using the Michigan imputation server (<https://imputationserver.sph.umich.edu/index.html#>) with the Haplotype Reference Consortium reference panel r1.1 2016 under default settings. Ancestry outliers were detected using HapMap3 principal component analysis (PCA) data in R version 4.0.1.

After imputation, we selected variants with $r^2 > 0.8$ and a minor allele frequency >0.05 , while retaining common risk variants in the *GBA1* (p.N370S, p.E326K, and p.T369M) and *LRRK2* (p.G2019S and p.M1646T) regions, to perform specific analyses on these variants (detailed below). These genes were specifically selected given their importance in PD etiology^{27,28} and recent clinical trials⁵⁴ as well as their previously suggested association with LID.²⁹⁻³⁴ The carrier status of *GBA1*/*LRRK2* risk variants in individuals with and without LID is detailed in Supplementary Tables S3 and S4. Carriers of variants in the same gene were combined, so that the carrier status for *GBA1* and *LRRK2* refers to any aforementioned *GBA1* and *LRRK2* variants, respectively. To examine the association between the *GBA1* and *LRRK2* risk variant carrier status and LID occurrence we performed logistic regression, and to evaluate the association between the carrier status and time to LID onset we performed Cox regression using the R package "survival" (<https://cran.r-project.org/web/packages/survival/>). The time to LID variable included in the Cox regression was defined as the period between the start of levodopa therapy and LID onset, as previously done.⁵⁵ When LID did not manifest, this parameter was right-censored at the last follow-up. To minimize biases because of possible confounders and further harmonize the analyses across the centers, we adjusted the analyses by multiple covariates including principal components (PCs), PD AAO, sex, levodopa dosage, levodopa equivalent daily dose

(LEDD),^{56,57} dopamine agonist use, BMI, Hoehn and Yahr score (HY) and, exclusively for logistic regression, disease duration. For logistic regression analyses, we included the cumulative levodopa dosage and LEDD starting from the baseline (ie, levodopa initiation) to the last time point (ie, LID onset or last follow-up when LID was not present). In Cox regression, to avoid collinearity with the time to LID onset dependent variable, we replaced cumulative doses with doses at the last time point. All the covariates were selected using an Akaike Information Criterion (AIC)-based stepwise regression approach, which evaluated the model goodness of fit and selected the most appropriate covariates to include in the model. We performed the analyses separately in each cohort and then meta-analyzed the results using the R package “metafor” (<https://cran.r-project.org/web/packages/metafor/index.html>). Because variants in these genes have been previously associated with LID, we used a significance threshold of $\alpha = 0.05$.

Similar to the analyses on specific genes, to investigate the impact of common genetic variants on LID risk and time to onset we also performed GWAS with, respectively, logistic and Cox regression adjusted for the above-specified covariates. Cox regression was performed using the SurvivalGWAS_SV software (<https://www.liverpool.ac.uk/population-health/research/groups/statistical-genetics/survival-gwas-sv/>).⁵⁸ We conducted the analyses in each cohort separately, and then meta-analyzed the results using the METAL software (https://genome.sph.umich.edu/wiki/METAL_Documentation) with a fixed effects model weighted by β coefficients and the inverse of the standard errors.

PD Risk Variant-Based PRS

To assess the impact on LID of the cumulative genetic risk for PD we calculated PRS for each PD patient including the 90 variants associated with PD in the most recent GWAS meta-analysis in Europeans.⁵⁹ PRS calculation was performed based on the weighted allele dose as implemented in PRSice2 using default clumping (<https://choishingwan.github.io/PRSice/>).⁶⁰ To investigate the association between the PRS and LID risk we performed logistic regression, whereas to evaluate the association between PRS and progression to LID we performed Cox regression. The analyses were adjusted for PCs, PD AAO, sex, HY, and levodopa dosage, cumulative in logistic regression and at the last time point in Cox regression. The covariates were selected to minimize collinearity based on the results of the previous model goodness-of-fit analyses. These analyses were repeated using PRS as a continuous variable and then as a discrete variable by dividing the PRS into quartiles. For the analysis using PRS quartiles, we separately compared the association of individual membership to the second, third, and fourth quartiles versus the first quartile with LID risk/progression.

Dopamine Pathway PRS

To assess the impact of genes involved in the dopaminergic transmission pathway we also constructed a pathway PRS, or polygenic effect score (PES)⁶¹ using the PRSet feature of PRSice2 (https://choishingwan.github.io/PRSice/prset_detail/). Genes involved in this pathway were obtained from Explore the Molecular Signatures Database (MSigDB, version 2023.1), a collection of annotated gene sets for use with Gene Set Enrichment Analysis (GSEA) software (<https://www.gsea-msigdb.org/gsea/msigdb/>). These genes included *CDK5*, *FLOT1*, *PARK7*, *CHRNA2*, *ADORA2A*, *CRH*, *CRHBP*, *DRD1*, *DRD2*, *DRD3*, *DRD4*, *DRD5*, *TOR1A*, *RASD2*, *PNKD*, *GDNF*, *ARRB2*, *PRKN*, *PTGS2*, *RAB3B*, *PINK1*, *SLC6A2*, *SLC6A3*, *SLC6A4*, *SNCA*, *TH*, and *CNTNAP4* (detailed at http://www.gsea-msigdb.org/gsea/msigdb/human/geneset/GOBP_SYNAPTIC_TRANSMISSION_DOPAMINERGIC). To investigate the role of dopamine genes overall in LID development, we also performed PES analyses including genes more strongly related to multiple dopamine pathways (ie, transmission, transportation, synthesis, and metabolism, including *DRD1*, *DRD2*, *DRD3*, *DRD4*, *SLC6A3*, *ANKK1*, *VMAT*, *DDC*, *TH*, *COMT*, *MAOA*, and *MAOB*). To select the variants in each of the genes to include in the PES analyses we used the LID GWAS meta-analysis summary statistics from the current study, filtering variants with a *P*-value less than or equal to 0.05. In addition, we performed linkage disequilibrium (LD) clumping using the default $r^2 = 0.1$ and selecting variants within 250 Kb of distance from the pathway-related genes. A total of 1000 permutations were implemented to generate the empirical *P*-value corresponding to the optimized PES prediction of the dependent variable in the target cohort. We then calculated individual PES for each target cohort. To avoid potential inflation because of the presence of the target cohort in the meta-analysis summary statistics, each time we calculated the PES for a target cohort we excluded such cohort from the meta-analysis using a leave-one-out approach. To investigate the association between the dopamine pathway PES and LID risk we performed logistic regression, whereas to evaluate the association between the PES and progression to LID we performed Cox regression, as specified above for the PRS analyses.

Results

GBA1 and LRRK2 Variants Show Significant Associations with LID Risk and Time to LID

Analyses focusing on *GBA1* showed that *GBA1* variants were significantly associated with LID risk (odds ratio [OR] = 1.65; 95% confidence interval [CI], 1.21–2.26; $P = 0.0017$) (Fig. 1A). No association was found with time to LID (hazard ratio [HR] = 1.25; 95% CI, 0.99–1.58; $P = 0.0635$) (Fig. 1B). In contrast,




FIG. 1. (A,B) Association between *GBA1* variants and levodopa-induced dyskinesia (LID). The meta-analysis forest plot shows the coefficient (black squares) and 95% confidence interval (bars) of the analyses in each single cohort. The size of the square is proportional to the weight the cohort had on the overall meta-analysis, based on their single standard error. The black diamond at the bottom represents the overall coefficient and confidence interval. (A) Logistic regression between *GBA1* variants and LID risk. (B) Cox regression between *GBA1* variants and time to development of LID. FE, fixed effect model; AMP-PD, Accelerating Medicines Partnership Parkinson's disease, including the New Discovery of Biomarkers (BioFIND), the Harvard Biomarker Study (HBS) and the Parkinson's Disease Biomarkers Program (PDBP) cohorts; Barcelona: Hospital Universitari Mutua de Terrassa, Spain; Coriell, National Institute for Neurological Disorders and Stroke (NINDS) Exploratory Trials in PD Long-Term Study 1 (NET-PD LS1), Coriell Institute for Medical Research, USA; DIGPD, Drug Interaction With Genes in Parkinson's Disease, France; LEAP, Levodopa in Early Parkinson's Disease, Netherlands; Luxemburg, Luxembourg Centre for Systems Biomedicine; Mayo, Mayo Clinic, USA; McGill, McGill University, Canada; Oviedo, Central University Hospital of Asturias, Spain; PreCEPT, Parkinson Research Examination of CEP-1347 Trial; SCOPA, Scales for Outcomes in Parkinson's disease; Sevilla, Universidad de Sevilla; Tartu, University of Tartu.

LRRK2 variants showed no association with LID risk (OR = 1.18; 95% CI, 0.84–1.67; $P = 0.3484$) (Fig. 2A), but were significantly associated with reduced time to development of LID (HR = 1.42; 95% CI, 1.09–1.84; $P = 0.0098$) (Fig. 2B). The *GBA1* and *LRRK2* meta-analyses were not significant for genetic heterogeneity ($I^2 = 2.8\%$, $P = 0.4187$ and $I^2 = 0\%$, $P = 0.7069$ for the *GBA1* logistic and Cox regression, respectively; $I^2 = 0\%$, $P = 0.8315$ and $I^2 = 0\%$, $P = 0.4886$ for the *LRRK2* logistic and Cox regression, respectively), suggesting that heterogeneity does not drive these associations.

In the GWAS, genomic inflation was evaluated using quantile-quantile plots (Q-Q plots) and the lambda factor, showing no inflation and a slight deflation (lambda logistic regression = 0.9709, lambda Cox regression = 0.9555) (Supplementary Figs. S1 and S2). GWAS using both logistic and Cox regression showed no significant association with LID risk or time to development of LID, respectively (Supplementary Figs. S3 and S4). We further examined whether variants previously associated with LID in the literature^{16–20,23–26} and from the LIDPD website (<http://LiDpd.eurac.edu/>) showed associations in the current GWAS, but we found no significant results (Supplementary Tables S5 and S6). A recent GWAS in LID⁶² nominated significant signals in a progression GWAS meta-analysis. However, our study

failed to confirm these findings and the reported variants did not reach the nominal significance of 0.05 in our GWAS (Supplementary Table S7).

PD Risk Variant-Based PRS Is Associated with Increased Risk for LID

PRS analyses aggregating PD-associated variants showed that higher values of PRS were associated with a very mild increase in LID risk (OR = 1.02; 95% CI, 1.002–1.035; $P = 0.0298$) (Fig. 3B). When dividing the PRS in quartiles, logistic regression showed a significant association between the fourth quartile and LID, with a greater risk compared to the analyses using PRS as a continuous variable (OR_{fourth quartile} = 1.27; 95% CI, 1.03–1.56; $P = 0.0210$) (Fig. 3A, Supplementary Table S8). Cox regression did not show any significant associations between PRS and time to development of LID (Supplementary Fig. S5A,B, Supplementary Table S9). The PD PRS logistic regression was significant for a moderate heterogeneity ($I^2 = 43.90\%$, $P = 0.0449$) and repeating the meta-analysis using a random-effect model, which accounts for heterogeneity, the results did not show statistically significant associations (OR = 1.02, $P = 0.2038$). PD PRS Cox regression did not show heterogeneity ($I^2 = 0\%$, $P = 0.6236$).




FIG. 2. (A,B) Association between *LRRK2* variants and levodopa-induced dyskinesia (LID). (A) Logistic regression between *LRRK2* variants and LID risk. (B) Cox regression between *LRRK2* variants and time to development of LID.

Dopaminergic Transmission Pathway Polygenic Effect Score Is Associated with a Reduced Time to Development of LID

Analyses on the dopaminergic transmission pathway PES showed that higher values of PES were associated with a reduced time to development of LID (HR = 1.10; 95% CI, 1.02–1.18; $P = 0.0088$) (Fig. 4B). In addition, the third and fourth PES quartile were also associated with a reduced time to development of LID with a more elevated effect size compared to the analyses on PES as a continuous variable (HR_{third_quartile} = 1.38; 95%

CI, 1.07–1.79; $P = 0.0128$; HR_{fourth_quartile} = 1.38; 95% CI, 1.06–1.78; $P = 0.0147$) (Fig. 4A, Supplementary Table S11). Logistic regression did not show any statistically significant associations between dopaminergic transmission PES and LID risk (Supplementary Fig. S6A, B, Supplementary Table S10). To ensure that the results were not driven by the *SNCA* variants included in this pathway we also repeated the analyses excluding variants in this gene. We confirmed the previous findings for Cox regression both using the PES as a continuous variable (HR = 1.13; 95% CI, 1.04–1.23; $P = 0.0056$)




FIG. 3. (A,B) Logistic regression between polygenic risk scores (PRS) aggregating PD risk variants and levodopa-induced dyskinesia (LID) risk. (A) The plot shows the association between each PRS quartile and LID risk compared with the first quartile, meta-analyzing the results across the cohorts. The Y axis represents the PRS quartile, the X axis the odds ratio (red dot) and 95% CI (red bar). The presence of an asterisk indicates a significant association ($P < 0.05$). (B) The forest plot shows the association between PRS as a continuous variable and LID risk. CI, confidence interval. [Color figure can be viewed at wileyonlinelibrary.com]




FIG. 4. (A,B) Cox regression between the dopaminergic transmission pathway polygenic effect score (PES) and time to development of levodopa-induced dyskinesia (LID). (A) The plot shows the association between each PES quartile and time to development of LID compared with the first quartile, meta-analyzing the results across the cohorts. The Y axis represents the polygenic risk score (PRS) quartile, the X axis the hazard ratio (red dot) and 95% confidence interval (red bar). (B) The forest plot shows the association between PES as a continuous variable and time to development of LID. [Color figure can be viewed at wileyonlinelibrary.com]

(Supplementary Fig. S7B) and divided in quartiles ($HR_{\text{third_quartile}} = 1.32$; 95% CI, 1.01–1.73; $P = 0.0429$; $HR_{\text{fourth_quartile}} = 1.31$; 95% CI, 1.004–1.71; $P = 0.0465$) (Supplementary Fig. S7A, Supplementary Table S12). Logistic regression showed a significant association between the fourth quartile of the dopamine transmission pathway PES and LID risk ($OR_{\text{fourth_quartile}} = 1.33$; 95% CI, 1.04–1.72; $P = 0.0249$) (Supplementary Fig. S8A, Supplementary Table S12), which, however, did not emerge when treating the PES as a continuous variable ($OR = 1.01$; 95% CI, 0.97–1.06; $P = 0.6099$) (Supplementary Fig. S8B, Supplementary Table S12). Finally, Cox regression for the dopamine PES including multiple pathways did not show any statistically significant results (Supplementary Fig. S9A,B, Supplementary Table S13), whereas it was significantly associated with increased LID risk ($OR = 1.12$; 95% CI, 1.02–1.22; $P = 0.0142$) (Supplementary Fig. S10B). This association also emerged between the fourth PES quartile and LID risk ($OR_{\text{fourth_quartile}} = 1.63$; 95% CI, 1.003–2.65; $P = 0.0483$) (Supplementary Fig. S10A, Supplementary Table S13). The dopamine transmission PES meta-analyses were not significant for genetic heterogeneity ($I^2 = 30.8\%$, $P = 0.1450$ dopamine transmission PES logistic regression; $I^2 = 0\%$, $P = 0.7934$ dopamine transmission PES Cox regression), suggesting that heterogeneity is not the driver of these associations.

Discussion

In this study, we confirmed that *GBA1* variants were associated with increased risk for LID and

demonstrated that *LRRK2* variants were associated with a reduced time to development of LID from the initiation of levodopa treatment. Additionally, we found that PD PRS was associated with mildly increased risk for LID and that the dopaminergic transmission pathway PES is associated with a reduced time to development of LID.

Albeit some studies found contradictory results on the association between the *GBA1* and *LRRK2* variants and LID,^{44–47} many others have shown that these variants play a role in LID development,^{29–34} and in this study we also demonstrated that *LRRK2* variants might also affect the time to development of LID. The power for the association analyses was optimal (99%, as calculated on https://csg.sph.umich.edu/abecasis/cats/gas_power_calculator/ using the following parameters: cases = 1612, controls = 3175, significance threshold = $5e-8$, prevalence in the general population = 0.5, allele frequency = 0.1, and genotype relative risk [as inferred by the power calculator] = 1.3). The absence of significant signals in the risk and progression GWAS and, in general, the difficulty finding congruent results between different genetic studies investigating LID, as also reflected by the divergent results between the recent LID progression GWAS⁶² and our study, may be because of the stronger contribution in LID development of environmental factors, especially pharmacologic- (dosage of dopaminergic drugs, use of amantadine) and disease-related factors.^{11–15} Our results suggest that *GBA1* variants affect risk for LID, but not LID onset, whereas *LRRK2* variants affect LID onset, but not risk. These results should be replicated by additional

studies in other populations, and the potential mechanisms behind these observations should be studied experimentally.

The significant association between the two PRS analyses suggests that aggregating multiple common variants that might have a scarce effect on LID individually could contribute to uncovering the overall genetic impact on LID. In particular, the association between the PRS including PD risk variants suggests that patients with a stronger genetic risk profile for PD are also more at risk for LID, a factor to consider for patient counselling and potential clinical trials, although the magnitude of the increased risk was small.

We demonstrated that the dopaminergic synaptic transmission pathway PES was associated with an increased rate of LID development, which is in line with previous pathophysiologic hypotheses^{8,9,63} and studies suggesting an implication of dopamine pathway genes in the development of LID.^{16-20,23-26} When excluding the SNCA variants from the dopaminergic transmission PES analyses, we observe that the dopamine transmission PES conserves the significant association with time to LID, suggesting that this association is not driven by the SNCA variants. We also observed that the PES including dopamine genes is associated with a mild increase in LID risk, suggesting that, whereas dopamine transmission is more strongly implicated in LID onset, other genes related to dopamine might play a role in LID risk. Additional studies will be necessary to further investigate the association between the PD and dopamine PRS and LID.

Unravelling the etiologic bases of LID is crucial to implementing a tailored therapy for PD patients taking levodopa, adapting the therapeutic choices, dosage, and management depending on the individual risk factors of each patient. Over time, it could be beneficial to define a risk profile accounting for the single genetic and environmental factors associated with LID as well as the cumulative genetic risk provided by the PRS. This might be used to stratify patients for LID prevention clinical trials and lead to a more refined and personalized therapeutic approach for each individual. In addition to the benefits of the current symptomatic therapies, uncovering and confirming genetic factors affecting the risk and time of development of LID could also have important implications for targeted therapies. In particular, *GBA1* and *LRRK2* pathways are already candidate targets for newly developing drugs in clinical trials.⁵⁴ A *LRRK2* inhibitor, BIIB122/DNL151, reached already experimental phase 3 (<https://www.denalitherapeutics.com>, 2021).⁶⁴ In addition, ambroxol, a pharmacological chaperone for glucocerebrosidase (GCase) capable of increasing its enzymatic levels, completed phase 2 and LTI-291, an activator of GCase, reached phase 1B.⁶⁵⁻⁶⁷ Because these drugs would likely be used in conjunction with symptomatic therapies, knowing that these pathways can be targeted to reduce the risk or delay the time of LID development

could considerably improve the compliance and quality of life of PD patients taking dopaminergic treatments.

The current study has several limitations. First, the subjects were all of European ancestry and, therefore, the results in other populations might be different. Despite an overall large sample size, most of the individual cohorts included a limited number of participants, especially those having longitudinal data necessary for Cox regression, this impacted the power of the study and could have contributed to the lack of association in the GWAS. Similarly, in the analyses focused on *GBA1* some cohorts showed a small number of carriers, however, this did not substantially affect the results as demonstrated by the optimal statistical power. NeuroX and NeuroChip can be limited in the detection of *GBA1* variants, however, any potential errors would be present in both the case and control groups, producing arguably a minimal impact on the results. Some studies suggested that LID is influenced more by the disease duration than by the therapy duration,⁶⁸ on this line PD AAO would represent a better baseline than levodopa initiation for the time to LID onset. However, this parameter was chosen in accordance with what was previously done with LID GWAS⁵⁵ and accounting for the recall bias that PD AAO suffers from, compared to levodopa initiation, which represents a report made by the physicians. In addition, understanding the genetic basis of the time to LID from levodopa initiation can be of considerable relevance for patient counselling at the time of treatment administration. Finally, we also accounted for the disease duration in each of our analyses with appropriate adjustments. The association between the PD PRS and LID risk is very mild, therefore, the impact of this finding in the clinical practice might be limited. Additionally, we found a moderate heterogeneity in the PD PRS logistic regression meta-analysis. Therefore, the effect of the PRS on LID risk in the fixed effect model could be affected by this slight heterogeneity, which is supported by the lack of statistical significance in the random effect model. Another limitation of this study was that not all the cohorts had the same amount of data available, which limited in part the design of the analytical model. An interesting future line of research could be to investigate the influence of genetics on specific features of LID, such as LID subtypes (ie, peak-dose, wearing-off-state, and diphasic LID) and progression (ie, severity over time), by collecting these data in a large group of patients.

In conclusion, in the current study, we demonstrated that PD risk variants and the dopaminergic transmission PRS are associated with risk of LID/time to development of LID. A better understanding of the role of genetics in LID development could reduce the impact of this adverse effect and enhance therapeutic management in PD. ■

Acknowledgments: We wholeheartedly thank the participants in this study. We thank the research participants and all members of IPDGC for making this work possible. The AMP-PD cohort data used in this study included the Fox Investigation for New Discovery of Biomarkers (BioFIND), HBS, PDBP, and PPMI cohorts. For up-to-date information on the study, visit <https://www.amp-pd.org>. AMP PD—a public-private partnership—is managed by the FNIH and funded by Celgene, GSK, The Michael J. Fox Foundation for Parkinson's Research (MJFF), the National Institute of Neurological Disorders and Stroke, Pfizer, Sanofi, and Verily. BioFIND is sponsored by MJFF with support from the National Institute for Neurological Disorders and Stroke (NINDS). The BioFIND Investigators have not participated in reviewing the data analysis or content of the manuscript. For up-to-date information on the study, visit <https://www.michaeljfox.org/news/biofind>. HBS is a collaboration of HBS investigators (full list of HBS investigators found at <https://www.bwhparkinsoncenter.org/biobank/> and funded through philanthropy and National Institutes of Health [NIH] and non-NIH funding sources). The HBS Investigators have not participated in reviewing the data analysis or content of the manuscript. PDBP consortium is supported by NINDS at the NIH. A full list of PDBP investigators can be found at <https://pdpb.ninds.nih.gov/policy>. The PDBP investigators have not participated in reviewing the data analysis or content of the manuscript. PPMI is sponsored by MJFF and supported by a consortium of scientific partners: (list of the full names of all of the PPMI funding partners can be found at <https://www.ppmi-info.org/about/ppmi/who-we-are-study-sponsors>). The PPMI investigators have not participated in reviewing the data analysis or content of the manuscript. For up-to-date information on the study, visit www.ppmi-info.org. This research was supported in part by the Intramural Research Program of the NIH, National Institute on Aging (NIA), NIH, Department of Health and Human Services; project number Z01 AG000535 and ZIA AG000949, as well as the NINDS (program, ZIANS003154).

We thank all participants of the Luxembourg Parkinson's Study for their important support of our research. Data used for the Luxembourg cohort in the preparation of this manuscript were obtained from the National Centre of Excellence in Research on Parkinson's Disease (NCER-PD). We acknowledge the joint effort of the NCER-PD consortium members from the partner institutions Luxembourg Centre for Systems Biomedicine, Luxembourg Institute of Health, Centre Hospitalier de Luxembourg, and Laboratoire National de Santé generally contributing to the Luxembourg Parkinson's Study as listed below: Geeta Acharya², Gloria Aguayo², Myriam Alexandre², Muhammad Ali², Wim Ammerlann², Giuseppe Arena¹, Rudi Balling¹, Michele Bassis¹, Katy Beaumont², Regina Becker², Camille Bellora², Guy Berchem³, Daniela Berg¹¹, Alexandre Bisdroff², Ibrahim Boussaad¹, Kathrin Brockmann¹¹, Jessica Calmes², Lorieza Castillo², Gessica Contesotto², Nico Diederich², Rene Dondelinger², Daniela Esteves², Guy Fagherazzi², Jean-Yves Ferrand², Manon Gantenbein², Thomas Gasser¹¹, Piotr Gawron¹, Soumyabrata Ghosh¹, Marijus Giraitis^{2,3}, Enrico Glaab¹, Elisa Gómez De Lope¹, Jérôme Graas², Mariella Graziano¹⁷, Valentin Groues¹, Anne Grünewald¹, Wei Gu¹, Gaël Hammo², Anne-Marie Hanff², Linda Hansen^{1,3}, Michael Heneka¹, Estelle Henry², Sylvia Herbrink⁶, Sascha Herzinger¹, Michael Heymann², Michele Hu⁸, Alexander Hundt², Nadine Jacoby¹⁸, Jacek Jaroslaw Lebiada¹, Yohan Jaroz¹, Sonja Jónsdóttir², Quentin Klopstein¹, Jochen Klucken^{1,2,3}, Rejko Krüger^{1,2,3}, Pauline Lambert², Zied Landoulsi¹, Roseline Lentz⁷, Inga Liepelt¹¹, Robert Liszka¹⁴, Laura Longhino³, Victoria Lorentz², Paula Cristina Lupu², Tainá M. Marques¹, Clare Mackay¹⁰, Walter Maetzler¹⁵, Katrin Marcus¹³, Guilherme Marques², Patricia Martins Conde¹, Patrick May¹, Deborah McIntyre², Chouaib Mediouni², Francoise Meisch¹, Myriam Menster², Maura Minelli², Michel Mittelbronn^{1,4}, Brit Mollenhauer¹², Friedrich Mühlshlegel⁴, Romain Nati³, Ulf Nehrbass², Sarah Nickels¹, Beatrice Nicolai³, Jean-Paul Nicolay¹⁹, Fozia Noor², Marek Ostaszewski¹, Clarissa P. C. Gomes¹, Sinthuja Pachchek¹, Claire Pauly^{1,3}, Laure Pauly¹, Lukas Pavelka^{1,3}, Magali Perquin², Rosalina Ramos Lima², Armin Rauschenberger¹, Rajesh Rawal¹, Dheeraj Reddy Bobbili¹, Kirsten Roomp¹, Eduardo Rosales², Isabel Rosety¹, Estelle Sandt², Stefano Sapientza¹, Venkata Satagopam¹, Margaux Schmitt², Sabine Schmitz¹, Reinhard Schneider¹, Jens Schwaborn¹, Amir Sharif², Ekaterina Soboleva¹, Kate Sokolowska², Hermann Thien², Elodie Thiry³, Rebecca Ting Jiin Loo¹, Christophe Trefois¹, Johanna Trouet², Olena Tsurkalenko², Michel Vaillant², Mesele Valenti¹, Gilles Van Cutsem^{1,3}, Carlos Vega¹, Liliana Vilas Boas³, Maharsi Vyas¹, Richard Wade-Martins⁹, Paul Wilmes¹, Evi Wollscheid-Lengeling¹, Gelani Zelimkhanov³.

¹Luxembourg Centre for Systems Biomedicine, University of Luxembourg, Esch-sur-Alzette, Luxembourg.

²Luxembourg Institute of Health, Strassen, Luxembourg.

³Centre Hospitalier de Luxembourg, Strassen, Luxembourg.

⁴Laboratoire National de Santé, Dudelange, Luxembourg.

⁵Centre Hospitalier Emile Mayrisch, Esch-sur-Alzette, Luxembourg.

⁶Centre Hospitalier du Nord, Ettelbrück, Luxembourg.

⁷Parkinson Luxembourg Association, Leudelange, Luxembourg.

⁸Oxford Parkinson's Disease Centre, Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, UK.

⁹Oxford Parkinson's Disease Centre, Department of Physiology, Anatomy and Genetics, University of Oxford, Oxford, UK.

¹⁰Oxford Centre for Human Brain Activity, Wellcome Centre for Integrative Neuroimaging, Department of Psychiatry, University of Oxford, Oxford, UK.

¹¹Center of Neurology and Hertie Institute for Clinical Brain Research, Department of Neurodegenerative Diseases, University Hospital Tübingen, Tübingen, Germany.

¹²Paracelsus-Elena-Klinik, Kassel, Germany.

¹³Ruhr-University of Bochum, Bochum, Germany.

¹⁴Westpfalz-Klinikum GmbH, Kaiserslautern, Germany.

¹⁵Department of Neurology, University Medical Center Schleswig-Holstein, Kiel, Germany.

¹⁶Department of Neurology Philipps, University Marburg, Marburg, Germany.

¹⁷Association of Physiotherapists in Parkinson's Disease Europe, Esch-sur-Alzette, Luxembourg.

¹⁸Private practice, Ettelbruck, Luxembourg.

¹⁹Private practice, Luxembourg, Luxembourg.

PreCEPT and PostCEPT were funded by NINDS 5U01NS050095-05, Department of Defense Neurotoxin Exposure Treatment Parkinson's Research Program (W23RRYX7022N606), MJFF, Parkinson's Disease Foundation, Lundbeck Pharmaceuticals, Cephalon, Lundbeck, John Blume Foundation, Smart Family Foundation, RJG Foundation, Kinetics Foundation, National Parkinson Foundation, Amarin Neuroscience LTD, CHDI Foundation, NIH (NHGRI and NINDS), and Columbia Parkinson's Disease Research Center. Z.G.O. is supported by the Fonds de recherche du Québec—Santé (FRQS) Chercheurs-boursiers award, and is a William Dawson Scholar. Y.L.S. is supported by the HBHL Graduate student fellowship. ProPARK is funded by the Alkemade-Keuls Foundation, Stichting Parkinson Fonds, Parkinson Vereniging, and The Netherlands Organization for Health Research and Development; Udall is supported by the NINDS. This work at the Mayo Clinic Florida was supported by the Haworth Family Professorship in Neurodegenerative Diseases fund, and the Albertson Parkinson's Research Foundation.

Ethics Statement

IRB study number A11-M60-21A (21-11-023) was reviewed and approved by the research ethics offices (REOs). Informed written patient consent was provided in each center before the inclusion of each in the study. We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this work is consistent with those guidelines.

Data Availability Statement

The LID GWAS summary statistics are publicly available on GWAS catalog (<https://www.ebi.ac.uk/gwas/>). All codes used for the analyses are available at <https://github.com/gan-orlab>.

References

1. Armstrong MJ, Okun MS. Diagnosis and Treatment of Parkinson Disease: A Review. *JAMA* 2020;323(6):548–560.
2. Thanvi B, Lo N, Robinson T. Levodopa-induced dyskinesia in Parkinson's disease: clinical features, pathogenesis, prevention and treatment. *Postgrad Med J* 2007;83(980):384–388.
3. Turcano P, Mielke MM, Bower JH, et al. Levodopa-induced dyskinesia in Parkinson disease: A population-based cohort study. *Neurology* 2018;91(24):e2238–e2243.
4. Sweet RD, McDowell FH. Five years' treatment of Parkinson's disease with levodopa. Therapeutic results and survival of 100 patients. *Ann Intern Med* 1975;83(4):456–463.

5. Impact of deprenyl and tocopherol treatment on Parkinson's disease in DATATOP patients requiring levodopa. Parkinson Study Group. *Ann Neurol* 1996;39(1):37–45.
6. Bezard E, Brotchie JM, Gross CE. Pathophysiology of levodopa-induced dyskinesia: potential for new therapies. *Nat Rev Neurosci* 2001;2(8):577–588.
7. Lindgren HS, Andersson DR, Lagerkvist S, Nissbrandt H, Cenci MA. L-DOPA-induced dopamine efflux in the striatum and the substantia nigra in a rat model of Parkinson's disease: temporal and quantitative relationship to the expression of dyskinesia. *J Neurochem* 2010;112(6):1465–1476.
8. Kwon DK, Kwatra M, Wang J, Ko HS. Levodopa-Induced Dyskinesia in Parkinson's Disease: Pathogenesis and Emerging Treatment Strategies. *Cells* 2022;11(23):3736.
9. Yang K, Zhao X, Wang C, Zeng C, Luo Y, Sun T. Circuit Mechanisms of L-DOPA-Induced Dyskinesia (LID). *Front Neurosci* 2021;15:614412.
10. Calabresi P, Ghiglieri V, Mazzocchi P, Corbelli I, Picconi B. Levodopa-induced plasticity: a double-edged sword in Parkinson's disease? *Philos Trans R Soc Lond B Biol Sci* 2015;370(1672):20140184.
11. Fahn S. Parkinson disease, the effect of levodopa, and the ELLDOPA trial, Earlier vs Later L-DOPA. *Arch Neurol* 1999;56(5):529–535.
12. Zappia M, Annesi G, Nicoletti G, et al. Sex differences in clinical and genetic determinants of levodopa peak-dose dyskinesias in Parkinson disease: an exploratory study. *Arch Neurol* 2005;62(4):601–605.
13. Stocchi F, Olanow CW. Continuous dopaminergic stimulation in early and advanced Parkinson's disease. *Neurology* 2004;62(1 Suppl 1):S56–S63.
14. Gilgun-Sherki Y, Djaldetti R, Melamed E, Offen D. Polymorphism in candidate genes: implications for the risk and treatment of idiopathic Parkinson's disease. *Pharmacogenomics J* 2004;4(5):291–306.
15. Tran TN, Vo TNN, Frei K, Truong DD. Levodopa-induced dyskinesia: clinical features, incidence, and risk factors. *J Neural Transm* 2018;125(8):1109–1117.
16. Rieck M, Schumacher-Schuh AF, Altmann V, et al. DRD2 haplotype is associated with dyskinesia induced by levodopa therapy in Parkinson's disease patients. *Pharmacogenomics* 2012;13(15):1701–1710.
17. Lee JY, Cho J, Lee EK, Park SS, Jeon BS. Differential genetic susceptibility in diphasic and peak-dose dyskinesias in Parkinson's disease. *Mov Disord* 2011;26(1):73–79.
18. Comi C, Ferrari M, Marino F, et al. Polymorphisms of Dopamine Receptor Genes and Risk of L-Dopa-Induced Dyskinesia in Parkinson's Disease. *Int J Mol Sci* 2017;18(2):242.
19. Kaiser R, Hofer A, Grapengiesser A, et al. L -dopa-induced adverse effects in PD and dopamine transporter gene polymorphism. *Neurology* 2003;60(11):1750–1755.
20. Kaplan N, Vituri A, Korczyn AD, et al. Sequence variants in SLC6A3, DRD2, and BDNF genes and time to levodopa-induced dyskinesias in Parkinson's disease. *J Mol Neurosci* 2014;53(2):183–188.
21. Tan YY, Jenner P, Chen SD. Monoamine Oxidase-B Inhibitors for the Treatment of Parkinson's Disease: Past, Present, and Future. *J Parkinsons Dis* 2022;12(2):477–493.
22. Rivest J, Barclay CL, Suchowersky O. COMT inhibitors in Parkinson's disease. *Can J Neurol Sci* 1999;26(Suppl 2):S34–S38.
23. de Lau LM, Verbaan D, Marinus J, Heutink P, van Hilten JJ. Catechol-O-methyltransferase Val158Met and the risk of dyskinesias in Parkinson's disease. *Mov Disord* 2012;27(1):132–135.
24. Cheshire P, Bertram K, Ling H, et al. Influence of single nucleotide polymorphisms in COMT, MAO-A and BDNF genes on dyskinesias and levodopa use in Parkinson's disease. *Neurodegener Dis* 2014;13(1):24–28.
25. Hao H, Shao M, An J, et al. Association of Catechol-O-Methyltransferase and monoamine oxidase B gene polymorphisms with motor complications in parkinson's disease in a Chinese population. *Parkinsonism Relat Disord* 2014;20(10):1041–1045.
26. Bialecka M, Drożdż M, Kłodowska-Duda G, et al. The effect of monoamine oxidase B (MAOB) and catechol-O-methyltransferase (COMT) polymorphisms on levodopa therapy in patients with sporadic Parkinson's disease. *Acta Neurol Scand* 2004;110(4):260–266.
27. Gan-Or Z, Amshalom I, Kilarski LL, et al. Differential effects of severe vs mild GBA mutations on Parkinson disease. *Neurology* 2015;84(9):880–887.
28. Ross OA, Soto-Ortolaza AI, Heckman MG, et al. Association of LRRK2 exonic variants with susceptibility to Parkinson's disease: a case-control study. *Lancet Neurol* 2011;10(10):898–908.
29. Olszewska DA, McCarthy A, Soto-Beasley AI, et al. Association Between Glucocerebrosidase Mutations and Parkinson's Disease in Ireland. *Front Neurol* 2020;11:527.
30. Jesús S, Huertas I, Bernal-Bernal I, et al. GBA Variants Influence Motor and Non-Motor Features of Parkinson's Disease. *PLoS One* 2016;11(12):e0167749.
31. Lesage S, Anheim M, Condroyer C, et al. Large-scale screening of the Gaucher's disease-related glucocerebrosidase gene in Europeans with Parkinson's disease. *Hum Mol Genet* 2011;20(1):202–210.
32. Bouhouche A, Tibar H, Ben El Haj R, et al. LRRK2 G2019S Mutation: Prevalence and Clinical Features in Moroccans with Parkinson's Disease. *Parkinson's Dis* 2017;2017:2412486.
33. Gao C, Pang H, Luo XG, Ren Y, Shang H, He ZY. LRRK2 G2385R variant carriers of female Parkinson's disease are more susceptible to motor fluctuation. *J Neurol* 2013;260(11):2884–2889.
34. Shu L, Zhang Y, Pan H, et al. Clinical Heterogeneity Among LRRK2 Variants in Parkinson's Disease: A Meta-Analysis. *Front Aging Neurosci* 2018;10:283.
35. Leaver K, Viser A, Kopell BH, et al. Clinical profiles and outcomes of deep brain stimulation in G2019S LRRK2 Parkinson disease. *J Neurosurg* 2021;137(1):1–8.
36. Stanic J, Mellone M, Cîrnuș MD, et al. LRRK2 phosphorylation level correlates with abnormal motor behaviour in an experimental model of levodopa-induced dyskinesias. *Mol Brain* 2016;9(1):53.
37. Wile DJ, Agarwal PA, Schulzer M, et al. Serotonin and dopamine transporter PET changes in the premotor phase of LRRK2 parkinsonism: cross-sectional studies. *Lancet Neurol* 2017;16(5):351–359.
38. Menozzi E, Schapira AHV. Exploring the Genotype-Phenotype Correlation in GBA-Parkinson Disease: Clinical Aspects, Biomarkers, and Potential Modifiers. *Front Neurol* 2021;12:694764.
39. Alcalay RN, Mejia-Santana H, Mirelman A, et al. Neuropsychological performance in LRRK2 G2019S carriers with Parkinson's disease. *Parkinsonism Relat Disord* 2015;21(2):106–110.
40. Foltynie T, Cheeran B, Williams-Gray CH, et al. BDNF val66met influences time to onset of levodopa induced dyskinesia in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2009;80(2):141–144.
41. Kusters CDJ, Paul KC, Guella I, et al. Dopamine receptors and BDNF-haplotypes predict dyskinesia in Parkinson's disease. *Parkinsonism Relat Disord* 2018;47:39–44.
42. Ivanova SA, Loonen AJ, Pechlivanoglou P, et al. NMDA receptor genotypes associated with the vulnerability to develop dyskinesia. *Transl Psychiatry* 2012;2(1):e67.
43. Rieck M, Schumacher-Schuh AF, Callegari-Jacques SM, et al. Is there a role for ADORA2A polymorphisms in levodopa-induced dyskinesia in Parkinson's disease patients? *Pharmacogenomics* 2015;16(6):573–582.
44. Oeda T, Umemura A, Mori Y, et al. Impact of glucocerebrosidase mutations on motor and nonmotor complications in Parkinson's disease. *Neurobiol Aging* 2015;36(12):3306–3313.
45. Tirozzi A, Modugno N, Palomba NP, et al. Analysis of Genetic and Non-genetic Predictors of Levodopa Induced Dyskinesia in Parkinson's Disease. *Front Pharmacol* 2021;12:640603.
46. Healy DG, Falchi M, O'Sullivan SS, et al. Phenotype, genotype, and worldwide genetic penetrance of LRRK2-associated Parkinson's disease: a case-control study. *Lancet Neurol* 2008;7(7):583–590.

47. Yahalom G, Kaplan N, Vituri A, et al. Dyskinesias in patients with Parkinson's disease: effect of the leucine-rich repeat kinase 2 (LRRK2) G2019S mutation. *Parkinsonism Relat Disord* 2012; 18(9):1039–1041.
48. Paus S, Gadow F, Knapp M, Klein C, Klockgether T, Wüllner U. Motor complications in patients from the German Competence Network on Parkinson's disease and the DRD3 Ser9Gly polymorphism. *Mov Disord* 2009;24(7):1080–1084.
49. Greenbaum L, Goldwurm S, Zozulinsky P, et al. Do tardive dyskinesia and L-dopa induced dyskinesia share common genetic risk factors? An exploratory study. *J Mol Neurosci* 2013;51(2): 380–388.
50. Hughes AJ, Daniel SE, Kilford L, Lees AJ. Accuracy of clinical diagnosis of idiopathic Parkinson's disease: a clinico-pathological study of 100 cases. *J Neurol Neurosurg Psychiatry* 1992;55(3):181–184.
51. Nalls MA, Bras J, Hernandez DG, et al. NeuroX, a fast and efficient genotyping platform for investigation of neurodegenerative diseases. *Neurobiol Aging* 2015;36(3):1605–1612.
52. Blauwendraat C, Faghri F, Pihlstrom L, et al. NeuroChip, an updated version of the NeuroX genotyping platform to rapidly screen for variants associated with neurological diseases. *Neurobiol Aging* 2017;57:247.e9–247.13.
53. Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007;81(3):559–575.
54. Cavallieri F, Cury RG, Guimarães T, et al. Recent Advances in the Treatment of Genetic Forms of Parkinson's Disease: Hype or Hope? *Cells* 2023;12(5):764.
55. Iwaki H, Blauwendraat C, Leonard HL, et al. Genetic risk of Parkinson disease and progression: An analysis of 13 longitudinal cohorts. *Neurol Genet* 2019;5(4):e348.
56. Julien C, Hache G, Dulac M, et al. The clinical meaning of levodopa equivalent daily dose in Parkinson's disease. *Fundam Clin Pharmacol* 2021;35(3):620–630.
57. Tomlinson CL, Stowe R, Patel S, Rick C, Gray R, Clarke CE. Systematic review of levodopa dose equivalency reporting in Parkinson's disease. *Mov Disord* 2010;25(15):2649–2653.
58. Syed H, Jorgensen AL, Morris AP. SurvivalGWAS_SV: software for the analysis of genome-wide association studies of imputed genotypes with "time-to-event" outcomes. *BMC Bioinf* 2017;18(1):265.
59. Nalls MA, Blauwendraat C, Vallerga CL, et al. Identification of novel risk loci, causal insights, and heritable risk for Parkinson's disease: a meta-analysis of genome-wide association studies. *Lancet Neurol* 2019;18(12):1091–1102.
60. Choi SW, O'Reilly PF. PRSice-2: Polygenic Risk Score software for biobank-scale data. *Gigascience* 2019;8(7):giz082.
61. Bandres-Ciga S, Saez-Atienzar S, Kim JJ, et al. Large-scale pathway specific polygenic risk and transcriptomic community network analysis identifies novel functional pathways in Parkinson disease. *Acta Neuropathol* 2020;140(3):341–358.
62. Martinez-Carrasco A, Real R, Lawton M, et al. Genetic meta-analysis of levodopa induced dyskinesia in Parkinson's disease. *npj Parkinson's Disease* 2023;9(1):128.
63. Espay AJ, Morgante F, Merola A, et al. Levodopa-induced dyskinesia in Parkinson disease: current and evolving concepts. *Ann Neurol* 2018;84(6):797–811.
64. Jennings D, Huntwork-Rodriguez S, Henry AG, et al. Preclinical and clinical evaluation of the LRRK2 inhibitor DNL201 for Parkinson's disease. *Sci Transl Med* 2022;14(648):eabj2658.
65. den Heijer JM, Kruithof AC, Moerland M, et al. A phase 1B trial in GBA1-associated Parkinson's Disease of BIA-28-6156, a glucocerebrosidase activator. *Mov Disord* 2023;38(7):1197–1208.
66. Silveira CRA, MacKinley J, Coleman K, et al. Ambroxol as a novel disease-modifying treatment for Parkinson's disease dementia: protocol for a single-centre, randomized, double-blind, placebo-controlled trial. *BMC Neurol* 2019;19(1):20.
67. Mullin S, Smith L, Lee K, et al. Ambroxol for the treatment of patients with Parkinson Disease with and without glucocerebrosidase gene mutations: a nonrandomized, noncontrolled trial. *JAMA Neurol* 2020;77(4):427–434.
68. Katzenschlager R, Head J, Schrag A, Ben-Shlomo Y, Evans A, Lees AJ. Fourteen-year final report of the randomized PDRG-UK trial comparing three initial treatments in PD. *Neurology* 2008;71(7): 474–480.

Supporting Data

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.