

Reduced Age-Dependent Penetrance of a Large *FGF14* GAA Repeat Expansion in a 74-Year-Old Woman from a German Family with SCA27B

Spinocerebellar ataxia 27B (SCA27B) is a recently described autosomal dominant cerebellar ataxia caused by an intronic GAA repeat expansion in the *FGF14* gene.¹ Since SCA27B is one of the most common genetic ataxias worldwide,^{2,3} a better understanding of disease penetrance is essential for improving genetic counseling and management of at-risk individuals, as well as for guiding eligibility assessment for future clinical trials.

Initial reports have suggested a pathogenic threshold of (GAA)_{≥250} repeat units, albeit with reduced penetrance for (GAA)_{250–300} alleles, since they were observed in ~1.5%–2% of controls.^{1,2} A subsequent study identified expansions ranging from 308 to 380 GAA repeat units in 5 of 475 controls, raising the possibility that (GAA)_{>300} expansions may also show incomplete

penetrance.⁴ Notwithstanding, direct evidence for the reduced penetrance of *FGF14* GAA repeat expansions remains limited. Here, we provide an in-depth, multimodal assessment of an asymptomatic 74-year-old woman carrying a (GAA)₄₈₃ expansion from a family with SCA27B, thus providing further direct evidence for the reduced age-dependent penetrance even of large *FGF14* GAA repeat expansions.

A 74-year-old woman of German descent presented to the Ataxia Clinic in Tübingen for evaluation of her genetic risk following the diagnosis of SCA27B in two of her relatives (Fig. 1A). She reported no symptoms suggestive of SCA27B,² such as episodic or progressive imbalance, visual disturbances, vertigo and/or dizziness, and alcohol sensitivity. Neurological examination by two independent SCA27B experts (M.S., A.T.) was unremarkable (including absence of downbeat nystagmus), except for questionable mild dysidiadochokinesia of the right upper extremity and mild reduction of vibration sense at the ankles bilaterally. The total score on the Scale for the Assessment and Rating of Ataxia (SARA) was 0.5/40 points (threshold for ataxia: ≥3 points⁵). Furthermore, the score on the Friedreich Ataxia Rating Scale (FARS) Part E, a gait and balance assessment tool with even higher sensitivity,⁶ was 0. We next performed quantitative gait analysis using body-worn sensors (see Supporting Information), which is able to detect even subtle gait and balance changes in the very early stages of cerebellar disease.^{7–9} Her gait performance, as assessed by the ataxic-sensitive measures “lateral step deviation” and “stride length variability”⁷ lay outside the 90% confidence intervals for aged-matched ataxic patients, but within the 90% confidence interval for age-matched healthy controls (Fig. 1B,C). Following a standardized protocol employing long-range polymerase chain reaction (PCR) and bidirectional repeat-primed PCRs,¹⁰ genetic testing revealed that her *FGF14* genotype was 483 and 9 GAA repeat units, without evidence of interruptions within the expanded allele. This result was confirmed with two independent blood samples. In comparison, the subject’s brother and cousin, who respectively carried a (GAA)₄₈₈ and (GAA)₄₉₆ expansion, developed slowly progressive cerebellar ataxia at the age of 55 and 63 years, respectively (Table S1).

© 2024 The Author(s). *Movement Disorders* published by Wiley Periodicals LLC on behalf of International Parkinson and Movement Disorder Society.

This is an open access article under the terms of the [Creative Commons Attribution](#) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Key Words: SCA27B, GAA-*FGF14* ataxia, repeat expansion disorder, penetrance

***Correspondence to:** Dr. Matthias Synofzik, Division Translational Genomics of Neurodegenerative Diseases, Hertie Institute for Clinical Brain Research and Center of Neurology, University of Tübingen, Hoppe-Seyler-Str. 3, 72076 Tübingen, Germany; E-mail: matthias.synofzik@uni-tuebingen.de

Relevant conflicts of interest/financial disclosures: D.P., J.S., A.T., B.B., and W.I. report no disclosures. M.S. has received consultancy honoraria from Janssen, Ionis, Orphazyme, Servier, Solaxa, Reata, Biohaven, Zevra, Lilly, GenOrph, and AviadoBio, all unrelated to the present work.

Funding agencies: This work was supported by the Clinician Scientist program “PRECISE.net” funded by the Else Kröner-Fresenius-Stiftung (to A.T. and M.S.) and the grant 779257 “Solve-RD” from the European’s Union Horizon 2020 research and innovation program (to M.S.). This work was also supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) No. 441409627, as part of the PROSPAX consortium under the frame of EJP RD, the European Joint Programme on Rare Diseases, under the EJP RD COFUND-EJP No. 825575 (to M.S. and B.B.), the Fondation Groupe Monaco (to B.B.), and the Canadian Institutes of Health Research (Grant No. 189963 to B.B.). The funders had no role in the conduct of this study.

Received: 30 March 2024; **Revised:** 29 May 2024; **Accepted:** 17 June 2024

Published online 1 July 2024 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.29915




FIG. 1. Pedigree of the German family with SCA27B and quantitative gait analysis. (A) Pedigree of the German family with spinocerebellar ataxia 27B (SCA27B). The allele sizes, expressed as numbers of GAA repeat units, are provided for the family members who underwent genotyping of the *FGF14* repeat locus. Squares represent male family members and circles female family members. Solid black shapes indicate affected persons. Slashed symbols indicate deceased persons. The pedigree has been abbreviated and family members randomly rearranged to preserve privacy. The subject of this study is indicated by a black arrow. Comparison of (B) lateral step deviation and (C) stride length variability by quantitative gait analysis using APDM wearable sensors between the index subject (red dot), healthy controls (HC, green), and patients with degenerative cerebellar ataxia (ATX, blue), according to their age at time of assessment. The green line indicates the upper bound of the 90% confidence interval around the average gait performance of healthy controls. The blue line indicates the lower bound of the 90% confidence interval around the average gait performance of ataxic patients. The color of the data points for the ataxic individuals is a function of the severity of ataxia as measured by the Scale for the Assessment and Rating of Ataxia (SARA) score, with dark blue indicating more severe ataxia (a hue scale is shown on the right y-axis). [Color figure can be viewed at wileyonlinelibrary.com]

Through in-depth, multimodal assessment that included highly sensitive standardized clinical scoring (FARS-E) and quantitative digital-motor gait analysis, this study provides first direct evidence for reduced age-dependent penetrance of a large *FGF14* (GAA)₄₈₃ repeat expansion in an otherwise still asymptomatic 74-year-old woman from a German family with SCA27B. Quantitative analysis of gait parameters shown to be highly sensitive to mild gait and balance deficits, namely lateral step deviation and stride length variability,⁷ showed that the subject had a performance within the range of age-matched healthy controls, but outside the range of ataxic individuals. Our findings extend previous evidence of reduced penetrance of *FGF14* GAA repeat expansions, as supported by their observation in non-ataxic controls and inferred from the frequent sporadic presentation of SCA27B (15%–50%),² where an asymptomatic parent must have been an obligate expansion carrier. Although it is not clear how thoroughly the relatives of sporadic cases reported in previous series were assessed, they were likely not assessed using highly sensitive clinical scales like the FARS-E and quantitative gait assessment.

While most patients with SCA27B develop ataxia in their 50s and 60s,² some occasionally manifest ataxia in their 80s,¹ thus not ruling out that the subject reported here may develop SCA27B later in life. While longitudinal follow-up is warranted to investigate this, our results support the reduced age-dependent penetrance of large *FGF14* repeat expansions past the age of 70 years. They also further corroborate the observed intrafamilial variability in the age at onset associated even with *FGF14* GAA repeat expansions of same size.² The reduced penetrance of *FGF14* expansions may be the result of modifying genetic factors and/or somatic mosaicism in the central nervous system, similar to what has previously been shown in Friedreich ataxia.¹¹

In conclusion, our findings show that reduced age-dependent penetrance may also be observed with large *FGF14* repeat expansions past the age of 70 years, which is of importance for genetic counseling and future trial planning. In addition, they highlight the need to comprehensively study factors that modulate age of onset and age-dependent penetrance of SCA27B, given its high frequency. ■

Acknowledgments: We thank the family for participating in this study. We thank Ms. Marie-Josée Dicaire (The Montreal Neurological Hospital-Institute, Montreal, Canada) for her assistance with the genotyping. D.P. holds a Fellowship Award from the Canadian Institutes of Health Research (CIHR).

Ethical Compliance Statement

The study was approved by the Institutional Review Board of the University of Tübingen (AZ 598/2011BO1).

Written informed consent was obtained from all study participants before enrollment.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

David Pellerin, MD,^{1,2}  Jens Seemann, MSc,^{3,4}

Andreas Traschütz, MD, PhD,^{5,6} 

Bernard Brais, MDCM, PhD,^{1,7,8} Winfried Ilg, PhD,^{3,4}  and

Matthis Synofzik, MD,^{5,6*} 

¹Department of Neurology and Neurosurgery, Montreal Neurological Hospital and Institute, McGill University, Montreal, Quebec, Canada, ²Department of Neuromuscular Diseases, UCL Queen Square Institute of Neurology and The National Hospital for Neurology and Neurosurgery, University College London, London, UK, ³Section Computational Sensomotrics, Hertie Institute for Clinical Brain Research, Tübingen, Germany, ⁴Centre for Integrative Neuroscience (CIN), Tübingen, Germany, ⁵Division Translational Genomics of Neurodegenerative Diseases, Hertie Institute for Clinical Brain Research and Center of Neurology, University of Tübingen, Tübingen, Germany, ⁶German Center for Neurodegenerative Diseases (DZNE), Tübingen, Germany, ⁷Department of Human Genetics, McGill University, Montreal, Quebec, Canada, and ⁸Centre de Réadaptation Lucie-Bruneau, Montreal, Quebec, Canada

References

- Pellerin D, Danzi MC, Wilke C, Renaud M, Fazal S, Dicaire MJ, et al. Deep intronic *FGF14* GAA repeat expansion in late-onset cerebellar ataxia. *N Engl J Med* 2023;388(2):128–141.
- Pellerin D, Danzi MC, Renaud M, Houlden H, Synofzik M, Zuchner S, et al. Spinocerebellar ataxia 27B: a novel, frequent and potentially treatable ataxia. *Clin Transl Med* 2024;14(1):e1504.
- Hengel H, Pellerin D, Wilke C, Fleszar Z, Brais B, Haack T, et al. As frequent as polyglutamine spinocerebellar ataxias: SCA27B in a large German autosomal dominant ataxia cohort. *Mov Disord* 2023;38(8):1557–1558.
- Méreaux JL, Davoine CS, Pellerin D, Coarelli G, Coutelier M, Ewencyk C, et al. Clinical and genetic keys to cerebellar ataxia due to *FGF14* GAA expansions. *EBioMedicine* 2024;99:104931.
- Maas RP, van Gaalen J, Klockgether T, van de Warrenburg BP. The preclinical stage of spinocerebellar ataxias. *Neurology* 2015;85(1):96–103.
- Rummey C, Harding IH, Delatycki MB, Tai G, Rezende T, Corben LA. Harmonizing results of ataxia rating scales: mFARS, SARA, and ICARS. *Ann Clin Transl Neurol* 2022;9(12):2041–2046.
- Seemann J, Daghsen L, Cazier M, Lamy JC, Welter ML, Giese MA, et al. Digital gait measures capture 1-year progression in early-stage spinocerebellar ataxia type 2. *Mov Disord* 2024;39(5):788–797.
- Seemann J, Traschütz A, Ilg W, Synofzik M. 4-Aminopyridine improves real-life gait performance in SCA27B on a single-subject level: a prospective n-of-1 treatment experience. *J Neurol* 2023;270(11):5629–5634.
- Ilg W, Seemann J, Giese M, Traschütz A, Schöls L, Timmann D, et al. Real-life gait assessment in degenerative cerebellar ataxia: toward ecologically valid biomarkers. *Neurology* 2020;95(9):e1199–e1210.
- Bonnet C, Pellerin D, Roth V, Clément G, Wandzel M, Lambert L, et al. Optimized testing strategy for the diagnosis of GAA-*FGF14* ataxia/spinocerebellar ataxia 27B. *Sci Rep* 2023;13(1):9737.
- De Biase I, Rasmussen A, Monticelli A, Al-Mahdawi S, Pook M, Coccozza S, et al. Somatic instability of the expanded GAA triplet-

repeat sequence in Friedreich ataxia progresses throughout life. *Genomics* 2007;90(1):1–5.

Supporting Data

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.

Cerebellar Ataxia Secondary to Leukodystrophy with a Frameshift *CST3* Variant



Leukodystrophies are a group of disorders involving the white matter of the central nervous system.¹ Recently, a novel adult onset leukodystrophy without amyloid angiopathy caused by specific dominant *CST3* truncating variants was reported.² Here, we describe a novel frameshift *CST3* variant causing leukodystrophy in a Chinese patient manifesting as predominant cerebellar ataxia.

The patient (Fig. 1A) was a 58-year-old female with a history of progressive gait imbalance and slurred speech for 5 years. The gait imbalance worsened progressively, and she gradually developed tremor of the head and both hands. Her mother had a similar history of gait imbalance before death. Neurological examinations showed dysarthria, dysdiadochokinesia, postural tremor of upper limbs, dysmetria on both finger-to-nose and heel-to-shin. Wide-based gait, tandem gait difficulty, and positive Romberg's sign were also observed. Deep tendon reflexes were normal, and Babinski's signs were negative (Video S1). Sensory examination was normal.

© 2024 International Parkinson and Movement Disorder Society.

Key Words: cerebellar ataxia, *CST3*, leukodystrophy

***Correspondence to:** Jiting Zhu, MD, and Zhi-Qiang Wang, MD, Department of Neurology and Institute of Neurology of First Affiliated Hospital, Institute of Neuroscience, and Fujian Key Laboratory of Molecular Neurology, Fujian Medical University, Fuzhou 350005, China. E-mail: jitingzhu@fjmu.edu.cn and wzhq1992@fjmu.edu.cn

Yi-Heng Zeng and Dan-Dan Zuo contributed equally to this work.

Relevant conflicts of interest/financial disclosures: None.

Funding agency: This work was supported by grants from the Natural Science Foundation of Fujian Province (2022J01703) to J.T.Z.

Received: 2 May 2024; **Revised:** 27 May 2024; **Accepted:** 17 June 2024

Published online 4 July 2024 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.29913