

ABSTRACT CITATION ID: NOAE064.236

EPID-04. CAUSES AND LONG-TERM COURSE OF OTOTOXICITY IN PEDIATRIC MEDULLOBLASTOMA AND EPENDYMOMA PATIENTS – CUMULATIVE PLATINUM AND COCHLEAR IRRADIATION DOSE MATTER

Elif Öz¹, Annika Hardt², Rolf-Dieter Kortmann³, Rudolf Schwarz², Beate Timmermann^{4,5}, Brigitte Bison⁶, Monika Warmuth-Metz⁷, Katja von Hoff^{8,9}, André O. von Bueren¹⁰, Nicolas U. Gerber¹¹, Gudrun Fleischhack¹², Torsten Pietsch¹³, Stefan M. Pfister^{14,15}, Dominik Sturm^{15,16}, Martin Mynarek^{1,17}, Stefan Rutkowski¹, Denise Obrecht-Sturm¹; ¹Pediatric Hematology and Oncology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany, ²Department for Radiotherapy, University Medical Center Hamburg-Eppendorf, Hamburg, Germany, ³Department of Radiation Oncology, University of Leipzig, Leipzig, Germany, ⁴West German Proton Therapy Center Essen (WPE), University Hospital Essen, Essen, Germany, ⁵Clinic for Particle Therapy, University Hospital Essen, Essen, Germany, ⁶Department of Neuroradiology, University Hospital Augsburg, Augsburg, Germany, ⁷Institute of Diagnostic and Interventional Neuroradiology, University Hospital Würzburg, Würzburg, Germany, ⁸Department of Pediatric Oncology and Hematology, Charité - Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany, ⁹Department of Pediatrics and Adolescent Medicine, Aarhus University Hospital, Aarhus, Denmark, ¹⁰Department of Pediatrics, Obstetrics and Gynecology, Division of Pediatric Hematology and Oncology, University Hospital of Geneva, Geneva, Switzerland, ¹¹Department of Pediatric Oncology, University Children's Hospital Zürich, Zürich, Switzerland, ¹²Pediatrics III, Pediatric Oncology and Hematology, University Hospital Essen, Essen, Germany, ¹³Institute of Neuropathology, Brain Tumor Reference Center of the German Society for Neuropathology and Neuroanatomy (DGNN), University of Bonn, DZNE German Center for Neurodegenerative Diseases, Bonn, Germany, ¹⁴Hopp Children's Cancer Center Heidelberg (KiTZ) and Division of Pediatric Neurooncology, German Cancer Research Center (DKFZ) and German Consortium for Translational Cancer Research (DKTK), Heidelberg, Germany, ¹⁵Department of Pediatric Hematology and Oncology, Heidelberg University Hospital, Heidelberg, Germany, ¹⁶Division of Pediatric Glioma Research, German Cancer Research Center (DKFZ) and German Consortium for Translational Cancer Research (DKTK), Heidelberg, Germany, ¹⁷Mildred Scheel Cancer Career Center HaTriCS4, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

BACKGROUND: Ototoxicity is a well-known side effect of treatment with platinum-containing chemotherapy and radiotherapy (RT). Nevertheless, precise dose-effect relationships are still lacking. **METHODS:** Patients from the HIT2000 trial receiving first-line treatment for medulloblastoma (MB) or ependymoma (EPN) who were alive at last follow-up and had at least one audiogram prior to and after administration of chemotherapy with cisplatin and radiotherapy were included into this study. Treatment details (incl. cochlear irradiation dose [CID]) and audiological outcome were analyzed. The ototoxicity was graded using the HIT scoring. **RESULTS:** Courses of 101 patients were analyzed. Median audiology follow-up was 6 years [range: 1.0–18.1 years]. Ototoxicity grade ≥ 1 was documented for n=66 patients (65.4%). For n=27 this resulted in treatment modification. Univariable analyses revealed different proportions of ototoxicity (grade ≥ 1) regarding histology (CMB: 50/65, LCAMB: 3/3, DMB/MBEN: 8/12, EPN: 5/21; $p < 0.001$), tumor location (supratentorial: 3/9, infratentorial: 63/92; $p < 0.001$), metastatic status at diagnosis (M0: 46/77, M+: 20/24; $p = 0.003$), radiation field (CSI: 59/75, local RT: 6/21, no RT: 1/5; $p < 0.001$), CID (continuous variable, $p = 0.001$) and cumulative cisplatin dose (continuous variable, $p < 0.001$). Certainly, disease specific parameters (histology/staging) were related to treatment intensity (cisplatin/irradiation dose). Multivariable analyses confirmed the impact of cumulative cisplatin dose ($p = 0.01$) and CID ($p = 0.04$). ROC-analyses allowed the definition of cut-offs for high risk of ototoxicity grade ≥ 1 (cisplatin: 70mg/m² BSA, CID: 37.5 Gy) and ototoxicity risk estimation, when applying doses above the respective cut-offs (cisplatin dose increase by 1mg/m² BSA results in 4% higher risk for ototoxicity grade ≥ 1 ; CID dose increase by 1 Gy results in 4.6% higher risk for ototoxicity grade ≥ 1). **CONCLUSIONS:** Ototoxicity affects a majority of children receiving adjuvant treatment for MB and EPN. To avoid disabilities and negative effect on quality of life, dosage of cisplatin and radiotherapy should be scrutinized and reduced whenever possible.