



Centrosomal and acentrosomal microtubule nucleation during neuronal development

Stanislav Vinopal¹ and Frank Bradke²

Neurons rely on the microtubule cytoskeleton to create and maintain their sophisticated cellular architectures. Advances in cryogenic electron microscopy, expansion microscopy, live imaging, and gene editing have enabled novel insights into mechanisms of centrosomal and acentrosomal microtubule nucleation, the key process generating new microtubules. This has paved the way for the functional dissection of distinct microtubule networks that regulate various processes during neuronal development, including neuronal delamination, polarization, migration, maturation, and synapse function. We review recent progress in understanding the molecular concepts of microtubule nucleation, how these concepts underlie neurodevelopmental processes, and pinpoint the open questions. Since microtubules play a pivotal role in axon regeneration within the adult central nervous system, understanding the processes of microtubule nucleation could inform strategies to enhance the regenerative capabilities of neurons in the future.

Addresses

¹ Centre for Nanomaterials and Biotechnology, Faculty of Science, Jan Evangelista Purkyně University (UJEP), Usti nad Labem, Czech Republic

² Laboratory of Axonal Growth and Regeneration, German Center for Neurodegenerative Diseases (DZNE), Bonn, Germany

Corresponding authors: Vinopal, Stanislav (stanislav.vinopal@ujep.cz); Bradke, Frank (Frank.Bradke@dzne.de)

Current Opinion in Neurobiology 2025, **92**:103016

This review comes from a themed issue on **Developmental Neuroscience 2025**

Edited by **Lisa Goodrich** and **Song-Hai Shi**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online xxx

<https://doi.org/10.1016/j.conb.2025.103016>

0959-4388/© 2025 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Microtubules are cytoskeletal filaments that drive myriads of cellular processes, governing cell shape, motility, intracellular transport, cell division, and signalling [1–4]. Microtubules are essential especially in neurons, highly polarized cells with sophisticated morphology

[4]. It enables them to form complex networks, giving rise to emergent properties that reach its pinnacle in human consciousness.

Yet, neurons are born as small, simple cells. They undergo complex morphological transformations, while growing and travelling to their destination in the brain [5–7]. Neurons make thousands of connections, typically via elaborate dendritic trees and axons [6], whose architecture relies on different microtubule networks. Some undergo massive expansion and become progressively stable, such as microtubule bundles in extending axons [8]. Others, such as microtubules in navigating growth cones, undergo cyclic remodelling [7,9,10]. As development progresses, microtubule networks gradually diversify. In axons, microtubules form a unipolar arrangement, whereas in neurites, they exhibit mixed polarity, varying by developmental stage and species [11,12]. This unique organization provides a basis for further regulatory mechanisms that coordinate cargo molecule sorting and are important for establishing neuronal polarity [11–14]. Microtubules also form a cage surrounding the nucleus during neuronal migration [7]. They are connected to the centrosome that is cyclically moving inside the cytoplasmic dilation [15], coupling the leading process and the nucleus in the neuronal rear [7]. Other microtubule networks form elaborate signalling structures, including dendritic trees, primary cilia and synapses [16,17].

Such massive expansion and remodeling requires highly regulated generation of new microtubules. They arise from severing of existing microtubules by microtubule severing enzymes, such as katanin, spastin or fidgetin, with microtubule remnants serving as polymerization seeds, or by *de novo* microtubule nucleation [4]. Microtubule nucleation has been implicated in many key events in neuronal lives and will be the main topic of this review.

First, we will discuss advances in the understanding of different modes of microtubule nucleation. Second, we will summarize when and where these modes of microtubule nucleation are employed, and how they are related to the diverse and dynamically changing microtubule networks during neuronal development. It will provide insights into how microtubule networks drive the form and function of neurons.

Modes of microtubule nucleation

Microtubule nucleation by γ -TuRC

Microtubules are stiff, 25 nm thick tubes. They form by unidirectional lateral association of typically 13 protofilaments, linear polymers of α - and β -tubulin heterodimers [1]. α -Tubulin is exposed on the minus end and β -tubulin on the plus end of each protofilament [4]. Minus and plus ends differ structurally and biochemically, which results in different kinetics of microtubule polymerization and depolymerization, with plus ends exhibiting much greater dynamicity [4]. This difference magnifies in cells, where specific proteins bind to the two different microtubule ends, modulating their dynamics [1,4].

In cells, microtubule nucleation is catalyzed primarily by the cone-shaped γ -Tubulin Ring Complex (γ -TuRC) [4], which contains seven main protein modules, each composed of a γ -tubulin complex protein (GCP) heterodimer and two γ -tubulins [18–20] (Figure 1a).

The assembled γ -TuRC is splayed. Thereby, it does not match the 13-protofilament microtubule, explaining its low nucleation activity [20–25]. Recent studies revealed that γ -TuRC can transition to a more closed ('early closed') conformation, which increases its microtubule nucleation efficiency [19–21,26] (Figure 1a).

γ -TuRC associates with many proteins that modulate its activity and subcellular targeting. For a more complete list of these proteins, we refer to recent excellent reviews [2,4,27]. An important class of γ -TuRC-interacting proteins are its stimulators. We will discuss CDK5 regulatory subunit-associated protein 2 (CDK5RAP2) [28,29], Targeting protein for Xklp2 (TPX2) [30], the augmin complex [31], AKNA [32], Cytoskeleton-associated protein 5 (CKAP5), and CLIP-associating protein 2 (CLASP2) [33].

Proteins stimulating γ -TuRC activity

The *bona fide* γ -TuRC activator is CDK5RAP2 (Centrosomin in *Drosophila*). It contains a conserved Centrosomin motif 1 (CM1, also termed γ -tubulin nucleation activator, γ -TuNA) that binds to the γ -TuRC via Mitotic-spindle organizing protein 2 (MZT2) and GCP2, changing γ -TuRC conformation to the 'early closed' state that better matches the 13-protofilament microtubules [28,29] (Figure 1b). Interestingly, only a fraction of γ -TuRCs interact with CDK5RAP2 in cells [15,34] and the interaction is likely transient [28]. CDK5RAP2 autoinhibition prevents uncontrolled γ -TuRC activation [35,36].

TPX2 stimulates γ -TuRC-dependent microtubule nucleation in several ways. It recruits γ -TuRC, soluble tubulin, and augmin to microtubules, enhancing branching microtubule nucleation [2,27] (Figure 1c).

However, TPX2 is dispensable for augmin-dependent branching microtubule nucleation in some systems [31].

Augmin is a heterooctameric complex of proteins HAUS1-8 that organizes branching microtubule nucleation [27,37]. Augmin facilitates γ -TuRC-dependent unidirectional microtubule nucleation from a template microtubule (Figure 1c). Augmin binds to γ -TuRC through an adaptor protein Neural precursor cell expressed developmentally down-regulated protein 1 (NEDD1) [31,38] (Figure 1c).

TPX2 is able to condense on microtubules and concentrate augmin, γ -TuRC and tubulin in patches, which serve as microtubule nucleation spots [30] (Figure 1c). Notably, TPX2's simultaneous binding across longitudinal and lateral tubulin interfaces may be sufficient to overcome the nucleation energy barrier, even in the absence of its liquid–liquid phase separation (LLPS) [39,40].


AKNA was recently identified as a γ -TuRC-interacting protein and a potent activator of centrosomal microtubule nucleation in neuronal progenitors and in normal murine mammary gland (NMG) epithelial cells [32]. Moreover, we have shown its very high potency to boost centrosomal microtubule nucleation in developing rodent neurons [15]. Nevertheless, the molecular mechanism of its action is still unclear.

Microtubule polymerase and plus-end tracking protein CKAP5 (also known as Xmap215 or chTOG) is another key protein that enhances γ -TuRC-dependent microtubule nucleation [33,41] (Figure 1d). Although CKAP5 can stimulate microtubule nucleation even without γ -TuRC [42,43], the kinetics of such nucleation reaction and, remarkably, the properties of the nucleated microtubules then differ from the canonical γ -TuRC-dependent pathway [43]. As CKAP5 plays key roles in neuronal development [44,45], it will be important to distinguish its functions at the microtubule plus and minus ends. Moreover, CKAP5 bundles actin filaments and links them to microtubules [46], which is crucial for microtubule advancement in neuronal growth cones and axonal growth [44].

The interplay of microtubule minus-end proteins in regulating γ -TuRC minus-end capping

CLIP-associating protein 2 (CLASP2) stimulates microtubule nucleation from purified γ -TuRCs [33]. Interestingly, γ -TuRCs are released from the nucleated microtubules by Calmodulin-regulated spectrin-associated protein 2 (CAMSAP2) and CAMSAP3 [33] (Figure 1e). CAMSAPs bind to microtubule minus ends, stabilize them [47] and facilitate their growth [33]. Katanin recognizes longer stretches of CAMSAPs and severs them [48]. CAMSAP-driven minus-end growth of

Figure 1



a subset of protofilaments on the γ -TuRC-attached microtubule leads to γ -TuRC removal and minus-end decapping [33]. Of note, CAMSAPs can also nucleate microtubules independently of γ -TuRC through LLPS [49].

Intriguingly, CDK5RAP2, but neither CKAP5 nor CLASP2, prevents CAMSAP2 and CAMSAP3 binding to the γ -TuRC: microtubule interface and subsequent microtubule decapping (Figure 1e) [33]. Thus, we speculate that CDK5RAP2 might, paradoxically, decrease cellular microtubule mass by preventing minus end decapping and subsequent stabilization by CAMSAPs and also by keeping γ -TuRCs locked on microtubules, making them unavailable for next rounds of microtubule nucleation.

Microtubule nucleation is also influenced by microtubule depolymerases [50,51]. Remarkably, KIF2A is a specific minus-end depolymerase [51]. Its affinity for the plus end is low, allowing unperturbed γ -TuRC-dependent nucleation in its presence. γ -TuRC-capped minus ends are protected from both KIF2C and KIF2A [51]. However, when KIF2A is combined with spastin, they recognize the γ -TuRC: microtubule interface and synergistically remove the γ -TuRC from the minus end, which is then further depolymerized by KIF2A [51] (Figure 1e).

Collectively, described mechanisms of γ -TuRC regulation at microtubule minus ends open new research perspectives on how microtubule networks might be regulated in neurons.

Microtubule nucleation by microtubule-associated proteins (MAPs)

Unexpectedly, microtubule nucleation activity resides in dozens of MAPs [52,53]. In addition, many of the MAPs undergo LLPS, which appears to be a dominant mechanism of their microtubule nucleation activity [52,54] (Figure 1f). In general, LLPS has emerged as an important mechanism of cellular homeostasis and organization [54,55], however, further studies are needed to specify the contributions of LLPS to regulation of

microtubule nucleation *in vivo*, both generally and in neurons specifically.

Outstanding questions

Discussed molecular mechanisms enable investigation of microtubule organization and regulation in neurons on a new level. What proportion of the new minus ends are capped by γ -TuRC vs. by CAMSAPs? What is the interplay between these systems? Are there different neuronal microtubule populations, some uncapped and some capped by various types of γ -TuRC complexes? How do these microtubule populations differ in terms of biophysical properties, abundance, subcellular localization, and among neuronal types? What is the contribution of γ -TuRC-independent microtubule nucleation by diverse MAPs to neuronal morphogenesis and migration *in vivo*? These questions will be central to future research.

Microtubule organizing centers (MTOCs)


In nascent neurons, the dominant MTOC is the centrosome [15] (Figure 2), composed of two microtubule-based centrioles, which are surrounded by intricate protein matrix called the pericentriolar material (PCM) [56,57]. The older, *mother* centriole, bears additional specific proteinaceous structures at its distal end – the subdistal (SDA) [58] and the distal (DA) appendages [59] (Figure 2). Both types of appendages play key roles in ciliogenesis and cilia positioning [58,59], with SDA mediating also microtubule anchoring [58].

The centrosome contains several pools of γ -TuRC (Figure 2), including at the base of the centrioles [56], in the centriolar lumen - anchored via augmin [56,60], in the outer layer of the PCM - anchored via Centrosomal protein of 192 kDa (Cep192)-NEDD1 or A-kinase anchor protein 9 (AKAP9)/Pericentrin-CDK5RAP2 modules [58,60], and at the SDA [58,60]. Remarkably, CKAP5 is necessary for γ -TuRC localization to the PCM and to the SDA [61]. At SDA, γ -TuRC interacts with ninein, important for microtubule anchoring [58], and with AKNA [32] (Figure 2).

Diverse Modes of Microtubule Generation.

(a) The core of the γ -TuRC is formed by GCP2-GCP3, GCP4-GCP5, GCP4-GCP6 pairs and MZT1, which are subsequently joined by actin, four other GCP2-GCP3 γ -Tubulin Small Complexes (γ -TuSCs), and MZT2 [18,28,29]. The γ -TuRC 'splayed' conformation correlates with negligible microtubule nucleation activity, while its partial closure increases it [19–21,26]. (b) CM1 domain dimers of CDK5RAP2 bind to the outer parts of the γ -TuRC via MZT2:GCP2 structural modules and push γ -TuRC towards closed conformations, while simultaneously destabilizing actin in the luminal bridge [28,29]. MZT2 not shown in the schematic. (c) Branching microtubule nucleation depends on augmin and γ -TuRC and can be enhanced by TPX2, which condenses on microtubules and concentrates components of the branching microtubule nucleation pathway [30,37]. NEDD1 seems to be required for augmin: γ -TuRC interaction [31]. (d) CKAP5 acts at both the plus and minus end of microtubules. It binds and stimulates γ -TuRC [41], likely by holding together γ -TuRC and the first tubulin layer, and by stabilization of nascent protofilaments [41]. (e) Interplay of novel mechanisms of microtubule minus end regulation. Microtubule minus ends can be timely protected by CAMSAPs or capped by various types of γ -TuRCs. They either protect or destabilize the minus ends by attracting or repulsing microtubule severases or depolymerizing kinesins, such as spastin, katanin or KIF2A, respectively. (f) Multivalent MAPs bundle microtubules, concentrate tubulin and many have the ability to undergo LLPS on microtubules or elsewhere [52,53], thus likely facilitating γ -TuRC-independent microtubule nucleation.

Figure 2



Neuronal MTOCs.

In nascent neurons, the centrosome is active as MTOC [15]. There are various γ -TuRC pools in and on the centrosome [60]. The Golgi is another important neuronal MTOC; perturbation of Golgi-dependent microtubule nucleation impairs axonal growth [15]. Studies in *Drosophila* and *C. elegans* implicated Rab5+ and Rab11+ endosomes, respectively, as MTOCs generating minus-end-out microtubules in dendrites [64–66]. Endosomal MTOCs still need to be shown in mammalian neurons.

Other neuronal MTOCs are the Golgi apparatus [62], probably some subsets of the Golgi outposts – although this is still unclear [63], endosomes [64–66] (Figure 2), and microtubules themselves – via augmin [67–70] (Figure 1c). The γ -TuRC anchoring and activating protein ensemble at the Golgi is partially different from the centrosomal machinery [71]. Golgi-associated γ -TuRC is bound and activated by the AKAP9-Myomegalin (MMG) protein complex, which is functionally related to Pericentrin-CDK5RAP2. Notably, AKAP9 also recruits CDK5RAP2 [71]. AKAP9 is anchored at the cis-Golgi membrane via Golgin subfamily A member 2 (GM130) [4,71]. Interestingly, only isoform 8 of MMG, but not CDK5RAP2, facilitates CAMSAP2 recruitment to the microtubules nucleated at the Golgi. Golgi-derived microtubules decorated by CAMSAP2 are further stabilized by CLASPs [71] (Figure 2).

Endosomes are a new class of neuronal MTOCs, recently discovered in *Drosophila* [65] and *Caenorhabditis elegans* [64] neurons. In *Drosophila*, Rab5-positive endosomes localize to the dendritic branch points and recruit γ -TuRC via Axin [65]. In *C. elegans*, γ -TuRC is anchored to Rab11-positive endosomes in the growth cone of outgrowing dendrites by a ninein homolog Non-centrosomal microtubule array protein 2 (NOCA-2) [66] (Figure 2). Microtubule nucleation at endosomes is

important for ensuring proper minus-end-out orientation of invertebrate dendritic microtubules (Figure 3).


Augmin guides branching microtubule nucleation in both axons and dendrites [67–70] (Figure 3). Although the importance of augmin for microtubule polarity and density in axons and dendrites was shown [67–70], the direct visualization of branching microtubule nucleation via augmin and its molecular mechanism in neurons has remained undefined.

Microtubule nucleation in the neuronal development space-time

Neuronal delamination

Neurons are born after neurogenic divisions of neuronal progenitors [72]. Depending on the type of the progenitor, some newly born neurons delaminate – disassemble their adhesions to neighboring cells in the differentiating neuroepithelium and retract their centrosome-containing apical tip [73] (Figure 3). This process requires active centrosomal microtubule nucleation and actin contractility, and involves N-cadherin downregulation [73]. While centrosomal Akna directs delamination of neuronal progenitors [32], it might also be involved in the delamination of nascent neurons (Figure 3).

Figure 3



Microtubule Nucleation During Neuronal Development.

A schematic of developing rodent neocortex. Neurons can arise from radial glia progenitors or from intermediate progenitors [72]. There are more modes of axon formation than shown in the schematic. VZ-ventricular zone, SVZ-subventricular zone, IZ-intermediate zone, CP-cortical plate. **1.** A speculative model of neuronal delamination in developing rodent neocortex is shown based on data from different systems [32,73]. **2.** Active centrosomal microtubule nucleation is required for radial migration of projection neurons [7,15]. **3.** Augmin-γ-TuRC complex is key for axonal branching, growth and plus-end-out microtubule polarity in axons [67,68,79]. Augmin-γ-TuRC is inhibited by Centrosomin (CDK5RAP2 homolog) in *Drosophila* [88] and might be influenced by the action of KIF2A (may be in complex with spastin), since KIF2A reduces axonal branching in mice [83]. **4.** γ-TuRC complexes nucleate minus-end-out microtubules at the distal dendrites and at dendrite branching points. Endosomes nucleate minus-end-out microtubules in invertebrates [64–66]. Dysregulation of augmin-γ-TuRC, γ-TuRC itself, and CAMSAPs leads to the reduction of dendritic arbor complexity, length and microtubule mass [68–70,79]. **5.** A speculative model of microtubule mass generation and accumulation of synaptic vesicles in *en passant* synapses by a combined action of augmin-γ-TuRC, spastin and KIF2A. Dashed arrows indicate the direction of depolymerization/degradation.

Neuronal differentiation in mice triggers drastic transcriptional downregulation of several players involved in microtubule nucleation, namely, NEDD1, TPX2 and important kinases such as Cyclin-dependent kinase 1 (CDK1), Serine/threonine-protein kinase (PLK1), and Aurora A [74]. However, AKAP9, MMG, CAMSAP2, and ninein are upregulated [74].

Neuronal alternative splicing generates a dominant non-centrosomal ninein isoform as well as variants of many other proteins, including AKAP9, CLASPs, and KIF2A [75]. While this uncovers that regulators of microtubule nucleation have a surprising diversity and richness in neurons, their functions are still unclear; this new territory is only beginning to be uncovered.

Neuronal polarization

The newly born neurons form dynamic neurites, quickly exploring their surroundings, while moving through the growing vertebrate neocortex [7] (Figure 3). This first step in neuronal polarization varies in different types of neurons, however, it relies on the reorganization of actin and microtubules [5,7]. The impact of microtubule nucleation on neurite formation has remained unclear. Since the centrosome is still active in nascent neurons [15], its localization was frequently studied in relation to the first neurite formation [5]. However, the observed patterns of centrosome behaviour before and during neurite outgrowth in different systems are variable. Moreover, the centrosome moves together with the Golgi, further complicating the interpretation of experiments [5]. Endogenous tagging of various regulators of microtubule nucleation with inducible degrons allows rapid and targeted protein depletion [42]. The employment of this technology in neurons might shed new light on the role of microtubule nucleation and respective MTOCs in the early developmental steps.

Axon specification comes as the next milestone in neuronal development. Here, one of the dynamic neurites acquires initial axonal characteristics, namely, faster and persistent growth and stabilization of microtubule bundles that become progressively unipolar with plus ends pointing towards the growth cone [76,77] (Figure 3). Axon specification depends on microtubule stabilization and bundling [78]. Furthermore, the centrosome has long been implicated in axon specification [11]. However, neither a specific centrosome position, nor nucleation activity is necessary for axon outgrowth, as was recently shown by two alternative approaches: by laser-ablating the centrosome in non-polarized projection neurons and by deactivating its microtubule nucleation capability by displacing centrosomal γ -TuRC with a dominant-negative-acting C-terminal part of NEDD1 [15].

Microtubule nucleation needed for the development of both axons and dendrites depends on the γ -TuRC and

augmin (Figure 3). Their depletion reduces microtubule density [67,68,79]. Proper microtubule density in dendrites and axons is also maintained by CAMSAPs [79], which have a profound impact on neuronal polarity and maturation [48,79,80]. Studies in *C. elegans* indicate that microtubule bundles in neurites are maintained by two main pathways: 1) protein complexes containing γ -tubulin and Ninein (NOCA-2 in *C. elegans*) and 2) CAMSAPs (PTRN-1 in *C. elegans*) [66,81]. Either of these pathways can compensate for the impairment of the other [81]. However, in mammalian neurons, loss of γ -TuRC is accompanied by a simultaneous loss of CAMSAP2-decorated microtubules [79], indicating a tighter link between both pathways.

CAMSAP-driven microtubule minus end decoration may be accomplished by decapping γ -TuRC-capped minus ends [20]. Since there is no augmin in *C. elegans*, it would be interesting to investigate whether the vertebrate γ -TuRC-augmin complexes fulfill similar roles as γ -TuRC-NOCA-2 in *C. elegans*, or whether the γ -TuRC-ninein and γ -TuRC-augmin complexes co-exist and cooperate (or compete) in both developing and mature vertebrate neurons.

Intriguingly, although Aprea et al. reported drastic transcriptional downregulation of TPX2 already in nascent neurons [74], Chen et al. detected TPX2 protein in hippocampal neurons [82], localizing besides cytoplasm also to the centrosome and after three days in culture to undefined filamentous structures. The activity of TPX2 depends on RanGTP and importins and its downregulation decreased neurite length [82]. The presence of sufficient amounts of TPX2 might have major effects on branching microtubule nucleation, both with or without augmin. Hence, it would be exciting to investigate the behaviour of this protein in detail using endogenous tagging *in vivo*.

Neuronal migration

While neurons are polarizing and grow their axon, their cell bodies simultaneously migrate to their final destination in the central nervous system (CNS) [7] (Figure 3). Whereas, for example, interneurons are bipolar, cortical projection neurons undergo complex switching of migratory modes during their journey through the cortex [7]. Both radial and tangential migration require a functional centrosome [7,15].

Only recently, it was uncovered that properly regulated centrosomal microtubule nucleation is essential for directional migration of cortical projection neurons [15]. Modulation of the neuronal centrosome to either downregulate or upregulate centrosomal microtubule nucleation resulted in stalled migration and accumulation of neurons in the subventricular (SVZ) and intermediate zone (IZ) of the developing neocortex [15]. Neurons with dysregulated centrosomes failed to switch

from multipolar to bipolar morphology to start locomoting along the radial glia towards the cortical plate [15]. Both too strong and too weak centrosomal microtubule nucleation disrupts proper cycling of the cytoplasmic dilation in the leading process of the migrating neuron [15], which is essential for radial migration [7].

Strikingly, dysregulation of expression and function of the major neuronal microtubule-depolymerising kinesin-13 KIF2A leads to a similar migratory block [83]. Since KIF2A is present at the centrosome [84] and in light of recent findings linking KIF2A and γ -TuRC [51] (Figure 1e), it is tempting to speculate that KIF2A might primarily regulate microtubule minus ends. This might be important for the cyclic remodelling of the cytoplasmic dilation in migrating neurons. Interestingly, KIF2A together with Centrosomal protein of 170 kDa B (Cep170B) locally antagonize microtubule minus end stabilization by CAMSAPs in epithelial cells [85]. Cep170B interaction with microtubules, which attracts KIF2A and repels CAMSAPs, is regulated by phosphorylation [85]. These novel mechanisms of the cross-talk between different microtubule minus end stabilization/destabilization pathways might have a profound impact on neuronal microtubule networks and would be an exciting new avenue for further studies.

In contrast to the centrosome dysregulation, impairment of microtubule nucleation from the Golgi by specific γ -TuRC-displacement, using a GM130-binding domain of AKAP9, attenuated axon growth, while radial migration was unaffected [15]. It indicates that different microtubule networks (e.g. centrosomal vs. Golgi) in neurons are specialized for different tasks (migration vs. axon growth, respectively).

Downregulation of NEDD1, and upregulation of CAMSAPs, AKAP9, MMG, and the acentrosomal ninein isoform [74,75] likely contributes to the progressive loss of microtubule nucleating and anchoring capacity of the centrosome that occurs during neuronal development [15] (Figure 3). Microtubule nucleation then becomes acentrosomal, with AKAP9/MMG, CAMSAPs, augmin and likely ninein becoming the major regulators. Since the centrosome can limit MT nucleation at other MTOCs [86], its switch-off may be required to ensure the massive growth of axons and dendrites.

Neuronal maturation, branching, synapse formation and regeneration

Microtubule nucleation in dendrites and axons is key not only for generation of microtubule mass, but also for orientation of microtubule bundles, and neurite branching [64]. In general, microtubule bundles in axons are uniformly plus-end-out, while they are of mixed polarity in mammalian dendrites and dominantly minus-end-out in longer branches of invertebrate dendrites [12,87]. Remarkably, minus-end-out microtubules are

located centrally in dendrites, while plus-end-out microtubules reside on the dendrite periphery [12].

A major player in acentrosomal microtubule nucleation in growing axons and dendrites is augmin [67–70] (Figure 3). Its downregulation leads to the loss of microtubule plus-end-out uniformity in axons and to the loss of microtubule mass [67]. The growth of both axons and dendrites is impaired after augmin or γ -TuRC depletion [68]. The importance of augmin for microtubule branching in neurons with complicated dendritic trees was recently shown in two independent elegant studies on *Drosophila* larval class IV dendritic arborization (da) neurons [69,70]. In addition, microtubule nucleation from endosomes in both *Drosophila* [65] and *C. elegans* [64] is crucial for generation of dendritic minus-end-out microtubules. How all these systems (ninein/NOCA-2-dependent vs. CAMSAPs/Patronin/PTRN-1-dependent vs. augmin-dependent) work together to establish the proper microtubule density and orientation needed for building elaborate neuronal morphology remains to be investigated.

Interestingly, *Drosophila* Centrosomin attenuates dendrite branching by suppressing or competing with the augmin pathway [88]. Indeed, CDK5RAP2- or NEDD1-attached γ -TuRCs are distinguished as two different γ -TuRC pools in other systems [15,34]. As described above, CDK5RAP2/Centrosomin might lock the γ -TuRC at the microtubule minus ends and limit its availability for augmin and other pathways. Similarly, KIF2A suppresses excessive collateral axonal branching [83]. It might act together with spastin on the microtubules capped by augmin- γ -TuRC close to or at the neurite branching points and destabilize them (Figure 3).

Augmin- γ -TuRC regulates axonal *en passant* synapses in hippocampal neurons, where localized bouts of microtubule nucleation are crucial for proper inter-bouton synaptic vesicle transport and exocytosis [89]. A recent study on human neurons confirmed that *en passant* presynaptic sites are hotspots of new microtubule growth, necessary for proper synapse function, and identified spastin as a generator of new microtubules [90]. Whether there is a spastin-KIF2A- γ -TuRC-augmin module acting at these presynaptic sites remains to be investigated (Figure 3).

Coupling microtubule nucleation with microtubule severing seems to be a powerful way to increase microtubule mass and turnover. Spastin, possibly acting with KIF2A close to the γ -TuRC: microtubule interface, could recover γ -TuRCs for new rounds of directed nucleation, ensured by augmin. The newly formed microtubule minus ends might be orderly depolymerized by KIF2A [51], increasing the amount of treadmilling microtubules. Spastin might also act at the microtubule shaft, generating shorter microtubules [90]. Thus, molecular

motors carrying synaptic vesicles along microtubules would reach a microtubule end with increased probability, facilitating synaptic vesicles unloading at the pre-synaptic site.

There are several indications that microtubule nucleation is neuroprotective after neuronal injury, however, the underlying mechanisms seem complex and invite further research [6,91]. For axon regeneration, microtubules are central. Upon CNS injury, they depolymerize in rodents and human patients [92,93], while they remain intact and bundled after an injury in the peripheral nervous system (PNS) [92]. Microtubule stabilization leads to axon regeneration in the injured spinal cord [93–97]. Yet, it is still unclear where and how the microtubules are nucleated to replenish the lost microtubule mass [6]. Investigation of the intricate interplay between various microtubule nucleation modes and related microtubule networks in neurons might bring novel insights into how neurons develop and regenerate.

Conclusions and future perspectives

We have learned a great deal about centrosomal and acentrosomal pathways of microtubule nucleation in neurons. Yet, we still lack a more comprehensive models of where, when and how the various microtubule nucleation modules are localized, regulated, and how they compete or work together.

The big task ahead is to disentangle various neuronal MT networks and their regulators. Moreover, the intriguing discrepancy between transcriptional down-regulation of TPX2 and NEDD1 and their reported (TPX2) or implied (Augmin requires NEDD1 for γ -TuRC binding) functioning later during neuronal development awaits reconciliation. The finding that the type of microtubule nucleation might influence microtubule properties [43] is interesting especially in relation to the different features of minus-end-out and plus-end-out dendritic microtubules in mammals [12]. Many important questions remain open for investigation, including dissecting the dominant minus-end protection and regulation mechanisms. Additionally, understanding the contributions of different microtubule nucleation modes—such as LLPS of MAPs, which requires clarification on its occurrence and role *in vivo*—is crucial for comprehending how microtubule mass is generated during neuronal development and maintained in neuronal homeostasis.

Addressing these questions will ultimately help us to enhance the regenerative potential of neurons.

Declaration of competing interest

The authors declare no conflicts of interest.

Acknowledgements

We thank Markéta Schmidt Černohorská, Emily Burnside, Juliane Schiweck, Kai Fährmann and Cedric Böger for reading our manuscript and for discussions. S.V. would like to extend special thanks to Jan Malý for his continuous support.

We apologize to all colleagues whose important contributions could not be cited due to space limitations.

S.V. is supported by Czech Science Foundation grant 25–14392L. F.B. is supported by the International Foundation for Research in Paraplegia, Wings for Life, ERANET AXON REPAIR, and ERANET RATER SCI. F.B. is also funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation): SFB 1089, as well as SFB 1158 and SPP 2395. F.B. is a member of the excellence cluster ImmunoSensation2 (EXC2151–390873048) and the iBehave NRW network. F.B. is a recipient of the Roger de Spoelberch Prize.

Data availability

No data was used for the research described in the article.

References

Papers of particular interest, published within the period of review, have been highlighted as:

- * of special interest
- ** of outstanding interest

1. Gudimchuk NB, McIntosh JR: **Regulation of microtubule dynamics, mechanics and function through the growing tip.** *Nat Rev Mol Cell Biol* 2021, **22**:777–795.
2. Valdez VA, Neahring L, Petry S, Dumont S: **Mechanisms underlying spindle assembly and robustness.** *Nat Rev Mol Cell Biol* 2023, **24**:523–542.
3. Sallee MD, Feldman JL: **Microtubule organization across cell types and states.** *Curr Biol* 2021, **31**:R506–R511.
4. Akhmanova A, Kapitein LC: **Mechanisms of microtubule organization in differentiated animal cells.** *Nat Rev Mol Cell Biol* 2022, **23**:541–558.
5. Higgs VE, Das RM: **Establishing neuronal polarity: microtubule regulation during neurite initiation.** *Oxford Open Neurosci* 2022, **1**, kvac007.
6. Hilton BJ, Griffin JM, Fawcett JW, Bradke F: **Neuronal maturation and axon regeneration: unfixing circuitry to enable repair.** *Nat Rev Neurosci* 2024, **25**:649–667.
7. Jossin Y: **Molecular mechanisms of cell polarity in a range of model systems and in migrating neurons.** *Mol Cell Neurosci* 2020, **106**, 103503.
8. Pinho-Correia LM, Prokop A: **Maintaining essential microtubule bundles in meter-long axons: a role for local tubulin biogenesis?** *Brain Res Bull* 2023, **193**:131–145.
9. Nakajima C, Sawada M, Umeda E, Takagi Y, Nakashima N, Kuboyama K, Kaneko N, Yamamoto S, Nakamura H, Shimada N, et al.: **Identification of the growth cone as a probe and driver of neuronal migration in the injured brain.** *Nat Commun* 2024, **15**:1877.
10. Atkins M, Nicol X, Fassier C: **Microtubule remodelling as a driving force of axon guidance and pruning.** *Semin Cell Dev Biol* 2023, **140**:35–53.

11. Kapitein LC, Hoogenraad CC: **Building the neuronal microtubule cytoskeleton.** *Neuron* 2015, **87**:492–506.
 12. Iwanski MK, Kapitein LC: **Cellular cartography: towards an atlas of the neuronal microtubule cytoskeleton.** *Front Cell Dev Biol* 2023, **11**, 1052245.
 13. Burute M, Kapitein LC: **Cellular logistics: unraveling the interplay between microtubule organization and intracellular transport.** *Annu Rev Cell Dev Biol* 2019, **35**:29–54.
 14. Kellihier MT, Saunders HA, Wildonger J: **Microtubule control of functional architecture in neurons.** *Curr Opin Neurobiol* 2019, **57**:39–45.
 15. Vinopal S, Dupraz S, Alfadil E, Pietralla T, Bendre S, Stiess M, Falk S, Camargo Ortega G, Maghelli N, Tolić IM, et al.: **Centrosomal microtubule nucleation regulates radial migration of projection neurons independently of polarization in the developing brain.** *Neuron* 2023, **111**:1241–1263.e16.
- This study shows how two intertwined dynamic processes - radial migration and axon formation - are selectively regulated through distinct microtubule nucleation systems. Microtubule nucleation at the centrosome is dispensable for axon formation and growth; however, it is required for radial migration of projection neurons. In contrast, disruption of Golgi-mediated microtubule nucleation did not impair radial migration, but decreased axonal growth. This provides the basis for understanding how certain neurodevelopmental disorders affect specifically migration but not axon growth and guidance.
16. Miryala CSJ, Holland ED, Dent EW: **Contributions of microtubule dynamics and transport to presynaptic and post-synaptic functions.** *Mol Cell Neurosci* 2022, **123**, 103787.
 17. Wu JY, Cho S-J, Descant K, Li PH, Shapson-Coe A, Januszewski M, Berger DR, Meyer C, Casingal C, Huda A, et al.: **Mapping of neuronal and glial primary cilia contactome and connectome in the human cerebral cortex.** *Neuron* 2024, **112**: 41–55.e3.
 18. Würtz M, Zupa E, Atorino ES, Neuner A, Böhler A, Rahadian AS, Vermeulen BJAA, Tonon G, Eustermann S, Schiebel E, et al.: **Modular assembly of the principal microtubule nucleator γ -TuRC.** *Nat Commun* 2022, **13**:473.
 19. Aher A, Urnavicius L, Xue A, Neselu K, Kapoor TM: **Structure of the γ -tubulin ring complex-capped microtubule.** *Nat Struct Mol Biol* 2024, **31**:1124–1133.
- The study brings high-resolution cryo-EM structural data of γ -TuRC-capped microtubules, showing the complete γ -TuRC-closure and 13- protofilament matching in agreement with [20].
20. Brito C, Serna M, Guerra P, Llorca O, Surrey T: **Transition of human γ -tubulin ring complex into a closed conformation during microtubule nucleation.** *Science* 2024, **383**:870–876.
- The authors show the changes in γ -TuRC structure during microtubule nucleation in high resolution using cryo-EM. They found that γ -TuRC ring progressively closes with increasing number of nucleated protofilaments and in the end, the γ -TuRC-structure matches the 13- protofilament structure of a microtubule.
21. Vermeulen BJA, Böhler A, Gao Q, Neuner A, Župa E, Chu Z, Würtz M, Jäkle U, Gruss OJ, Pfeffer S, et al.: **γ -TuRC asymmetry induces local protofilament mismatch at the RanGTP-stimulated microtubule minus end.** *EMBO J* 2024, **43**: 2062–2085.
- The authors found that the γ -TuRC activated by Ran-GTP pathway closes only partially, leading to imperfections in inter-protofilament and protofilament- γ -tubulin contacts in the nucleated microtubules. Such structures open several possibilities how microtubule minus ends might be modified by several mechanisms discussed in this review.
22. Liu P, Zupa E, Neuner A, Böhler A, Loerke J, Flemming D, Ruppert T, Rudack T, Peter C, Spahn C, et al.: **Insights into the assembly and activation of the microtubule nucleator γ -TuRC.** *Nature* 2020, **578**:467–471.
 23. Wieczorek M, Urnavicius L, Ti SC, Molloy KR, Chait BT, Kapoor TM: **Asymmetric molecular architecture of the human γ -tubulin ring complex.** *Cell* 2020, **180**:165–175.e16.
 24. Consolati T, Locke J, Roostalu J, Chen ZA, Gannon J, Asthana J, Lim WM, Martino F, Cvetkovic MA, Rappsilber J, et al.: **Microtubule nucleation properties of single human γ TuRCs explained by their cryo-EM structure.** *Dev Cell* 2020, **53**: 603–617.e8.
 25. Zimmermann F, Serna M, Ezquerro A, Fernandez-Leiro R, Llorca O, Lüdgers J: **Assembly of the asymmetric human γ -tubulin ring complex by RUVBL1-RUVBL2 AAA ATPase.** *Sci Adv* 2020, **6**:1–20.
 26. Gao Q, Vermeulen BJA, Würtz M, Shin H, Erdogdu D, Zheng A, Hofer FW, Neuner A, Pfeffer S, Schiebel E: **The structure of the γ -TuRC at the microtubule minus end – not just one solution.** *Bioessays* 2024, **46**, e2400117.
 27. Kraus J, Alfaro-Aco R, Gouveia B, Petry S: **Microtubule nucleation for spindle assembly: one molecule at a time.** *Trends Biochem Sci* 2023, **48**:761–775.
 28. Xu Y, Muñoz-Hernández H, Krutyholowa R, Marxer F, Cetin F, Wieczorek M: **Partial closure of the γ -tubulin ring complex by CDK5RAP2 activates microtubule nucleation.** *Dev Cell* 2024, **59**:3161–3174.e15.
- This interesting ultrastructural study identifies structural motifs of CDK5RAP2 interaction with the γ -TuRC, which leads to its activation by favouring more closed conformation. Back-to-back with [29].
29. Serna M, Zimmermann F, Vineethakumari C, Gonzalez-Rodriguez N, Llorca O, Lüdgers J: **CDK5RAP2 activates microtubule nucleator γ TuRC by facilitating template formation and actin release.** *Dev Cell* 2024, **59**:3175–3188.e8.
- An interesting study independently identifies CDK5RAP2: γ -TuRC interaction sites and the induction of γ -TuRC partial closure by CDK5RAP2. In addition, the authors found that CDK5RAP2 binding is correlated with the loss of actin from γ -TuRC luminal bridge. Back-to-back with [28].
30. Setru SU, Gouveia B, Alfaro-Aco R, Shaevitz JW, Stone HA, Petry S: **A hydrodynamic instability drives protein droplet formation on microtubules to nucleate branches.** *Nat Phys* 2021, **17**:493–498.
 31. Zhang Y, Hong X, Hua S, Jiang K: **Reconstitution and mechanistic dissection of the human microtubule branching machinery.** *J Cell Biol* 2022, **221**, e202109053.
 32. Camargo Ortega G, Falk S, Johansson PA, Peyre E, Broix L, Sahu SK, Hirst W, Schlichthaerle T, De Juan Romero C, Draganova K, et al.: **The centrosome protein AKNA regulates neurogenesis via microtubule organization.** *Nature* 2019, **567**: 113–117.
 33. Rai D, Song Y, Hua S, Stecker K, Monster JL, Yin V, Stucchi R, Xu Y, Zhang Y, Chen F, et al.: **CAMSAPs and nucleation-promoting factors control microtubule release from γ -TuRC.** *Nat Cell Biol* 2024, **26**:404–420.
- The authors of this exciting study revealed novel mechanisms of microtubule minus end regulation by uncovering the interplay of CDK5RAP2, CAMSAPs, and TOG-domain containing proteins CKAP5 and CLASP2 with γ -TuRC.
34. Muroyama A, Seldin L, Lechler T: **Divergent regulation of functionally distinct γ -tubulin complexes during differentiation.** *J Cell Biol* 2016, **213**:679–692.
 35. Yang S, Au FKC, Li G, Lin J, Li XD, Qi RZ: **Autoinhibitory mechanism controls binding of centrosomin motif 1 to γ -tubulin ring complex.** *J Cell Biol* 2023, **222**, e202007101.
- An elegant study identifying the molecular mechanism of CDK5RAP2 autoinhibition.
36. Tovey CA, Tsuji C, Egerton A, Bernard F, Guichet A, de la Roche M, Conduit PT: **Autoinhibition of Cnn binding to γ -TuRCs prevents ectopic microtubule nucleation and cell division defects.** *J Cell Biol* 2021, **220**, e202010020.
 37. Travis SM, Mahon BP, Huang W, Ma M, Rale MJ, Kraus J, Taylor DJ, Zhang R, Petry S: **Integrated model of the vertebrate augmin complex.** *Nat Commun* 2023, **14**:1–10.
 38. Muñoz-Hernández H, Xu Y, Zhang D, Xue A, Aher A, Carmadiel Pellicer A, Walker E, Marxer F, Kapoor TM, Wieczorek M: **Structure of the microtubule anchoring factor NEDD1 bound to the γ -tubulin ring complex.** *bioRxiv* 2024, <https://doi.org/10.1101/2024.11.05.622067>.
- The authors solved the partial structure of γ -TuRC interacting with its key targeting factor NEDD1.
39. Zhang R, Roostalu J, Surrey T, Nogales E: **Structural insight into TPX2-stimulated microtubule assembly.** *Elife* 2017, **6**: 1–22.

40. Roostalu J, Cade NI, Surrey T: **Complementary activities of TPX2 and chTOG constitute an efficient importin-regulated microtubule nucleation module.** *Nat Cell Biol* 2015, **17**: 1422–1434.
41. McManus CT, Travis SM, Jeffrey PD, Zhang R, Petry S: **Mechanism of how the universal module XMAP215 γ -TuRC nucleates microtubules.** *bioRxiv* 2024, <https://doi.org/10.1101/2024.06.03.597159>.
An important study revealing novel structural organization and function of XMAP215 (CKAP5 in mammals) during its interaction with γ -TuRC.
42. Tsuchiya K, Goshima G: **Microtubule-associated proteins promote microtubule generation in the absence of γ -tubulin in human colon cancer cells.** *J Cell Biol* 2021, **220**, e202104114.
43. Zhu Z, Becam I, Tovey CA, Elfarkouchi A, Yen EC, Bernard F, Guichet A, Conduit PT: **Multifaceted modes of γ -tubulin complex recruitment and microtubule nucleation at mitotic centrosomes.** *J Cell Biol* 2023, **222**, e202212043.
This elegant study in *Drosophila* brought into focus partially redundant, but distinct γ -TuRC-dependent and -independent microtubule nucleation pathways in the mitotic spindle. Remarkably, the authors showed that microtubules nucleated by CKAP5 homolog Msps in the absence of γ -TuRC are stabler than in the wildtype.
44. Cammarata GM, Erdogan B, Sabo J, Kayaer Y, Dujava Zdimalova M, Engström F, Gupta U, Senel J, O'Brien T, Sibanda C, et al.: **The TOG5 domain of CKAP5 is required to interact with F-actin and promote microtubule advancement in neurons.** *Mol Biol Cell* 2024, **35**, br24.
This interesting study highlights the ability of CKAP5 to bind actin, impacting the morphological development of neurons.
45. Hahn I, Voelzmann A, Parkin J, Fülle JB, Slater PG, Lowery LA, Sanchez-Soriano N, Prokop A: **Tau, XMAP215/Msps and Eb1 co-operate interdependently to regulate microtubule polymerisation and bundle formation in axons.** *PLoS Genet* 2021, **17**: 1–30.
46. Sabo J, Dujava Zdimalova M, Slater PG, Dostal V, Herynek S, Libusova L, Lowery LA, Braun M, Lansky Z: **CKAP5 enables formation of persistent actin bundles templated by dynamically unstable microtubules.** *Curr Biol* 2024, **34**:260–272.e7.
The authors of this elegant *in vitro* study found that CKAP5 guides formation of actin bundles along microtubules.
47. Akhmanova A, Steinmetz MO: **Microtubule minus-end regulation at a glance.** *J Cell Sci* 2019, **132**:1–7.
48. Buijs RR, Hummel JJA, Burute M, Pan X, Cao Y, Stucchi R, Altelaar M, Akhmanova A, Kapitein LC, Hoogenraad CC: **WDR47 protects neuronal microtubule minus ends from katanin-mediated severing.** *Cell Rep* 2021, **36**, 109371.
49. Imasaki T, Kikkawa S, Niwa S, Saijo-Hamano Y, Shigematsu H, Aoyama K, Mitsuoka K, Shimizu T, Aoki M, Sakamoto A, et al.: **CAMSAP2 organizes a γ -tubulin-independent microtubule nucleation centre through phase separation.** *Elife* 2022, **11**: 1–28.
50. Wieczorek M, Bechstedt S, Chaaban S, Brouhard GJ: **Microtubule-associated proteins control the kinetics of microtubule nucleation.** *Nat Cell Biol* 2015, **17**.
51. Henkin G, Brito C, Thomas C, Surrey T: **The minus-end depolymerase KIF2A drives flux-like treadmilling of γ TuRC-uncapped microtubules.** *J Cell Biol* 2023, **222**, e202304020.
An important *in vitro* study discovering previously unknown interplay of γ -TuRC, KIF2A and spastin.
52. Jijumon AS, Bodakuntla S, Genova M, Bangera M, Sackett V, Besse L, Maksud F, Henriot V, Magiera MM, Sirajuddin M, et al.: **Lysate-based pipeline to characterize microtubule behaviour.** *Nat Cell Biol* 2022, **24**:253–267.
53. Duan D, Lyu W, Chai P, Ma S, Wu K, Wu C, Xiong Y, Sestan N, Zhang K, Koleske AJ: **Abl2 repairs microtubules and phase separates with tubulin to promote microtubule nucleation.** *Curr Biol* 2023, **33**:4582–4598.e10.
This important study highlights the ability of Abl2, both a MAP and a kinase, to condense and drive γ -TuRC-independent microtubule nucleation.
54. Volkov VA, Akhmanova A: **Phase separation on microtubules: from droplet formation to cellular function?** *Trends Cell Biol* 2024, **34**:18–30.
55. Gouveia B, Kim Y, Shaevitz JW, Petry S, Stone HA, Brangwynne CP: **Capillary forces generated by biomolecular condensates.** *Nature* 2022, **609**:255–264.
56. Laporte MH, Gambarotto D, Bertiaux É, Bournonville L, Louvel V, Nunes JM, Borgers S, Hamel V, Guichard P: **Time-series reconstruction of the molecular architecture of human centriole assembly.** *Cell* 2024, **187**:2158–2174.e19.
This landmark study solved the structural changes during centriole biogenesis in ultra-high resolution using ultrastructure expansion microscopy.
57. Lawo S, Hasegan M, Gupta GD, Pelletier L: **Subdiffraction imaging of centrosomes reveals higher-order organizational features of pericentriolar material.** *Nat Cell Biol* 2012, **14**: 1148–1158.
58. Chong WM, Wang WJ, Lo CH, Chiu TY, Chang TJ, Liu YP, Tanos B, Mazo G, Tsou MFB, Jane WN, et al.: **Super-resolution microscopy reveals coupling between mammalian centriole subdistal appendages and distal appendages.** *Elife* 2020, **9**: 1–21.
59. Chang T-J Ben, Hsu JC-C, Yang TT: **Single-molecule localization microscopy reveals the ultrastructural constitution of distal appendages in expanded mammalian centrioles.** *Nat Commun* 2023, **14**:1688.
60. Schweizer N, Lüders J: **From tip to toe - dressing centrioles in γ TuRC.** *J Cell Sci* 2021, **134**, jcs258397.
61. Ali A, Vineethakumari C, Lacasa C, Lüders J: **Microtubule nucleation and γ TuRC centrosome localization in interphase cells require ch-TOG.** *Nat Commun* 2023, **14**:24–26.
The authors of this solid and interesting study discovered that interphase γ -TuRC requires for its localization in the centrosomal PCM and SDAs its potent stimulator CKAP5 (ch-TOG).
62. Mukherjee A, Brooks PS, Bernard F, Guichet A, Conduit PT: **Microtubules originate asymmetrically at the somatic golgi and are guided via Kinesin2 to maintain polarity within neurons.** *Elife* 2020, **9**:1–33.
63. Nguyen MM, McCracken CJ, Milner ES, Goetschius DJ, Weiner AT, Long MK, Michael NL, Munro S: **Rolls MM: γ -Tubulin controls neuronal microtubule polarity independently of Golgi outposts.** *Mol Biol Cell* 2014, **25**:2039–2050.
64. Liang X, Kokes M, Fetter RD, Sallee MD, Moore AW, Feldman JL, Shen K: **Growth cone-localized microtubule organizing center establishes microtubule orientation in dendrites.** *Elife* 2020, **9**: 1–28.
65. Weiner AT, Seebold DY, Torres-Gutierrez P, Folker C, Swope RD, Kothe GO, Stoltz JG, Zalenski MK, Kozlowski C, Barbera DJ, et al.: **Endosomal Wnt signaling proteins control microtubule nucleation in dendrites.** *PLoS Biol* 2020, **18**, e3000647.
66. He L, van Beem L, Snel B, Hoogenraad CC, Harterink M: **PTRN-1 (CAMSAP) and NOCA-2 (NINEIN) are required for microtubule polarity in *Caenorhabditis elegans* dendrites.** *PLoS Biol* 2022, **20**, e3001855.
67. Sánchez-Huertas C, Freixo F, Viais R, Lacasa C, Soriano E, Lüders J: **Non-centrosomal nucleation mediated by augmin organizes microtubules in post-mitotic neurons and controls axonal microtubule polarity.** *Nat Commun* 2016, **7**, 12187.
68. Cunha-Ferreira I, Chazeau A, Buijs RR, Stucchi R, Will L, Pan X, Adolfs Y, van der Meer C, Wolthuis JC, Kahn OI, et al.: **The HAUS complex is a key regulator of non-centrosomal microtubule organization during neuronal development.** *Cell Rep* 2018, **24**: 791–800.
69. Mukherjee A, Andrés Jeske Y, Becam I, Taieb A, Brooks P, Aouad J, Monguillon C, Conduit PT: **γ -TuRCs and the augmin complex are required for the development of highly branched dendritic arbors in *Drosophila*.** *J Cell Sci* 2024, **137**, jcs261534.
In this study, the authors show the importance of augmin for the development of branched dendritic arbors in *Drosophila* da4 neurons. Back to back with [70].

70. Zhang Y, Sung H-H, Ziegler AB, Wu Y-C, Vlais R, Sánchez-Huertas C, Kilo L, Agircan FG, Cheng Y-J, Mourri K, *et al.*: **Augmin complex activity finetunes dendrite morphology through non-centrosomal microtubule nucleation in vivo.** *J Cell Sci* 2024, **137**, jcs261512.

This interesting paper demonstrates that augmin is required for physiological branching of dendritic arbors in *Drosophila* da4 neurons. Back to back with [69].

71. Wu J, de Heus C, Liu Q, Bouchet BP, Noordstra I, Jiang K, Hua S, Martin M, Yang C, Grigoriev I, *et al.*: **Molecular pathway of microtubule organization at the golgi apparatus.** *Dev Cell* 2016, **39**:44–60.
72. Kalebic N, Huttner WB: **Basal progenitor morphology and neocortex evolution.** *Trends Neurosci* 2020, **43**:843–853.
73. Kawaguchi A: **Neuronal delamination and outer radial glia generation in neocortical development.** *Front Cell Dev Biol* 2021, **8**:1–10.
74. Aprea J, Prenninger S, Dori M, Ghosh T, Monasor LS, Wessendorf E, Zocher S, Massalini S, Alexopoulou D, Lesche M, *et al.*: **Transcriptome sequencing during mouse brain development identifies long non-coding RNAs functionally involved in neurogenic commitment.** *EMBO J* 2013, **32**:3145–3160.
75. Zhang X, Chen MH, Wu X, Kodani A, Fan J, Doan R, Ozawa M, Ma J, Yoshida N, Reiter JF, *et al.*: **Cell-type-specific alternative splicing governs cell fate in the developing cerebral cortex.** *Cell* 2016, **166**:1147–1162.e15.
76. Schelski M, Bradke F: **Microtubule retrograde flow retains neuronal polarization in a fluctuating state.** *Sci Adv* 2022, **8**:1–19.

This study uncovers a completely new way of microtubule dynamics: in developing neurites, the whole microtubule array moves retrogradely to the cell body. In the newly formed axon this retrograde movement ceases. Back to back with [77].

77. Burute M, Jansen KI, Mihajlovic M, Vermonden T, Kapitein LC: **Local changes in microtubule network mobility instruct neuronal polarization and axon specification.** *Sci Adv* 2022, **8**, eabo2343.

This study reveals a new way of microtubule dynamics, microtubule retrograde flow. Back to back with [76].

78. Schelski M, Bradke F: **Neuronal polarization: from spatiotemporal signaling to cytoskeletal dynamics.** *Mol Cell Neurosci* 2017, **84**:11–28.
79. Yau KW, van Beuningen SFBB, Cunha-Ferreira I, Cloin BMCC, van Battum EY, Will L, Schätzle P, Tas RP, van Krugten J, Katrukha EA, *et al.*: **Microtubule minus-end binding protein CAMSAP2 controls axon specification and dendrite development.** *Neuron* 2014, **82**:1058–1073.
80. Zhou Z, Xu H, Li Y, Yang M, Zhang R, Shiraishi A, Kiyonari H, Liang X, Huang X, Wang Y, *et al.*: **CAMSAP1 breaks the homeostatic microtubule network to instruct neuronal polarity.** *Proc Natl Acad Sci U S A* 2020, **117**:22193–22203.
81. Wang S, Wu D, Quintin S, Green RA, Cheerambathur DK, Ochoa SD, Desai A, Oegema K: **NOCA-1 functions with γ -tubulin and in parallel to Patronin to assemble non-centrosomal microtubule arrays in *C. elegans*.** *Elife* 2015, **4**, e08649.
82. Chen W-S, Chen Y-J, Huang Y-A, Hsieh B-Y, Chiu H-C, Kao P-Y, Chao C-Y, Hwang E: **Ran-dependent TPX2 activation promotes acentrosomal microtubule nucleation in neurons.** *Sci Rep* 2017, **7**, 42297.

83. Ruiz-Reig N, Hakanen J, Tissir F: **Connecting neurodevelopment to neurodegeneration: a spotlight on the role of kinesin superfamily protein 2A (KIF2A).** *Neural Regen Res* 2024, **19**:375–379.
84. Kodani A, Kenny C, Lai A, Gonzalez DM, Stronge E, Sejourne GM, Isacco L, Partlow JN, O'Donnell A, McWalter K, *et al.*: **Posterior neocortex-specific regulation of neuronal migration by CEP85L identifies maternal centriole-dependent activation of CDK5.** *Neuron* 2020, **106**:246–255.e6.
85. Guan C, Hua S, Jiang K: **The CEP170B-KIF2A complex destabilizes microtubule minus ends to generate polarized microtubule network.** *EMBO J* 2023, **42**, e112953.
86. Gavilan MP, Gandolfo P, Balestra FR, Arias F, Bornens M, Rios RM: **The dual role of the centrosome in organizing the microtubule network in interphase.** *EMBO Rep* 2018, **19**:1–21.
87. Ori-McKenney KM, Jan LY, Jan Y-N: **Golgi outposts shape dendrite morphology by functioning as sites of acentrosomal microtubule nucleation in neurons.** *Neuron* 2012, **76**:921–930.
88. Yalgin C, Ebrahimi S, Delandre C, Yoong LF, Akimoto S, Tran H, Amikura R, Spokony R, Torben-Nielsen B, White KP, *et al.*: **Centrosomin represses dendrite branching by orienting microtubule nucleation.** *Nat Neurosci* 2015, **18**:1437–1445.
89. Qu X, Kumar A, Blockus H, Waites C, Bartolini F: **Activity-dependent nucleation of dynamic microtubules at presynaptic boutons controls neurotransmission.** *Curr Biol* 2019, **29**:4231–4240.e5.
90. Aiken J, Holzbaur ELF: **Spastin locally amplifies microtubule dynamics to pattern the axon for presynaptic cargo delivery.** *Curr Biol* 2024, **34**:1687–1704.e8.

The authors demonstrate that local activity of spastin is key for increased microtubule dynamics in *en passant* synapses. While the authors did not observe augmin- γ -TuRC complexes in their system as shown previously [89], they corroborated that *en passant* synapses are hotspots of increased microtubule polymerization.

91. Rolls MM, Thyagarajan P, Feng C: **Microtubule dynamics in healthy and injured neurons.** *Dev Neurobiol* 2021, **81**:321–332.
92. Ertürk A, Hellal F, Enes J, Bradke F: **Disorganized microtubules underlie the formation of retraction bulbs and the failure of axonal regeneration.** *J Neurosci* 2007, **27**:9169–9180.
93. Ruschel J, Hellal F, Flynn KC, Dupraz S, Elliott DA, Tedeschi A, Bates M, Sliwinski C, Brook G, Dobrindt K, *et al.*: **Systemic administration of epothilone B promotes axon regeneration after spinal cord injury.** *Science* 2015, **348**:347–352.
94. Hellal F, Hurtado A, Ruschel J, Flynn KC, Laskowski CJ, Umlauf M, Kapitein LC, Strikis D, Lemmon V, Bixby J, *et al.*: **Microtubule stabilization reduces scarring and causes axon regeneration after spinal cord injury.** *Science* 2011, **331**:928–931.
95. Ruschel J, Bradke F: **Systemic administration of epothilone D improves functional recovery of walking after rat spinal cord contusion injury.** *Exp Neurol* 2018, **306**:243–249.
96. Griffin JM, Hingorani Jai Prakash S, Bockemühl T, Benner JM, Schaffran B, Moreno-Manzano V, Büschges A, Bradke F: **Rehabilitation enhances epothilone-induced locomotor recovery after spinal cord injury.** *Brain Commun* 2023, **5**:1–19.
97. Stern S, Hilton BJ, Burnside ER, Dupraz S, Handley EE, Gonyer JM, Brakebusch C, Bradke F: **RhoA drives actin compaction to restrict axon regeneration and astrocyte reactivity after CNS injury.** *Neuron* 2021, **109**:3436–3455.e9.