



Increased variability in response to transcranial direct current stimulation in healthy older compared to young adults: A systematic review and meta-analysis

Anna Elisabeth Fromm^a, Catalina Trujillo-Llano^a, Ulrike Grittner^{b,c}, Marcus Meinzer^a, Agnes Flöel^{a,d}, Daria Antonenko^{a,*}

^a Department of Neurology, Universitätsmedizin Greifswald, Greifswald, Germany

^b Berlin Institute of Health (BIH), Berlin, Germany

^c Charité – Universitätsmedizin Berlin, Humboldt-Universität zu Berlin, Berlin Institute of Health, Institute of Biometry and Clinical Epidemiology, Berlin, Germany

^d German Centre for Neurodegenerative Diseases (DZNE) Standort Greifswald, Greifswald, Germany

ARTICLE INFO

Keywords:

Aging
Variability
Stimulation effect
Transcranial electrical stimulation (tES)
Transcranial direct current stimulation (tDCS)
Meta-analysis

ABSTRACT

Background: Healthy aging is associated with a decline in cognitive and motor functions, affecting daily activities and quality of life. Combining transcranial direct current stimulation (tDCS) with behavioral training may be a promising intervention against this decline. However, individual response variability may obscure group-level effects and mislead conclusions about tDCS efficacy. Quantifying this variability is crucial for accurately assessing stimulation effects and understanding individual response factors, like age. Yet, no study has quantitatively compared tDCS variability across age groups. This systematic review and meta-analysis examine age-related variability in cognitive and motor responses to tDCS.

Methods: Following PRISMA guidelines, we searched PubMed and Cochrane for studies directly comparing young and healthy older adults under similar experimental conditions. Across 19 studies comprising 390 older adults (mean \pm SD age: 67 \pm 5 years) and 384 young adults (mean \pm SD age: 24 \pm 3 years) receiving transcranial direct current (tDCS), we quantified behavioral variability using the log-transformed coefficient of variation ratio (lnCVR).

Results: Results revealed substantially higher response variability in healthy older compared to young adults during active tES (21 %, lnCVR_{active} = -0.24 [-0.43, -0.04], p = 0.02), but not during sham conditions (lnCVR_{sham} = -0.18 [-0.42, 0.05], p = 0.13).

Conclusion: These findings provide the first quantitative evidence that advanced age increases behavioral tDCS response variability, highlighting the need to develop personalized tDCS approaches to optimize their efficacy in older populations.

1. Introduction

Healthy aging is associated with declines in cognitive and motor functions, impacting everyday activities and quality of life [1–3]. Cognitive decline is evident in domains such as memory, executive functions, attention, and processing speed [1,4,5], while motor impairment manifests in reduced accuracy, slower movement, and increased risk of falling [6,7]. Transcranial direct current stimulation (tDCS) has emerged as a promising, noninvasive and cost-effective approach to counteract age-related functional decline [8,9]. This

technique applies direct low-intensity current (typically 1–2 mA) via two or more scalp-attached electrodes to modulate neuronal excitability and promote synaptic plasticity by altering membrane potentials [10, 11]. While tDCS does not directly induce action potentials, it can influence network-level activity and facilitate long-lasting changes in synaptic strength through plasticity-related mechanisms [12–15]. When applied during cognitive or motor tasks, tDCS has been shown to enhance performance both in young and older adults [3,6,16,17]. These findings highlight its potential as a compensatory intervention [18,19]. However, tDCS effects are inconsistent across studies, particularly in

* Corresponding author. Department of Neurology, Universitätsmedizin Greifswald, Ferdinand-Sauerbruch-Straße, 17475, Greifswald, Germany.

E-mail address: daria.antonenko@med.uni-greifswald.de (D. Antonenko).

<https://doi.org/10.1016/j.brs.2025.06.005>

Received 18 March 2025; Received in revised form 21 May 2025; Accepted 6 June 2025

Available online 8 July 2025

1935-861X/© 2025 Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

older adults [20,21]. Several studies report only small or even a lack of stimulation effects [22–24], while others suggest that older adults experience greater performance gains than young adults [3,25,26]. This divergence may be because individuals with lower baseline performance, having more room for improvement, often gain greater benefits from tDCS interventions than those with initially higher performance levels [27–29]. However, older adults may exhibit reduced plasticity than young adults, diminishing their responsiveness to both training-based and tDCS approaches [23,30].

In addition to conflicting findings between studies, responses to tDCS vary widely between individuals [31–34]. Interindividual differences in brain anatomy and function likely influence this variability. The former affects the current flow to the target regions for tDCS, hence its functional network organization and activity [35,36]. Such response variability can obscure group-level effects, potentially leading to the

incorrect assumption that tDCS lacks efficacy when, in fact, individual differences may drive mixed outcomes [31,37]. In some cases, this variability can lead to highly significant effects, driven by a subset of individuals who exhibit exceptionally strong responses, even though the majority experience minimal or no benefit [33,38]. Quantifying this response variability is a crucial step in accurately evaluating tDCS efficacy. In other therapeutic contexts, such as antipsychotic treatment for schizophrenia, this variability quantification can enable the identification of responder subgroups [39,40], potentially supporting the development of personalized or stratified treatment approaches [41]. Variability measurement provides insight into the consistency of an intervention's effects across individuals, helps to identify factors that influence differential outcomes like age and might guide the refinement of interventional strategies [39]. Notably, reduced response variability often reflects enhanced treatment effectiveness [39,42].




Fig. 1. PRISMA flowchart for the study identification procedure.

So far, no study has systematically examined and quantified age-related variability in response to tDCS. Greater variability within an age group would indicate more heterogeneous responses due to underlying biological factors, allowing a quantitative assessment of this hypothesized variability for the first time. To address this gap, we conducted a systematic review [43] to identify studies directly comparing young and older adults across motor and cognitive domains within the same experimental setup. Then, we quantified cognitive and motor performance variability using the log-transformed coefficient of variation ratio (lnCVR) to systematically assess age-related differences in tDCS response [44,45]. We hypothesized that age-related variability would differ substantially between young and older adults, with older adults exhibiting higher variability. Moreover, we hypothesized that this difference would be more pronounced in active tES than sham stimulation, allowing us to identify a unique factor distinct from any general differences in behavioral training effectiveness between age groups.

2. Methods

2.1. Search strategy and study selection

A comprehensive literature search was conducted in the PubMed and Cochrane databases from January to August 2024. The used search string was: “(transcranial electrical stimulation OR tES OR transcranial direct current stimulation) AND (old OR elder OR older) AND (young).” The reference lists of the retrieved studies were also checked for additional eligible studies. A total of 411 studies were identified through the database search, with one additional study found through reference checking (see Fig. 1).

Studies were included if they met the following criteria: (a) applied tDCS in young (YA, between 18 and 35 years) and older adults (OA, ≥ 60 years) using identical stimulation parameters and tasks, (b) involved only healthy participants to minimize variability related to pathological conditions, (c) included both an active (treatment) and a sham (placebo) group, (d) examined motor and/or cognitive functions operationalized by behavioral outcomes (e) single- and multisession tDCS studies, (f) that employed either cross-over or between-subject designs, (g) were published in peer-reviewed journals, (h) were written in English, and (i) were not case reports, case series, reviews or opinion pieces.

2.2. Data extraction

Following the PRISMA guidelines [43], two independent researchers (A.E.F and D.A.) conducted the literature search, study assessment and data extraction. We extracted the mean and standard deviation for motor and cognitive outcomes for both the active and sham stimulation conditions post-intervention, and the sample size. When numerical values were not provided in the manuscript, the “metaDigitise” R package (version 1.0.1) [46] was used to extract values from figures. Otherwise, authors were contacted directly (4, with all responding). Additionally, we extracted information on the study design, participant characteristics, stimulation parameters and electrode placement (Table 1). The final sample included 19 studies: 7 investigated motor and 12 cognitive functions. Please note that we used additional information from the following 2023 publication [16] to extract the contrast between older and young adults for Meinzer and colleagues (2012). The systematic review was not pre-registered.

2.3. Age-related response variability

Using the formula below, we calculated the log coefficient of variation ratio (lnCVR), which adjusts for mean differences between groups [39,44,45]. A lnCVR below 0 suggests higher variability in older compared to young adults, a lnCVR of 0 implies equal variability between young and older adults, and a value above 0 suggests lower variability in older compared to young adults. To facilitate

interpretation, we converted the lnCVR into a percentage-based variability score using an exponential function [45].

$$\ln\text{CVR} = \ln \frac{\frac{s_{\text{young}}}{\bar{x}_{\text{young}}}}{\frac{s_{\text{old}}}{\bar{x}_{\text{old}}}} + \frac{1}{2(n_{\text{young}} - 1)} - \frac{1}{2(n_{\text{old}} - 1)}$$

Note. lnCVR = Log coefficient of variation ratio; s = Standard deviation; \bar{x} = Mean; n = sample size.

We pooled the lnCVR across all domains and conditions, with contrasting young and older adults to account for age-related variability in response to active and sham tDCS conditions. For additional exploratory analyses, we performed meta-regressions using mixed-effects models, including the task domain (cognitive or motor) as moderator in the lnCVR analysis. This approach allowed us to specifically assess the effect of task domain on response variability. We report the moderator estimates along with a 95 % confidence interval and the corresponding p-value.

We used the Higgins and Green I^2 test to assess heterogeneity, which quantifies heterogeneity as a percentage from 0 % to 100 %. According to guidelines, I^2 values of 25 %, 50 %, and 75 % correspond to low, moderate, and high heterogeneity levels, respectively [47,48].

2.4. Variability and stimulation effect within age groups

We conducted additional analyses within each age group. To investigate the link between variability and stimulation effects across different age groups, we calculated the lnCVR and the standardized mean difference (SMD) by comparing active and sham conditions for each study. A lnCVR below 0 indicates reduced variability under active stimulation, values around 0 indicate no difference, and values above 0 indicate increased variability under active stimulation. SMD values below 0 indicate a reduced effect under active stimulation, 0 reflects no difference, and values above 0 indicate a higher effect under active stimulation. Finally, we used the extracted effect sizes for a Pearson correlation analysis to explore potential associations between variability and stimulation effects within each age group.

All analyses were conducted in R (4.4.1, <https://www.R-project.org/>), using the “metafor” package for lnCVR and SMD analyses [45,49]. All effects are reported with 95 % confidence intervals.

3. Results

3.1. Descriptive statistics

A total of 19 studies were included in this meta-analysis, with 7 studies investigating motor functions [24,50–55] and 12 studies examining cognitive functions [29,56–66]. Motor functions included learning new skills [24], eye-movement testing [50] or visuomotor tracking [54], Go/NoGo tasks [51,53], postural control tasks, [52], proprioceptive assessment task [55] and treadmill walking [63]. Cognitive functions included memory paradigms, such as language learning [29,58,60,64,66,67], visual working memory tasks [56] or naming tasks [57,61,62,65].

Two were multisession studies [24,29], while the remaining studies applied active tDCS once. Five studies were between-subject studies [24,29,59,63,64] and the remaining 19 were cross-over designs. All participants were healthy, with no history of neurological or psychiatric disorders or use of psychoactive medications. For older adults, normal cognitive functioning was confirmed using standard screening tools (e. g., Montreal Cognitive Assessment, Mini-Mental-Status-Test) with established cut-off scores.

Overall, 774 subjects were included, comprising 384 young adults (mean \pm SD age: 24 ± 3 years) and 390 old participants (67 ± 5 years). For an overview of the study characteristics of the included studies, see Table 1. For further information on the extracted mean and standard deviation, see Table S1.

Table 1
Study characteristics of included studies for quantitative synthesis.

Study	Parameters	Design and Domain	Task	Outcome	Sample
Study 1: Arciniega et al., 2018 10.3389/fnagi.2018.00057	2 mA	Cross	visual working memory	accuracy (%)	Young
	20 min frontoparietal cortex	Single-session Cognitive			22 ± 3 Old
Study 2: Chen & Machado 2017 10.16910/jemr.10.3.5	1 mA	Cross	eye movement testing (prosaccade + antisaccade blocks)	RT (ms)	Young
	10 min dorsolateral prefrontal cortex	Single-session Motor			23 ± 1 Old
Study 3: Conley et al., 2016 10.3389/fnhum.2016.00384	1 mA	Cross	Go/NoGo task	RT (ms)	Young
	20 min motor cortex	Single-session Motor			22 ± 3 Old
Study 4: Craig & Dumas 2017 10.1371/journal.pone.0170331	2 mA	Cross	postural control task	length (cm)	Young
	20 min primary motor cortex	Single-session Motor			21 ± 3 Old
Study 5: Fertoni et al., 2014 10.3389/fnagi.2014.00131	2 mA	Cross	picture-naming task	RT (ms)	Young
	5 min dorsolateral prefrontal cortex	Single-session Cognitive			22 ± 1 Old
Study 6: Fiori et al., 2017 10.1016/j.bbr.2016.12.044	2 mA	Cross	language learning (pseudoword-picture pairs)	correct response (%)	Young
	20 min temporal cortex	Single-session Cognitive			29 ± 6 Old
Study 7: Fujiyama et al., 2022 10.1016/j.neurobiolaging.2021.09.014	1.5 mA	Cross	Flashing grid task combined with perceptual decision-making task and Stop Signal Task	RT (ms)	Young
	20 min pre-supplementary motor area	Single-session Motor			25 ± 5 Old
Study 8: Goodwill et al., 2015 10.1016/j.clinph.2015.01.006	1 mA	Cross	visuomotor tracking task	MEP (%)	Young
	15 min motor cortex	Single-session Motor			26 ± 2 Old
Study 9: Habich et al., 2020 10.1155/2020/8896791	1 mA	Cross	verbal episodic memory task	correct response (#)	Young
	20 min dorsolateral prefrontal cortex	Single-session Cognitive			25 ± 3 Old
Study 10: Kaminski et al., 2021 10.1038/s41598-021-82275-4	1 mA	Between	arc pointing task	accuracy (%)	Young
	20 min primary motor cortex	Multisession Motor			28 ± 4 Old
Study 11: Leach et al., 2019 10.1093/geronb/gby003	1.5 mA	Between Single-session	Associative memory (face-name) task	correct response (SDT)	Young
	25 min dorsolateral prefrontal cortex	Cognitive			23 ± 5 Old
Study 12: Manenti et al., 2013 10.3389/fnagi.2013.00049	1.5 mA	Cross	verbal episodic memory task	RT (ms)	Young
	6 min dorsolateral prefrontal cortex	Single-session Cognitive			24 ± 3 Old
Study 13: Martin et al., 2017 10.1162/jocn_a_01166	1 mA	Cross	semantic word generation task	errors (#)	Young
	30 min primary motor cortex	Single-session Cognitive			27 ± 4 Old
Study 14: Meinzer et al., 2012 10.1523/JNEUROSCI.4812-11.2012 Meinzer et al., 2013 10.1523/JNEUROSCI.5743-12.2013	1 mA	Cross	semantic word generation task	RT (ms)	Young
	20 min inferior frontal gyrus	Single-session Cognitive			27 ± 4 Old
Study 15: Muffel et al., 2019 10.3389/fnagi.2019.00264	1 mA	Cross	proprioceptive assessment task (KINARM)	error (degree)	Young
	15 min primary somatosensory cortex	Single-session Motor			27 ± 3 Old
Study 16: Orcioli-Silva et al., 2021 10.3389/fnagi.2021.739998	0.6 mA	Between Single-session	Treadmill walking task	errors (%)	Young
	20 min prefrontal cortex and vertex	Cognitive			21 ± 3 Old
Study 17: Perceval et al., 2020 10.1016/j.bandl.2020.104788	1 mA	Between	language learning (pseudoword-picture pairs)	correct response (#)	Young
	20 min inferior frontal gyrus	Multisession Cognitive			22 ± 4 Old

(continued on next page)

Table 1 (continued)

Study	Parameters	Design and Domain	Task	Outcome	Sample
Study 18: Peter et al., 2019 10.1007/s00429-019-01946-1	1 mA 15 min dorsolateral prefrontal cortex	Between Single-session Cognitive	verbal episodic memory task	correct response (#)	Young 25 ± 3 Old 69 ± 6
Study 19: Ross et al., 2011 10.3389/fnagi.2011.00016	1.5 mA 15 min anterior temporal lobe	Cross Single-session Cognitive	name recall task	correct response (%)	Young 19–37 Old 65 ± 4

Note. The table summarizes the characteristics of the studies for quantitative synthesis, including key parameters and outcomes for the transcranial direct current stimulation (tDCS) intervention. The reference, including the DOI, is listed in the “study.” “Parameters” contains information regarding the stimulation intensity (in mA), stimulation duration (in minutes) and the stimulation location. “Design and Domain” contain information regarding the study design, hence cross-over studies (cross) or between-subject studies (between), the number of sessions (single- or multisession studies) and the task domain (cognitive or motor). More information on the paradigm can be found in “Task” and the corresponding extracted variable in “Outcome.” The sample describes the age distribution of the young and older participants. The Age is described with mean ± standard deviation or age range if there was no information on the mean.




Fig. 2. Age-related response variability. Left panel: Forest plot illustrates the log coefficient of variation ratio (lnCVR) comparing the active young sample to the active older sample. Values below 0 indicate greater variability in the older sample under active transcranial direct current stimulation (tDCS). Effects are displayed as squares (black representing motor domain, while white represent cognitive domain) with 95 % confidence intervals. The overall effect was: $\text{lnCVR}_{\text{active}} = -0.24$ [-0.43, -0.04], $p = 0.02$. The lnCVR was converted into a percentage-based variability score using an exponential function to facilitate its interpretation: 21.3 % (95 %CI: [4.3, 35.2]). Additional analyses on the task domain revealed a more consistent pattern of age-related variability for motor tasks respectively ($\text{lnCVR}_{\text{active_motor}} = -0.19$, 95 % CI: [-0.38, 0.00], $p = 0.053$; $\text{lnCVR}_{\text{active_cognition}} = -0.25$, 95 % CI: [-0.56, 0.06], $p = 0.11$). Right Panel: Forest plot illustrates the lnCVR comparing the sham young sample to the sham older sample. Values below 0 indicate greater variability in the older sample under sham tDCS. Effects are displayed as squares (black representing motor domain, while white represents cognitive domain) with 95 % confidence intervals. The overall effect was: $\text{lnCVR}_{\text{sham}} = -0.18$ [-0.42, 0.05], $p = 0.13$. Additional analyses on the task domain revealed no difference between motor and cognitive tasks ($\text{lnCVR}_{\text{sham_motor}} = -0.04$, 95 % CI: [-0.23, 0.15], $p = 0.70$; Cognition: $\text{lnCVR}_{\text{sham_cognition}} = -0.27$, 95 % CI: [-0.63, 0.08], $p = 0.13$). Studies were ordered according to the degree of variability under active tDCS. tDCS, Transcranial direct current stimulation. lnCVR, log coefficient of variation ratio.

3.2. Age-related response variability

Under active tDCS, older adults exhibited higher variability compared to young adults ($\ln\text{CVR}_{\text{active}} = -0.24 [-0.43, -0.04]$, $p = 0.02$), although moderate heterogeneity was detected ($I^2 = 62.30\%$ [31.21, 83.93]). This finding represents a 21 % higher variability in older adults compared to young adults (95 % CI: [4.3, 35.2]). In contrast, no age-related difference in variability was observed under the sham condition ($\ln\text{CVR}_{\text{sham}} = -0.18 [-0.42, 0.05]$, $p = 0.13$). Again, the heterogeneity was moderate ($I^2 = 73.39\%$ [50.56, 87.51]), indicating that the observed variability may differ across studies. See Fig. 2 for the forest plots.

Subgroup analyses on task domain (cognitive versus motor) revealed a more reliable pattern of increased age-related variability in motor tasks compared to cognitive tasks (for more information see Supplements).

To further assess potential publication bias, we performed the Egger's test, which indicated no significant funnel plot asymmetry ($z = 0.5$, $p = 0.62$), with the intercept estimate supporting symmetry ($b = -0.38$, 95 % CI: [-0.98, 0.22]). Complementary Trim and Fill analysis identified five potentially missing studies on the left side of the funnel plot, suggesting underrepresentation of smaller studies with negative or null effects. After adjustment, the effect size remained directionally consistent and slightly stronger ($\ln\text{CVR}_{\text{active}} = -0.41$, 95 % CI: [-0.63, -0.19], $p = 0.0002$), indicating that any publication bias may have led to an underestimation of age-related variability. High heterogeneity persisted ($I^2 = 75.5\%$), indicating that methodological variation across studies was the likely source of variability rather than publication bias (see Supplementary Fig. 1 for the funnel plot).

3.3. Variability and stimulation effect within age groups

In older adults, effect and variability did not differ substantially between active tDCS and sham (effect: $\text{SMDOA} = 0.14 [-0.02, 0.31]$, $p = 0.09$, variability: $\ln\text{CVR}_{\text{OA}} = -0.07 [-0.18, 0.05]$, $p = 0.27$), showing a small heterogeneity (effect: $I^2 = 24.95\%$ [0, 73.28] variability: $I^2 = 0\%$ [0, 66.84]). In young adults, effect and variability differed between active tDCS and sham (effect: $\text{SMDYA} = 0.27 [0.1, 0.45]$, $p = 0.02$, variability: $\ln\text{CVR}_{\text{YA}} = -0.11 [-0.22, 0.01]$, $p = 0.06$), showing moderate heterogeneity (effect: $I^2 = 32.16\%$ [0, 80.72], variability: $I^2 = 0\%$ [0, 46.17]). Overall, a higher stimulation effect was associated with reduced variability in both age groups, but was statistically significant only for older adults ($r_{\text{OA}} = -0.78 [-0.77, -0.03]$, $p = 0.04$, $r_{\text{YA}} = -0.38 [-0.71, 0.08]$, $p = 0.1$).

4. Discussion

This meta-analysis aimed to systematically assess age-related differences in tDCS response variability across motor and cognitive domains. Our results revealed that older adults exhibited substantially higher response variability than young adults during active, but not sham, tDCS. This effect was consistent across motor and cognitive domains. Moreover, lower response variability was associated with enhanced treatment efficacy. These results highlight the critical need to understand and address variability in tDCS applications to maximize their therapeutic potential.

Research investigating response variability in noninvasive brain stimulation remains limited [41,68]. Our findings diverge from previous meta-analyses, which indicated minimal response variability under active stimulation for disease-related outcome variables in psychiatric populations [41] or showed no variability in response inhibition among healthy and pathological participants [68]. Such discrepancy may arise from differences in study populations [41] and methodological focus, as previous analyses emphasized task type and stimulation protocols rather than biological factors such as age, sex and health status [68]. Homan et al. (2021) investigated treatment variability in psychiatric disorders

by using questionnaires focused on symptom reduction, primarily for transcranial magnetic stimulation (TMS, which comprised 82 % of the included studies) and tDCS. Most participants received additional medication. Their results showed consistently low variability in treatment effects, except for schizophrenia, suggesting a potential need for personalized approaches in pathological populations. However, they did not consider moderators such as age or sex, which limited their ability to identify biological contributors to variability [41]. Similarly, a recent meta-analysis examined tDCS effects on response inhibition and found no difference in variability between active tDCS and sham conditions, indicating that inter-individual variability may not significantly contribute to the heterogeneity observed in this domain [68]. Again, this study primarily focused on methodological variations, such as differences in stimulation intensity and duration, rather than on biological characteristics of participants, like age and sex [69].

In contrast, our findings revealed an increased variability with age, which likely reflects an interaction between stimulation effects and age-related structural and functional brain changes [3,23,70,71]. Compared to young adults, older individuals may show greater within-group differences in skull thickness, sulcal morphology, and brain tissue composition, which can result in more variable electric field distributions under identical stimulation settings [35,36,72]. For example, increased cerebrospinal fluid (CSF) volume disperses current away from the cortex, reducing field intensity at the target site [36,69,73] and also creates more diffuse and inconsistent field patterns in older adults [36, 69]. Moreover, CSF amplifies the impact of sulcal morphology on current flow, further increasing current-flow variability between individuals [35]. This interaction between stimulation application and age-related anatomical heterogeneity may amplify variability in stimulation-induced behavioral outcomes in older adults, challenging the reliability and effectiveness of standard stimulation protocols [74, 75].

Domain-specific variability in stimulation outcomes likely reflects age-related neurobiological changes that alter brain network function. Structural decline is often accompanied by functional reorganization to maintain performance, often through compensatory mechanisms such as bilateral activation or the recruitment of alternative networks [76–78]. This reorganization complicates the identification of optimal stimulation targets [72,79], as – in addition to more variable neural networks in older adults – older compared to young adults may engage different networks for the same task [23,80]. Age-related differences in network integration are further supported by studies showing that identical tDCS targets can produce opposing changes in functional connectivity across different age groups [16,61,81]. Moreover, individual connectivity profiles have been linked to behavioral outcomes in older adults [12,82]. These findings underscore the importance of individualized stimulation approaches, which may be informed by functional connectivity analyses [81,83,84].

An optimal approach to enhance performance more uniformly in older adults likely depends on multiple interacting factors, including cognitive domain, baseline performance, and individual network integrity [3,23,36]. For instance, individuals with lower baseline learning capacity may derive a greater benefit from stimulation, highlighting the relevance of stratifying participants based on pre-intervention performance metrics [29,30,85]. In addition, domain-specific compensatory strategies, such as increased reliance on executive control in memory tasks, may shape not only behavioral outcomes but also the neural susceptibility to tDCS modulation [3,86, 87]. Further fluctuations in alertness, attentional engagement, and general health status dynamically influence cortical excitability and, consequently, the brain's responsiveness to stimulation [32,88]. For instance, older adults may show more variability in tDCS responses because they often experience fluctuations in attention and vigilance [4, 88,89]. Hence, the interaction of state-dependent factors and tDCS adds another layer of complexity to its application across diverse populations [37,88,90].

In summary, our meta-analysis demonstrated that tDCS induces more heterogeneous responses in older than young adults across motor and cognitive domains. Differences in neuroanatomy, functional reorganization, and brain state may drive this increased variability between age groups. Addressing these multifaceted challenges will require innovative, personalized approaches that integrate anatomical, functional, and state-dependent factors to optimize tDCS efficacy in aging populations [36,81].

4.1. Limitations and future directions

Some limitations of our study should be considered. The limited number of studies directly comparing young and older adults within the same experimental design constrains our sample size and analytical scope. The limitation precluded detailed subgroup analyses of different tDCS modalities and paradigms. Direct comparisons are crucial for isolating age-related effects on tDCS response, as they help control for confounding factors such as variations in study design, stimulation parameters (e.g., intensity, duration, electrode positioning), and task paradigms [32,34,91]. In addition, our analysis suggested the possibility of publication bias, as indicated by a small number of potentially missing studies. However, adjusting for this using a trim-and-fill method did not alter the direction of our findings, supporting the robustness of the main results. The absence of sex-stratified outcome data in the included studies limited our ability to examine potential sex differences in tDCS responsiveness. Anatomical factors such as skull thickness and porosity, which differ between sexes, can influence current distribution and affect stimulation efficacy [69,92]. Future research should address this gap by reporting outcomes separately for males and females and incorporating sex as a key biological variable. Additionally, research with larger datasets is needed to explore how stimulation modalities and task domains influence variability in response patterns across age groups, providing deeper insights into the factors driving age-related differences in tDCS efficacy.

This study offers a novel contribution as the first meta-analysis to systematically quantify age-related variability in tDCS responses. Our findings suggest that lower response variability is generally linked to greater stimulation efficacy, aligning with theoretical assumptions [23, 35]. This may indicate that tDCS induces a more uniform and robust response in young adults, potentially explaining age-related differences in variability, though further empirical validation is needed. We encourage future research to adopt the lnCVR as a variability effect size in meta-analyses across cognitive domains such as working memory, attention, and learning. To reduce response variability in older adults, future research should investigate personalized strategies that consider age-related anatomical and functional changes. One approach might be to increase electric field strength to compensate for atrophy and altered current distribution in aging brains [36,82,93]. Increasing regional precision through individualized electrode positioning may further improve accuracy in targeting and stimulation effectiveness [94,95]. Additionally, combining structural modeling with functional connectivity analyses may help to identify task-relevant network nodes for more effective stimulation in older adults [12,83,84].

5. Conclusion

Our findings showed that behavioral responses during active tDCS were more variable in older than young participants, providing the first quantitative evidence for previous theoretical frameworks on increased response variability in older adults. Our study highlights the importance of personalized approaches to optimizing tDCS interventions, particularly in aging populations, to increase their benefit.

CRedit authorship contribution statement

Anna Elisabeth Fromm: Writing – original draft, Visualization,

Methodology, Investigation, Formal analysis, Conceptualization. **Catalina Trujillo-Llano:** Writing – review & editing, Writing – original draft, Visualization. **Ulrike Grittner:** Writing – review & editing, Methodology, Formal analysis. **Marcus Meinzer:** Writing – review & editing, Conceptualization. **Agnes Flöel:** Writing – review & editing, Conceptualization. **Daria Antonenko:** Writing – review & editing, Supervision, Project administration, Funding acquisition, Conceptualization.

Availability of data and materials

The data supporting this study's findings are available from the corresponding authors upon reasonable request.

Funding

The project was funded by the Deutsche Forschungsgemeinschaft (DFG, project number: 497919823, AN 1103/4-1 to DA and Research Unit 5429/1, project number: 467143400, FL 379/34-1, FL 379/35-1 to AF, ME 3161/5-1, ME 3161/6-1 to MM, AN 1103/5-1 to DA, and project number: 539593253, AN 1103/6-1 to DA). CTLL is a PhD fellow funded by the “Deutscher Akademischer Austauschdienst” (DAAD), German Academic Exchange Service, scholarship: 91828451).

The funders had no role in the design and conduct of the study, data collection, management, analyses, and interpretation, manuscript preparation, review, or approval, or decision to submit the manuscript for publication.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

We would like to thank Daniel Schulze for the helpful methodical exchange.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.brs.2025.06.005>.

References

- [1] Baltes PB, Staudinger UM, Lindenberger U. Lifespan PSYCHOLOGY: theory and application to intellectual functioning. *Annu Rev Psychol* 1999;50:471–507. <https://doi.org/10.1146/annurev.psych.50.1.471>.
- [2] Lövdén M, Fratiglioni L, Glymour MM, Lindenberger U, Tucker-Drob EM. Education and cognitive functioning across the life span. *Psychol Sci Public Interest* 2020;21:6–41. <https://doi.org/10.1177/1529100620920576>.
- [3] Perceval G, Flöel A, Meinzer M. Can transcranial direct current stimulation counteract age-associated functional impairment? *Neurosci Biobehav Rev* 2016;65: 157–72. <https://doi.org/10.1016/j.neubiorev.2016.03.028>.
- [4] Fjell AM, Walhovd KB. Structural brain changes in aging: courses, causes and cognitive consequences. *Rev Neurosci* 2010;21. <https://doi.org/10.1515/REVNEURO.2010.21.3.187>.
- [5] Goldthorpe RA, Rapley JM, Violante IR. A systematic review of non-invasive brain stimulation applications to memory in healthy aging. *Front Neurol* 2020;11: 575075. <https://doi.org/10.3389/fneur.2020.575075>.
- [6] Baharlouei H, Saba MA, Shaterzadeh Yazdi MJ, Jaberzadeh S. The effect of transcranial direct current stimulation on balance in healthy young and older adults: a systematic review of the literature. *Neurophysiol Clin* 2020;50:119–31. <https://doi.org/10.1016/j.neucli.2020.01.006>.
- [7] Cuevas-Trisan R. Balance problems and fall risks in the elderly. *Phys Med Rehabil Clin N Am* 2017;28:727–37. <https://doi.org/10.1016/j.pmr.2017.06.006>.
- [8] Antal A, Alekseichuk I, Bikson M, Brockmüller J, Brunoni AR, Chen R, et al. Low intensity transcranial electric stimulation: safety, ethical, legal regulatory and application guidelines. *Clin Neurophysiol* 2017;128:1774–809. <https://doi.org/10.1016/j.clinph.2017.06.001>.

- [9] Antal A, Luber B, Brem A-K, Bikson M, Brunoni AR, Cohen Kadosh R, et al. Non-invasive brain stimulation and neuroenhancement. *Clin Neurophysiol Pract* 2022; 7:146–65. <https://doi.org/10.1016/j.cnp.2022.05.002>.
- [10] Nitsche MA, Paulus W. Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *J Physiol* 2000;527:633–9. <https://doi.org/10.1111/j.1469-7793.2000.t01-1-00633.x>.
- [11] Stagg CJ, Antal A, Nitsche MA. Physiology of transcranial direct current stimulation. *J ECT* 2018;34:144–52. <https://doi.org/10.1097/YCT.0000000000000510>.
- [12] Antonenko D, Hayek D, Netzband J, Grittnier U, Flöel A. tDCS-induced episodic memory enhancement and its association with functional network coupling in older adults. *Sci Rep* 2019;9:2273. <https://doi.org/10.1038/s41598-019-38630-7>.
- [13] Barbati SA, Podda MV, Grassi C. Tuning brain networks: the emerging role of transcranial direct current stimulation on structural plasticity. *Front Cell Neurosci* 2022;16. <https://doi.org/10.3389/fncel.2022.945777>.
- [14] Kronberg G, Bridi M, Abel T, Bikson M, Parra LC. Direct current stimulation modulates LTP and LTD: activity dependence and dendritic effects. *Brain Stimulat* 2017;10:51–8. <https://doi.org/10.1016/j.brs.2016.10.001>.
- [15] Paciello F, Podda MV, Rolesi R, Cocco S, Petrosini L, Troiani D, et al. Anodal transcranial direct current stimulation affects auditory cortex plasticity in normal-hearing and noise-exposed rats. *Brain Stimulat* 2018;11:1008–23. <https://doi.org/10.1016/j.brs.2018.05.017>.
- [16] Meinzer M, Lindenberg R, Antonenko D, Flaisch T, Flöel A. Anodal transcranial direct current stimulation temporarily reverses age-associated cognitive decline and functional brain activity changes. *J Neurosci* 2013;33:12470–8. <https://doi.org/10.1523/JNEUROSCI.5743-12.2013>.
- [17] Pino-Esteban A, Megía-García Á, Martín-Caro Álvarez D, Beltran-Alacreu H, Avendaño-Coy J, Gómez-Soriano J, et al. Can transcranial direct current stimulation enhance functionality in older adults? A systematic review. *J Clin Med* 2021;10:2981. <https://doi.org/10.3390/jcm10132981>.
- [18] Panouillères MTN, Joundri RA, Brittian J, Jenkinson N. Reversing motor adaptation deficits in the ageing brain using non-invasive stimulation. *J Physiol* 2015;593: 3645–55. <https://doi.org/10.1113/JP270484>.
- [19] Zimmerman M, Nitsch M, Giraux P, Gerloff C, Cohen LG, Hummel FC. Neuroenhancement of the ageing brain: restoring skill acquisition in old subjects. *Ann Neurol* 2013;73:10–5. <https://doi.org/10.1002/ana.23761>.
- [20] Galli G, Vadillo MA, Sirota M, Feurra M, Medvedeva A. A systematic review and meta-analysis of the effects of transcranial direct current stimulation (tDCS) on episodic memory. *Brain Stimulat* 2019;12:231–41. <https://doi.org/10.1016/j.brs.2018.11.008>.
- [21] Siegert A, Diedrich L, Antal A. New methods, old brains—A systematic review on the effects of tDCS on the cognition of elderly people. *Front Hum Neurosci* 2021; 15:730134. <https://doi.org/10.3389/fnhum.2021.730134>.
- [22] Emonson MRL, Fitzgerald PB, Rogasch NC, Hoy KE. Neurobiological effects of transcranial direct current stimulation in younger adults, older adults and mild cognitive impairment. *Neuropsychologia* 2019;125:51–61. <https://doi.org/10.1016/j.neuropsychologia.2019.01.003>.
- [23] Habich A, Fehér KD, Antonenko D, Boraxbekk C-J, Flöel A, Nissen C, et al. Stimulating aged brains with transcranial direct current stimulation: opportunities and challenges. *Psychiatry Res Neuroimaging* 2020;306:111179. <https://doi.org/10.1016/j.pscychres.2020.111179>.
- [24] Kaminski E, Engelhardt M, Hoff M, Steele C, Villringer null A, Ragert P. tDCS effects on pointing task learning in young and old adults. *Sci Rep* 2021;11:3421. <https://doi.org/10.1038/s41598-021-82275-4>.
- [25] Hsu Ku Y, Zanto TP, Gazzaley A. Effects of noninvasive brain stimulation on cognitive function in healthy aging and Alzheimer's disease: a systematic review and meta-analysis. *Neurobiol Aging* 2015;36:2348–59. <https://doi.org/10.1016/j.neurobiolaging.2015.04.016>.
- [26] Indahlastari A, Hardcastle C, Albizu A, Alvarez-Alvarado S, Boutzoukas EM, Evangelista ND, et al. A systematic review and meta-analysis of transcranial direct current stimulation to remediate age-related cognitive decline in healthy older adults. *Neuropsychiatr Dis Treat* 2021;17:971–90. <https://doi.org/10.2147/NDT.S259499>.
- [27] Bagattini C, Cid-Fernández S, Bulgari M, Miniussi C, Bortoletto M. Opposite pattern of transcranial direct current stimulation effects in middle-aged and older adults: behavioral and neurophysiological evidence. *Front Aging Neurosci* 2023;15: 1087749. <https://doi.org/10.3389/fnagi.2023.1087749>.
- [28] Hsu T-Y, Juan C-H, Tseng P. Individual differences and state-dependent responses in transcranial direct current stimulation. *Front Hum Neurosci* 2016;10. <https://doi.org/10.3389/fnhum.2016.00643>.
- [29] Perceval G, Martin AK, Copland DA, Laine M, Meinzer M. Multisession transcranial direct current stimulation facilitates verbal learning and memory consolidation in young and older adults. *Brain Lang* 2020;205:104788. <https://doi.org/10.1016/j.bandl.2020.104788>.
- [30] De Lange A-MG, Bråthen ACS, Rohani DA, Grydeland H, Fjell AM, Walhovd KB. The effects of memory training on behavioral and microstructural plasticity in young and older adults. *Hum Brain Mapp* 2017;38:5666–80. <https://doi.org/10.1002/hbm.23756>.
- [31] Nitsche MA, Bikson M, Bestmann S. On the use of meta-analysis in neuromodulatory non-invasive brain stimulation. *Brain Stimulat* 2015;8:666–7. <https://doi.org/10.1016/j.brs.2015.03.008>.
- [32] Polanía R, Nitsche MA, Ruff CC. Studying and modifying brain function with non-invasive brain stimulation. *Nat Neurosci* 2018;21:174–87. <https://doi.org/10.1038/s41593-017-0054-4>.
- [33] Vergallito A, Feroldi S, Pisoni A, Romero Lauro LJ. Inter-individual variability in tDCS effects: a narrative review on the contribution of stable, variable, and contextual factors. *Brain Sci* 2022;12:522. <https://doi.org/10.3390/brainsci12050522>.
- [34] Wansbrough K, Tan J, Vallence A-M, Fujiyama H. Recent advancements in optimising transcranial electrical stimulation: reducing response variability through individualised stimulation. *Curr Opin Behav Sci* 2024;56:101360. <https://doi.org/10.1016/j.cobeha.2024.101360>.
- [35] Antonenko D, Grittnier U, Saturnino G, Nierhaus T, Thielscher A, Flöel A. Inter-individual and age-dependent variability in simulated electric fields induced by conventional transcranial electrical stimulation. *Neuroimage* 2021;224:117413. <https://doi.org/10.1016/j.neuroimage.2020.117413>.
- [36] Indahlastari A, Albizu A, O'Shea A, Forbes MA, Nissim NR, Kraft JN, et al. Modeling transcranial electrical stimulation in the aging brain. *Brain Stimulat* 2020;13:664–74. <https://doi.org/10.1016/j.brs.2020.02.007>.
- [37] Antal A, Keeser D, Priori A, Padberg F, Nitsche MA. Conceptual and procedural shortcomings of the systematic review “Evidence That Transcranial Direct Current Stimulation (tDCS) Generates Little-to-no Reliable Neurophysiologic Effect Beyond MEP Amplitude Modulation in Healthy Human Subjects: a Systematic Review” by horvath and Co-workers. *Brain Stimulat* 2015;8:846–9. <https://doi.org/10.1016/j.brs.2015.05.010>.
- [38] Wiethoff S, Hamada M, Rothwell JC. Variability in response to transcranial direct current stimulation of the motor cortex. *Brain Stimulat* 2014;7:468–75. <https://doi.org/10.1016/j.brs.2014.02.003>.
- [39] McCutcheon RA, Pillinger T, Mizuno Y, Montgomery A, Pandian H, Vano L, et al. The efficacy and heterogeneity of antipsychotic response in schizophrenia: a meta-analysis. *Mol Psychiatry* 2021;26:1310–20. <https://doi.org/10.1038/s41380-019-0502-5>.
- [40] Merritt K, McCutcheon RA, Aleman A, Ashley S, Beck K, Block W, et al. Variability and magnitude of brain glutamate levels in schizophrenia: a Meta and mega-analysis. *Mol Psychiatry* 2023;28:2039–48. <https://doi.org/10.1038/s41380-023-01991-7>.
- [41] Homan S, Muscat W, Joanlanne A, Marousis N, Cecere G, Hofmann L, et al. Treatment effect variability in brain stimulation across psychiatric disorders: a meta-analysis of variance. *Neurosci Biobehav Rev* 2021;124:54–62. <https://doi.org/10.1016/j.neubiorev.2020.11.033>.
- [42] Mizuno Y, McCutcheon RA, Brugger SP, Howes OD. Heterogeneity and efficacy of antipsychotic treatment for schizophrenia with or without treatment resistance: a meta-analysis. *Neuropsychopharmacology* 2020;45:622–31. <https://doi.org/10.1038/s41386-019-0577-3>.
- [43] Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;n71. <https://doi.org/10.1136/bmj.n71>.
- [44] Nakagawa S, Poulain R, Mengersen K, Reinhold K, Engqvist L, Lagisz M, et al. Meta-analysis of variation: ecological and evolutionary applications and beyond. *Methods Ecol Evol* 2015;6:143–52. <https://doi.org/10.1111/2041-210X.12309>.
- [45] Senior AM, Viechtbauer W, Nakagawa S. Revisiting and expanding the meta-analysis of variation: the log coefficient of variation ratio. *Res Synth Methods* 2020; 11:553–67. <https://doi.org/10.1002/rsrm.1423>.
- [46] Pick JL, Nakagawa S, Noble DWA. Reproducible, flexible and high-throughput data extraction from primary literature: the metaDigitiser R package. *Methods Ecol Evol* 2019;10:426–31. <https://doi.org/10.1111/2041-210X.13118>.
- [47] Higgins JPT. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557–60. <https://doi.org/10.1136/bmj.327.7414.557>.
- [48] Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58. <https://doi.org/10.1002/sim.1186>.
- [49] Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw* 2010;36. <https://doi.org/10.18637/jss.v036.i03>.
- [50] Chen PL, Machado L. Developing clinically practical transcranial direct current stimulation protocols to improve saccadic eye movement control. *J Eye Mov Res* 2017;10. <https://doi.org/10.16910/jemr.10.3.5>.
- [51] Conley AC, Fulham WR, Marquez JL, Parsons MW, Karayianidis F. No effect of anodal transcranial direct current stimulation over the motor cortex on response-related ERPs during a conflict task. *Front Hum Neurosci* 2016;10. <https://doi.org/10.3389/fnhum.2016.00384>.
- [52] Craig CE, Doumas M. Anodal transcranial direct current stimulation shows minimal, measure-specific effects on dynamic postural control in young and older adults: a double blind, sham-controlled study. *PLoS One* 2017;12:e0170331. <https://doi.org/10.1371/journal.pone.0170331>.
- [53] Fujiyama H, Tan J, Puri R, Hinder MR. Influence of tDCS over right inferior frontal gyrus and pre-supplementary motor area on perceptual decision-making and response inhibition: a healthy ageing perspective. *Neurobiol Aging* 2022;109: 11–21. <https://doi.org/10.1016/j.neurobiolaging.2021.09.014>.
- [54] Goodwill AM, Daly RM, Kidgell DJ. The effects of anodal-tDCS on cross-limb transfer in older adults. *Clin Neurophysiol Off J Int Fed Clin Neurophysiol* 2015; 126:2189–97. <https://doi.org/10.1016/j.clinph.2015.01.006>.
- [55] Muffel T, Kirsch F, Shih P-C, Kallach B, Schaumberg S, Villringer A, et al. Anodal transcranial direct current stimulation over S1 differentially modulates proprioceptive accuracy in young and old adults. *Front Aging Neurosci* 2019;11: 264. <https://doi.org/10.3389/fnagi.2019.00264>.
- [56] Arciniegua H, Gözenman F, Jones KT, Stephens JA, Berryhill ME. Frontoparietal tDCS benefits visual working memory in older adults with low working memory capacity. *Front Aging Neurosci* 2018;10:57. <https://doi.org/10.3389/fnagi.2018.00057>.
- [57] Fertonani A, Brambilla M, Cotelli M, Miniussi C. The timing of cognitive plasticity in physiological aging: a tDCS study of naming. *Front Aging Neurosci* 2014;6. <https://doi.org/10.3389/fnagi.2014.00131>.

- [58] Fiori V, Nitsche M, Iasevoli L, Cucuzza G, Caltagirone C, Marangolo P. Differential effects of bihemispheric and unihemispheric transcranial direct current stimulation in young and elderly adults in verbal learning. *Behav Brain Res* 2017;321:170–5. <https://doi.org/10.1016/j.bbr.2016.12.044>.
- [59] Leach RC, McCurdy MP, Trumbo MC, Matzen LE, Leshikar ED. Differential age effects of transcranial direct current stimulation on associative memory. *J Gerontol B Psychol Sci Soc Sci* 2019;74:1163–73. <https://doi.org/10.1093/geronb/gby003>.
- [60] Manenti R, Brambilla M, Petesi M, Ferrari C, Cotelli M. Enhancing verbal episodic memory in older and young subjects after non-invasive brain stimulation. *Front Aging Neurosci* 2013;5. <https://doi.org/10.3389/fnagi.2013.00049>.
- [61] Martin AK, Meinzer M, Lindenberg R, Sieg MM, Nachtigall L, Flöel A. Effects of transcranial direct current stimulation on neural networks in young and older adults. *J Cogn Neurosci* 2017;29:1817–28. https://doi.org/10.1162/jocn_a.01166.
- [62] Meinzer M, Antonenko D, Lindenberg R, Hetzer S, Ulm L, Avirame K, et al. Electrical brain stimulation improves cognitive performance by modulating functional connectivity and task-specific activation. *J Neurosci* 2012;32:1859–66. <https://doi.org/10.1523/JNEUROSCI.4812-11.2012>.
- [63] Orcioli-Silva D, Islam A, Baker MR, Gobbi LTB, Rochester L, Pantall A. Bi-Anodal transcranial direct current stimulation combined with treadmill walking decreases motor cortical activity in young and older adults. *Front Aging Neurosci* 2021;13:739998. <https://doi.org/10.3389/fnagi.2021.739998>.
- [64] Peter J, Neumann-Dunayevska E, Geugelin F, Ninosu N, Plewnia C, Klöppel S. Reducing negative affect with anodal transcranial direct current stimulation increases memory performance in young-but not in elderly-individuals. *Brain Struct Funct* 2019;224:2973–82. <https://doi.org/10.1007/s00429-019-01946-1>.
- [65] Ross LA, McCoy D, Coslett HB, Olson IR, Wolk DA. Improved proper name recall in aging after electrical stimulation of the anterior temporal lobes. *Front Aging Neurosci* 2011;3. <https://doi.org/10.3389/fnagi.2011.00016>.
- [66] Habich A, Slotboom J, Peter J, Wiest R, Klöppel S. No effect of anodal tDCS on verbal episodic memory performance and neurotransmitter levels in young and elderly participants. *Neural Plast* 2020;2020:8896791. <https://doi.org/10.1155/2020/8896791>.
- [67] Lindberg PG, Verneau M, Boterff QL, Cuenca-Maia M, Baron J-C, Maier MA. Age- and task-dependent effects of cerebellar tDCS on manual dexterity and motor learning-A preliminary study. *Neurophysiol Clin Clin Neurophysiol* 2022;52:354–65. <https://doi.org/10.1016/j.neucli.2022.07.006>.
- [68] Lasogga L, Gramegna C, Müller D, Habel U, Mehler DMA, Gur RC, et al. Meta-analysis of variance in tDCS effects on response inhibition. *Sci Rep* 2024;14:19197. <https://doi.org/10.1038/s41598-024-70065-7>.
- [69] Bhattacharjee S, Kashyap R, Goodwill AM, O'Brien BA, Rapp B, Oishi K, et al. Sex difference in tDCS current mediated by changes in cortical anatomy: a study across young, middle and older adults. *Brain Stimulat* 2022;15:125–40. <https://doi.org/10.1016/j.brs.2021.11.018>.
- [70] Fertonani A, Miniussi C. Transcranial electrical stimulation: what we know and do not know about mechanisms. *Neuroscientist* 2017;23:109–23. <https://doi.org/10.1177/1073858416631966>.
- [71] Stagg CJ. Magnetic resonance spectroscopy as a tool to study the role of GABA in motor-cortical plasticity. *Neuroimage* 2014;86:19–27. <https://doi.org/10.1016/j.neuroimage.2013.01.009>.
- [72] Evans C, Zich C, Lee JSA, Ward N, Bestmann S. Inter-individual variability in current direction for common tDCS montages. *Neuroimage* 2022;260:119501. <https://doi.org/10.1016/j.neuroimage.2022.119501>.
- [73] Roheger M, Mäder A, Riemann S, Niemann F, Kessler K, Martin AK, et al. Intact embodiment during perspective-taking in older adults is not affected by focal tDCS. *GeroScience* 2025. <https://doi.org/10.1007/s11357-025-01554-4>.
- [74] Saturnino GB, Antunes A, Thielscher A. On the importance of electrode parameters for shaping electric field patterns generated by tDCS. *Neuroimage* 2015;120:25–35. <https://doi.org/10.1016/j.neuroimage.2015.06.067>.
- [75] Thielscher A, Antunes A, Saturnino GB. Field modeling for transcranial magnetic stimulation: a useful tool to understand the physiological effects of TMS? 2015 37th annu. Int Conf IEEE Eng Med Biol Soc EMBC 2015:222–5. <https://doi.org/10.1109/EMBC.2015.7318340>. Milan: IEEE.
- [76] Cabeza R. Hemispheric asymmetry reduction in older adults: the HAROLD model. *Psychol Aging* 2002;17:85–100. <https://doi.org/10.1037/0882-7974.17.1.85>.
- [77] Reuter-Lorenz PA, Cappel KA. Neurocognitive aging and the compensation hypothesis. *Curr Dir Psychol Sci* 2008;17:177–82. <https://doi.org/10.1111/j.1467-8721.2008.00570.x>.
- [78] Stern Y, Barnes CA, Grady C, Jones RN, Raz N. Brain reserve, cognitive reserve, compensation, and maintenance: operationalization, validity, and mechanisms of cognitive resilience. *Neurobiol Aging* 2019;83:124–9. <https://doi.org/10.1016/j.neurobiolaging.2019.03.022>.
- [79] Lee JSA, Bestmann S, Evans C. A future of current flow modelling for transcranial electrical stimulation? *Curr Behav Neurosci Rep* 2021;8:150–9. <https://doi.org/10.1007/s40473-021-00238-5>.
- [80] Cabeza R, Albert M, Belleville S, Craik FIM, Duarte A, Grady CL, et al. Maintenance, reserve and compensation: the cognitive neuroscience of healthy ageing. *Nat Rev Neurosci* 2018;19:701–10. <https://doi.org/10.1038/s41583-018-0068-2>.
- [81] Antonenko D, Nierhaus T, Meinzer M, Prehn K, Thielscher A, Ittermann B, et al. Age-dependent effects of brain stimulation on network centrality. *Neuroimage* 2018;176:71–82. <https://doi.org/10.1016/j.neuroimage.2018.04.038>.
- [82] Meinzer M, Shahbabaie A, Antonenko D, Blankenburg F, Fischer R, Hartwigsen G, et al. Investigating the neural mechanisms of transcranial direct current stimulation effects on human cognition: current issues and potential solutions. *Front Neurosci* 2024;18:1389651. <https://doi.org/10.3389/fnins.2024.1389651>.
- [83] Menardi A, Momi D, Vallesi A, Barabási A-L, Towson EK, Santarnecchi E. Maximizing brain networks engagement via individualized connectome-wide target search. *Brain Stimulat* 2022;15:1418–31. <https://doi.org/10.1016/j.brs.2022.09.011>.
- [84] Wang JX, Rogers LM, Gross EZ, Ryals AJ, Dokucu ME, Brandstatt KL, et al. Targeted enhancement of cortical-hippocampal brain networks and associative memory. *Science* 2014;345:1054–7. <https://doi.org/10.1126/science.1252900>.
- [85] Antonenko D, Külzow N, Sousa A, Prehn K, Grittner U, Flöel A. Neuronal and behavioral effects of multi-day brain stimulation and memory training. *Neurobiol Aging* 2018;61:245–54. <https://doi.org/10.1016/j.neurobiolaging.2017.09.017>.
- [86] Fertonani A, Pirulli C, Bollini A, Miniussi C, Bortoletto M. Age-related changes in cortical connectivity influence the neuromodulatory effects of transcranial electrical stimulation. *Neurobiol Aging* 2019;82:77–87. <https://doi.org/10.1016/j.neurobiolaging.2019.07.009>.
- [87] Fromm AE, Grittner U, Brodt S, Flöel A, Antonenko D. No object–location memory improvement through focal transcranial direct current stimulation over the right temporoparietal cortex. *Life* 2024;14:539. <https://doi.org/10.3390/life14050539>.
- [88] Schutter DJLG, Smits F, Klaus J. Mind matters: a narrative review on affective state-dependency in non-invasive brain stimulation. *Int J Clin Health Psychol* 2023;23:100378. <https://doi.org/10.1016/j.ijchp.2023.100378>.
- [89] Majdi A, Van Boekholdt L, Sadigh-Eteghad S, Mc Laughlin M. A systematic review and meta-analysis of transcranial direct-current stimulation effects on cognitive function in patients with Alzheimer's disease. *Mol Psychiatry* 2022;27:2000–9. <https://doi.org/10.1038/s41380-022-01444-7>.
- [90] Nissim NR, O'Shea A, Indahlstari A, Telles R, Richards L, Porges E, et al. Effects of in-Scanner bilateral frontal tDCS on functional connectivity of the working memory network in older adults. *Front Aging Neurosci* 2019;11:51. <https://doi.org/10.3389/fnagi.2019.00051>.
- [91] Narmashiri A, Akbari F. The effects of transcranial direct current stimulation (tDCS) on the cognitive functions: a systematic review and meta-analysis. *Neuropsychol Rev* 2023. <https://doi.org/10.1007/s11065-023-09627-x>.
- [92] Rudroff T, Workman CD, Fietsam AC, Kamholz J. Response variability in transcranial direct current stimulation: why sex matters. *Front Psychiatry* 2020;11:585. <https://doi.org/10.3389/fpsy.2020.00585>.
- [93] Evans C, Bachmann C, Lee JSA, Gregoriou E, Ward N, Bestmann S. Dose-controlled tDCS reduces electric field intensity variability at a cortical target site. *Brain Stimulat* 2020;13:125–36. <https://doi.org/10.1016/j.brs.2019.10.004>.
- [94] Niemann F, Shababaie A, Paßmann S, Riemann S, Malinowski R, Kocatay H, et al. Neuronavigated focalized transcranial direct current stimulation administered during functional magnetic resonance imaging. *J Vis Exp* 2024:67155. <https://doi.org/10.3791/67155>.
- [95] Rasmussen ID, Boayue NM, Mittner M, Bystad M, Grønli OK, Vangberg TR, et al. High-definition transcranial direct current stimulation improves delayed memory in alzheimer's disease patients: a pilot study using computational modeling to optimize electrode position. *J Alzheimers Dis* 2021;83:753–69. <https://doi.org/10.3233/JAD-210378>.