

7T MRS Glutamate and GABA in Alzheimer's Disease

Laura Göschel^{1,2} | Andrea Dell'Orco^{1,3,4} | Bernd Ittermann⁵ |
Charlotte E. Teunissen⁶ | Patty Hoede⁷ | Jeanette Melin⁸ | Leslie Pendrill⁸ |
Sebastian Roemer-Cassiano^{9,10,11} | Nicolai Franzmeier^{12,13,14} | Peter Koertvelyessy^{15,16} |
Agnes Flöel^{17,18} | Ariane Fillmer⁵

¹Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Department of Neurology, Charitéplatz 1, Berlin, Germany

²Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Neuroscience Clinical Research Center, Charitéplatz 1, Berlin, Germany

³Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Neuroscience Clinical Research Center, Charitéplatz 1, Berlin, Germany

⁴Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Department of Neuroradiology, Charitéplatz 1, Berlin, Germany

⁵Physikalisch-Technische Bundesanstalt (PTB), Braunschweig and Berlin, Germany

⁶Neurochemistry Laboratory, Department of Laboratory Medicine, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, Amsterdam, Netherlands

⁷Neurochemistry Laboratory, Department of Laboratory Medicine, Vrije Universiteit Amsterdam, Amsterdam UMC, Amsterdam, Netherlands, Amsterdam, Netherlands

⁸RISE, Research Institutes of Sweden, Division Safety and Transport, Measurement Science and Technology, Göteborg, Sweden

⁹Department of Neurology, University Hospital, LMU Munich, Munich, Bavaria, Germany

¹⁰Max Planck School of Cognition, Leipzig, Sachsen, Germany

Abstract

Background: Functional neuroimaging studies suggest a dynamic trajectory in Alzheimer's disease (AD), with early hyperconnectivity followed by hypoconnectivity due to pathologic progression. This study aimed to investigate this hypothesis using neurochemical markers derived from high-field magnetic resonance spectroscopy (MRS).

Method: We analyzed data from 126 older adults enrolled in the NeuroMET studies, spanning from normal aging to dementia due to suspected AD. Using ultra-high-field 7 Tesla MRS, we quantified levels of the neurotransmitters glutamate (excitatory) and GABA (inhibitory) in the precentral cortex. Plasma *p*-Tau181 was used as a proxy for AD pathology, and memory was assessed using the NeuroMET Memory Metric (NMM). A *p*-Tau181 threshold of >2.08 pg/mL was applied as an estimated marker of amyloid positivity (Aβ+). Linear mixed-effects models, adjusted for age at baseline, were used to evaluate associations and interaction effects.

Result: Glutamate levels showed a non-linear association with age, decreasing up to around 70 years and increasing thereafter, while GABA levels remained stable (Figure 1A–C). Before reaching the suggested pathological conversion threshold for amyloid positivity, both glutamate and GABA showed a slight, non-significant increase in individuals with higher *p*-Tau181 levels (Figure 1D–F). Notably, amyloid status moderated the relationship between memory ability and glutamate levels (Figure 1G–I). Among Aβ-negative individuals, lower memory ability was associated with elevated glutamate, potentially reflecting a very early compensatory response.

Conclusion: Our findings support the hypothesis of a non-linear neurochemical trajectory in aging and early AD pathology. The observed age-dependent fluctuations in glutamate, together with subtle shifts in response to rising plasma *p*-Tau181, may reflect early pathologic or compensatory mechanisms. The interaction between

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Alzheimer's Association. *Alzheimer's & Dementia* published by Wiley Periodicals LLC on behalf of Alzheimer's Association.

¹¹Institute for Stroke and Dementia Research (ISD), University Hospital, LMU Munich, Munich, Bavaria, Germany

¹²University of Gothenburg, The Sahlgrenska Academy, Institute of Neuroscience and Physiology, Psychiatry and Neurochemistry, Gothenburg, Sweden

¹³Institute for Stroke and Dementia Research (ISD), LMU University Hospital, LMU Munich, Munich, Germany

¹⁴Munich Cluster for Systems Neurology (SyNergy), Munich, Bavaria, Germany

¹⁵Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Department of Neurology, Charitéplatz 1, Berlin, Berlin, Germany

¹⁶German Center for Neurodegenerative Diseases (DZNE), Magdeburg, Germany

¹⁷University Medicine Greifswald, Greifswald, Mecklenburg-Vorpommern, Germany

¹⁸German Centre for Neurodegenerative Diseases (DZNE), Standort Rostock/Greifswald, Greifswald, Germany

Correspondence

Laura Göschel, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Department of Neurology, Charitéplatz 1, Berlin, Germany.

Email: laura.goeschel@charite.de

glutamate and memory ability, particularly in supposedly A β -negative individuals, suggests that excitatory neurotransmission could transiently support cognitive function in the face of emerging pathology. Notably, the threshold of plasma *p*-Tau181 used to define amyloid positivity may identify individuals in progressed stages, potentially missing earlier windows of therapeutic opportunity. As potential treatments may have markedly different effects depending on the stage of disease progression, investigating the trajectory of neuronal connectivity and activity is critical.

Table 1. Linear mixed-effects models assessing associations between plasma p-Tau181 levels and glutamate, GABA, and the glutamate/GABA ratio across time, while adjusting for age (at baseline) and accounting for random intercepts per subject. Significant associations (p < 0.05) are bolded.

Model Glutamate			Model GABA			Model Glutamate/GABA		
lmer(glu ~ ns(pTau, df = 3) * ns(year, df = 3) + ns(age_bl, df = 2) + (1 record_id))			lmer(gaba ~ ns(pTau, df = 3) * ns(year, df = 4) + ns(age_bl, df = 1) + (1 record_id))			lmer(glu_gaba ~ ns(pTau, df = 3) * ns(year, df = 3) + ns(age_bl, df = 2) + (1 record_id))		
age_bl	β [95% CI]	p	age_bl	β [95% CI]	p	age_bl	β [95% CI]	p
60	-0,12 [-0,20; -0,03]	0,006	60	-0,01 [-0,03; 0,01]	0,201	60	-0,02 [-0,06; 0,01]	0,192
65	-0,08 [-0,14; -0,02]	0,012				65	-0,01 [-0,04; 0,01]	0,331
70	-0,01 [-0,05; 0,02]	0,435				70	0,01 [-0,01; 0,02]	0,510
75	0,06 [0,00; 0,11]	0,046				75	0,03 [0,00; 0,05]	0,042
80	0,10 [0,02; 0,18]	0,016				80	0,04 [0,00; 0,07]	0,037
p-Tau181			p-Tau181			p-Tau181		
1,25	0,42 [-0,90; 1,74]	0,535	1,25	0,54 [-0,02; 1,11]	0,059	1,25	-0,41 [-1,01; 0,19]	0,183
1,88	-0,10 [-0,50; 0,30]	0,612	1,88	0,12 [-0,05; 0,30]	0,156	1,88	-0,16 [-0,34; 0,02]	0,082
2,5	-0,35 [-0,86; 0,15]	0,166	2,5	-0,10 [-0,32; 0,12]	0,359	2,5	-0,02 [-0,25; 0,21]	0,877
5	-0,06 [-0,28; 0,15]	0,560	5	-0,01 [-0,11; 0,08]	0,786	5	-0,01 [-0,11; 0,09]	0,864
Model Glutamate			Model GABA			Model Glutamate/GABA		
lmer(glu ~ pTau_group * ns(nmm, df = 4) + ns(year, df = 3) + ns(age_bl, df = 2) + (1 record_id))			lmer(gaba ~ pTau_group * ns(nmm, df = 3) + ns(year, df = 3) + ns(age_bl, df = 2) + (1 record_id))			lmer(glu_gaba ~ pTau_group * ns(nmm, df = 3) + ns(year, df = 3) + ns(age_bl, df = 2) + (1 record_id))		
Aβ-			Aβ-			Aβ-		
NMM	β [95% CI]	p	NMM	β [95% CI]	p	NMM	β [95% CI]	p
-2.0	1,52 [0,17; 2,88]	0,028	-2.0	0,10 [-0,45; 0,64]	0,728	-2.0	0,03 [-0,53; 0,59]	0,915
-1.0	-0,39 [-1,05; 0,26]	0,239	-1.0	0,01 [-0,28; 0,31]	0,921	-1.0	-0,02 [-0,32; 0,27]	0,876
-0.5	-1,57 [-2,52; -0,62]	0,001	-0.5	-0,04 [-0,26; 0,19]	0,758	-0.5	-0,06 [-0,29; 0,17]	0,630
0.5	0,15 [-0,73; 1,02]	0,745	0.5	-0,10 [-0,40; 0,19]	0,491	0.5	-0,09 [-0,40; 0,22]	0,580
1.0	0,49 [-0,29; 1,27]	0,214	1.0	-0,07 [-0,26; 0,12]	0,454	1.0	-0,04 [-0,25; 0,17]	0,708
2.0	-0,56 [-1,59; 0,46]	0,278	2.0	-0,03 [-0,40; 0,35]	0,891	2.0	0,03 [-0,40; 0,46]	0,886
Aβ+			Aβ+			Aβ+		
NMM	β [95% CI]	p	NMM	β [95% CI]	p	NMM	β [95% CI]	p
-2.0	0,31 [-0,08; 0,7]	0,122	-2.0	0,17 [-0,02; 0,35]	0,075	-2.0	-0,09 [-0,28; 0,11]	0,379
-1.0	0,07 [-0,29; 0,44]	0,689	-1.0	0,10 [-0,03; 0,23]	0,143	-1.0	-0,06 [-0,20; 0,07]	0,342
-0.5	-0,07 [-0,63; 0,49]	0,814	-0.5	0,05 [-0,12; 0,23]	0,549	-0.5	-0,05 [-0,23; 0,13]	0,584
0.5	0,25 [-0,94; 1,45]	0,675	0.5	0,01 [-0,20; 0,22]	0,914	0.5	-0,04 [-0,26; 0,17]	0,693
1.0	0,00 [-0,90; 0,90]	0,998	1.0	0,07 [-0,31; 0,45]	0,715	1.0	-0,07 [-0,49; 0,34]	0,731
2.0	-0,80 [-3,85; 2,25]	0,606	2.0	0,16 [-0,72; 1,03]	0,723	2.0	-0,12 [-1,08; 0,84]	0,811

Abbreviations: Aβ = Amyloid-beta; CI = Confidence Interval; GABA = Gamma-Aminobutyric Acid; Glu = Glutamate; nmm = NeuroMET Memory Metric; p-Tau = Phosphorylated Tau.




Figure 1: Association between neurotransmitter levels and age, plasma pTau181 and memory ability indicated by the NMM. (A–C) Metabolite levels across age for glutamate (A), GABA (B), and glutamate/GABA ratio (C), based on linear mixed-effects models controlling for pTau181, and weighted by metabolite-specific measures of uncertainties (CRLBs). **(D–F)** Predicted metabolite levels across plasma pTau181 concentrations for glutamate (D), GABA (E), and glutamate/GABA ratio (F), adjusted for age at baseline and weighted by metabolite-specific CRLBs. Dashed vertical lines indicate the previously established cut-off of 2.08 pg/mL for elevated pTau181. **(G–I)** Predicted metabolite levels across memory ability (NMM scores) stratified by Aβ status (estimated by plasma pTau181), again adjusted for age at baseline and weighted by metabolite-specific CRLBs. Shaded areas represent 95% confidence intervals; data points reflect individual measurements. **Abbreviations:** Aβ = amyloid-beta; CRLB = Cramér–Rao lower bound; GABA = γ -aminobutyric acid; NMM = NeuroMET Memory Metric.