

MOLECULAR AND CELL BIOLOGY

Cell-Type Specific Contributions to Metabolic Connectivity in an Alzheimer's Disease Mouse Model

Manvir Lalia¹ | Stephan Wagner¹ | Selina Hummel¹ | Justus Thevis¹ |
 Danilo Prtvar² | Artem Zatcepin¹ | Valerio Zenatti² | Laura Bartos¹ |
 Sabina Tahirovic² | Matthias Brendel^{1,2,3} | Johannes Gnörich^{1,2}

¹LMU University Hospital, Munich, Germany

²German Center for Neurodegenerative Diseases (DZNE), Munich, Germany

³Munich Cluster for Systems Neurology (SyNergy), Munich, Bavaria, Germany

Correspondence

Manvir Lalia, LMU University Hospital,
Munich, Germany.
Email: Manvir.Lalia@med.uni-muenchen.de

Abstract

Background: The integration of molecular imaging and multivariate connectivity approaches has emerged as a novel approach to gain insights into the underlying pathophysiology in neurodegenerative diseases. Metabolic connectivity, in particular, has already demonstrated disease-related pattern changes in both human and mammalian brains. However, the cellular sources of disconnected brain regions have not been investigated in detail. This study aimed to elucidate the driving cellular sources of metabolic connectivity in an Alzheimer's disease (AD) mouse model and wild-type mice (WT).

Method: After intravenous injection of 45MBq F-18-FDG, a static PET/MRI was performed on APP^{NL-G-F} and age- and sex-matched WT controls to obtain maps of regional FDG uptake and metabolic connectivity. To calculate the inter-regional correlations for metabolic connectivity, 26 delineated brain regions were used, resulting in a 26 × 26 matrix of Pearson's correlation coefficient pairs. Subsequently, the brain was extracted and separated into fore- and hindbrain to achieve region-specific isolation of microglia, astrocytes, oligodendrocytes, and neurons. The radioactivity of each cell fraction was measured to quantify the cell-specific FDG uptake (Figure 1D).

Result: APP^{NL-G-F} mice demonstrated higher FDG uptake compared to WT, along with a significantly increased metabolic connectivity between fore- and hindbrain (Figure 1A-C). Among all cell types, microglial exhibited the highest single-cell FDG uptake, in both mouse models (Figure 1E). In APP^{NL-G-F} mice, microglia, astrocytes, and oligodendrocytes displayed increased FDG uptake, while neurons exhibited reduced uptake compared to WT. The correlation between forebrain and hindbrain cellular FDG uptake was significant across all cell types in the APP^{NL-G-F} model (microglia $r=0.89$, $p=0.0006$; astrocytes $r=0.65$, $p=0.042$; oligodendrocytes $r=0.77$, $p=0.025$).

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Alzheimer's Association. *Alzheimer's & Dementia* published by Wiley Periodicals LLC on behalf of Alzheimer's Association.

and neurons $r=0.51$, $p = 0.005$). In contrast, WT mice did not exhibit any significant correlation in single-cell uptake between forebrain and hindbrain. Notably, region-specific microglial FDG uptake correlated significantly with respective FDG-PET signals in APP^{NL-G-F} mice (forebrain $r=0.89$, $p = 0.007$; hindbrain $r=0.8$, $p = 0.014$), whereas no significant correlation was observed for other cell types.

Conclusion: These findings suggest that microglia are the primary drivers of the increased forebrain-hindbrain metabolic connectivity observed in the AD mouse model. Further RNA expression analyses could provide valuable insights into the molecular mechanisms underlying microglial metabolic coupling in neurodegeneration.

Fig. 1

